

B3

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
23 August 2001 (23.08.2001)

PCT

(10) International Publication Number
WO 01/61009 A2(51) International Patent Classification⁷: C12N 15/12,
15/57, C07K 14/47, 14/555, 14/705, C12N 9/64, C12Q
1/68, A61K 38/17, 38/48, 48/00, G01N 33/53, 33/68

(21) International Application Number: PCT/US01/04828

(22) International Filing Date: 15 February 2001 (15.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/182,723	15 February 2000 (15.02.2000)	US
60/182,724	15 February 2000 (15.02.2000)	US
60/182,733	15 February 2000 (15.02.2000)	US
60/183,896	22 February 2000 (22.02.2000)	US
60/184,275	23 February 2000 (23.02.2000)	US
60/184,482	23 February 2000 (23.02.2000)	US
60/184,497	23 February 2000 (23.02.2000)	US
60/184,744	24 February 2000 (24.02.2000)	US
60/197,083	13 April 2000 (13.04.2000)	US
60/224,157	10 August 2000 (10.08.2000)	US
60/233,405	18 September 2000 (18.09.2000)	US
60/236,060	27 September 2000 (27.09.2000)	US
60/259,414	2 January 2001 (02.01.2001)	US
60/262,454	18 January 2001 (18.01.2001)	US
09/783,429	14 February 2001 (14.02.2001)	US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:

US	60/182,723 (CIP)
Filed on	15 February 2000 (15.02.2000)
US	60/182,724 (CIP)
Filed on	15 February 2000 (15.02.2000)
US	60/182,733 (CIP)
Filed on	15 February 2000 (15.02.2000)
US	60/183,896 (CIP)
Filed on	22 February 2000 (22.02.2000)
US	60/184,275 (CIP)
Filed on	23 February 2000 (23.02.2000)
US	60/184,482 (CIP)
Filed on	23 February 2000 (23.02.2000)
US	60/184,497 (CIP)
Filed on	23 February 2000 (23.02.2000)

US	60/184,744 (CIP)
Filed on	24 February 2000 (24.02.2000)
US	60/197,083 (CIP)
Filed on	13 April 2000 (13.04.2000)
US	60/224,157 (CIP)
Filed on	10 August 2000 (10.08.2000)
US	60/233,405 (CIP)
Filed on	18 September 2000 (18.09.2000)
US	60/236,060 (CIP)
Filed on	27 September 2000 (27.09.2000)
US	60/259,414 (CIP)
Filed on	2 January 2001 (02.01.2001)
US	60/262,454 (CIP)
Filed on	18 January 2001 (18.01.2001)
US	09/783,429 (CIP)
Filed on	14 February 2001 (14.02.2001)

(71) Applicant (for all designated States except US): CURA-
GEN CORPORATION [US/US]; 11th Floor, 555 Long
Wharf Drive, New Haven, CT 06511 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): MALYANKAR,
Uriel, M. [IN/US]; 35 Averill Place, Branford, CT 06405
(US). TCHERNEV, Velizar, T. [BG/US]; Apt. 156, 1216
SW 2nd Avenue, Gainesville, FL 32601 (US). PADIGARU,
Muralidhara [IN/US]; 1579 Rhinelander Avenue, Bronx,
NY 10461 (US). TAUPIER, Raymond, J., Jr. [US/US];
47 Holmes Street, East Haven, CT 06512 (US). SPYTEK,
Kimberly, Ann [US/US]; 28 Court Street, #1, New Haven,
CT 06511 (US). MAJUMDER, Kumud [IN/US]; 140
Silver Hill Lane, Stamford, CT 06905 (US). GUO, Xiaojia
[CN/US]; 713 Robert Frost Drive, Branford, CT 06405
(US). SPADERNA, Steven, K. [US/US]; 261 Deerfield
Drive, Berlin, CT 06037 (US). BOLDOG, Ferenc, L.
[HU/US]; 1687 Hartford Turnpike, North Haven, CT
06473 (US).(74) Agent: ELRIFI, Ivor, R.; Mintz, Levin, Cohn, Ferris,
Glovsky, and Popeo, P.C., One Financial Center, Boston,
MA 02111 (US).(81) Designated States (national): AF, AG, AI, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GE, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,

[Continued on next page]

(54) Title: NOVEL POLYPEPTIDES AND NUCLEIC ACIDS ENCODING SAME

(57) Abstract: The present invention provides novel isolated NOVX polynucleotides and polypeptides encoded by the NOVX polynucleotides. Also provided are the antibodies that immunospecifically bind to a NOVX polypeptide or any derivative, variant, mutant or fragment of the NOVX polypeptide, polynucleotide or antibody. The invention additionally provides methods in which the NOVX polypeptide, polynucleotide and antibody are utilized in the detection and treatment of a broad range of pathological states, as well as to other uses.

WO 01/61009 A2



LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

Published:

— without international search report and to be republished
upon receipt of that report

- (84) **Designated States (regional):** ARIPO patent (GI, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

NOVEL POLYPEPTIDES AND NUCLEIC ACIDS ENCODING SAME

TECHNICAL FIELD OF THE INVENTION

5 The invention generally relates to nucleic acids and polypeptides encoded therefrom.

BACKGROUND OF THE INVENTION

The invention generally relates to nucleic acids and polypeptides encoded therefrom. More specifically, the invention relates to nucleic acids encoding cytoplasmic, nuclear,
10 membrane bound, and secreted polypeptides, as well as vectors, host cells, antibodies, and recombinant methods for producing these nucleic acids and polypeptides.

SUMMARY OF THE INVENTION

The invention is based, in part, upon the discovery of novel polynucleotide sequences
15 encoding novel polypeptides.

Accordingly, in one aspect, the invention provides an isolated nucleic acid molecule that includes the sequence of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25 or a fragment, homolog, analog or derivative thereof. The nucleic acid can include, *e.g.*, a nucleic acid sequence encoding a polypeptide at least 85% identical to a polypeptide that includes the
20 amino acid sequences of SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26. The nucleic acid can be, *e.g.*, a genomic DNA fragment, or a cDNA molecule.

Also included in the invention is a vector containing one or more of the nucleic acids described herein, and a cell containing the vectors or nucleic acids described herein.

The invention is also directed to host cells transformed with a vector comprising any of
25 the nucleic acid molecules described above.

In another aspect, the invention includes a pharmaceutical composition that includes a NOVX nucleic acid and a pharmaceutically acceptable carrier or diluent.

In a further aspect, the invention includes a substantially purified NOVX polypeptide, *e.g.*, any of the NOVX polypeptides encoded by an NOVX nucleic acid, and fragments,
30 homologs, analogs, and derivatives thereof. The invention also includes a pharmaceutical composition that includes an NOVX polypeptide and a pharmaceutically acceptable carrier or diluent.

In still a further aspect, the invention provides an antibody that binds specifically to an NOVX polypeptide. The antibody can be, *e.g.*, a monoclonal or polyclonal antibody, and fragments, homologs, analogs, and derivatives thereof. The invention also includes a pharmaceutical composition including NOVX antibody and a pharmaceutically acceptable carrier or diluent. The invention is also directed to isolated antibodies that bind to an epitope on a polypeptide encoded by any of the nucleic acid molecules described above.

The invention also includes kits comprising any of the pharmaceutical compositions described above.

The invention further provides a method for producing an NOVX polypeptide by providing a cell containing an NOVX nucleic acid, *e.g.*, a vector that includes an NOVX nucleic acid, and culturing the cell under conditions sufficient to express the NOVX polypeptide encoded by the nucleic acid. The expressed NOVX polypeptide is then recovered from the cell. Preferably, the cell produces little or no endogenous NOVX polypeptide. The cell can be, *e.g.*, a prokaryotic cell or eukaryotic cell.

The invention is also directed to methods of identifying an NOVX polypeptide or nucleic acid in a sample by contacting the sample with a compound that specifically binds to the polypeptide or nucleic acid, and detecting complex formation, if present.

The invention further provides methods of identifying a compound that modulates the activity of an NOVX polypeptide by contacting an NOVX polypeptide with a compound and determining whether the NOVX polypeptide activity is modified.

The invention is also directed to compounds that modulate NOVX polypeptide activity identified by contacting an NOVX polypeptide with the compound and determining whether the compound modifies activity of the NOVX polypeptide, binds to the NOVX polypeptide, or binds to a nucleic acid molecule encoding an NOVX polypeptide.

In another aspect, the invention provides a method of determining the presence of or predisposition of an NOVX-associated disorder in a subject. The method includes providing a sample from the subject and measuring the amount of NOVX polypeptide in the subject sample. The amount of NOVX polypeptide in the subject sample is then compared to the amount of NOVX polypeptide in a control sample. An alteration in the amount of NOVX polypeptide in the subject protein sample relative to the amount of NOVX polypeptide in the control protein sample indicates the subject has a tissue proliferation-associated condition. A control sample is preferably taken from a matched individual, *i.e.*, an individual of similar age,

sex, or other general condition but who is not suspected of having a tissue proliferation-associated condition. Alternatively, the control sample may be taken from the subject at a time when the subject is not suspected of having a tissue proliferation-associated disorder. In some embodiments, the NOVX is detected using an NOVX antibody.

5 In a further aspect, the invention provides a method of determining the presence of or predisposition of an NOVX-associated disorder in a subject. The method includes providing a nucleic acid sample, *e.g.*, RNA or DNA, or both, from the subject and measuring the amount of the NOVX nucleic acid in the subject nucleic acid sample. The amount of NOVX nucleic acid sample in the subject nucleic acid is then compared to the amount of an NOVX nucleic acid in a control sample. An alteration in the amount of NOVX nucleic acid in the sample
10 relative to the amount of NOVX in the control sample indicates the subject has a NOVX-associated disorder.

In a still further aspect, the invention provides a method of treating or preventing or delaying an NOVX-associated disorder. The method includes administering to a subject in
15 which such treatment or prevention or delay is desired an NOVX nucleic acid, an NOVX polypeptide, or an NOVX antibody in an amount sufficient to treat, prevent, or delay a NOVX-associated disorder in the subject.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention
20 belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In the case of conflict, the present specification, including definitions, will control. In addition, the materials, methods,
25 and examples are illustrative only and not intended to be limiting.

Other features and advantages of the invention will be apparent from the following detailed description and claims.

DETAILED DESCRIPTION OF THE INVENTION

30 The present invention provides novel nucleotides and polypeptides encoded thereby. Included in the invention are the novel nucleic acid sequences and their polypeptides. The sequences are collectively referred to as "NOVX nucleic acids" or "NOVX polynucleotides"

and the corresponding encoded polypeptides are referred to as "NOVX polypeptides" or "NOVX proteins." Unless indicated otherwise, "NOVX" is meant to refer to any of the novel sequences disclosed herein. Table 1 provides a summary of the NOVX nucleic acids and their encoded polypeptides. Example 1 provides a description of how the novel nucleic acids were identified.

TABLE 1. Sequences and Corresponding SEQ ID Numbers

NOVX Assignment	Internal Identification	SEQ ID NO (nucleic acid)	SEQ ID NO (polypeptide)	Homology
1	83350421_EXT_REVCOMP	1	2	GAGE-like proteins
2	83350434_EXT_REVCOMP	3	4	GAGE-like proteins
3	83350421.0.46	5	6	GAGE-like proteins
4	8361984_EXT	7	8	GAGE-like proteins
5	Ba403c19_A	9	10	Interferons
6	AC021427_A_da1	11	12	Interferons
7	30179370_EXT	13	14	GPCR-like proteins
8	c333e1_A	15	16	Mast Cell Proteases
9	Ba328m14_A	17	18	Hepatocyte Nuclear Factors
10	C333e1_B	19	20	Mast Cell Proteases
11	AL031711_A_EX T	21	22	Mast Cell Proteases
12	S562_7F	23	24	Mast Cell Proteases
13	CG56242-01	25	26	Mast Cell Proteases

Where GAGE is the G Antigen protein family and GPCR is a G-Protein Coupled Receptor.

NOVX nucleic acids and their encoded polypeptides are useful in a variety of applications and contexts. The various NOVX nucleic acids and polypeptides according to the invention are useful as novel members of the protein families according to the presence of domains and sequence relatedness to previously described proteins. Additionally, NOVX nucleic acids and polypeptides can also be used to identify proteins that are members of the family to which the NOVX polypeptides belong.

For example, NOV1-4 are homologous to members of the G-Antigen (GAGE) family of proteins that are expressed on cancer cells, *e.g.* human melanoma, and recognized by immune cells, *e.g.* cytolytic T lymphocytes (CTLs). Thus, the NOV1-4 nucleic acids and

polypeptides, antibodies and related compounds according to the invention will be useful in therapeutic and diagnostic applications in proliferative disorders, *e.g.* malignant cancer.

Also, NOV5-6 are homologous to the Trophoblast Protein-1 protein family, belonging to the INF-alpha II subclass of the INF-alpha family. Thus, the NOV5-6 nucleic acids and polypeptides, antibodies and related compounds according to the invention will be useful in therapeutic applications in disorders of maternal recognition, proliferative disorders, *e.g.* cancer, and viral infections, *e.g.* AIDS and hepatitis.

Also, a NOV7 polypeptide is homologous to members of the seven-pass transmembrane receptor family, specifically the G-protein coupled receptors (GPCRs). Thus, the NOV7 nucleic acids and polypeptides, antibodies and related compounds according to the invention will be useful in therapeutic applications in neurological and olfactory disorders, and proliferative disorders, *e.g.* cancer.

Further, NOV8 and NOV10-13 are homologous to members of the mast cell protease family. Thus, the NOV8 and NOV10-13 nucleic acids and polypeptides, antibodies and related compounds according to the invention will be useful in therapeutic applications in proliferative disorders, *e.g.* mastocytosis.

Also, NOV9 is homologous to the hepatocyte nuclear factor-3/forkhead family of proteins. Thus, the NOV9 nucleic acids and polypeptides, antibodies and related compounds according to the invention will be useful in therapeutic applications in hepatic disorders, *e.g.* liver cancer, cirrhosis, ischaemia-re-perfusion injury, and diabetes.

The NOVX nucleic acids and polypeptides can also be used to screen for molecules, which inhibit or enhance NOVX activity or function. Specifically, the nucleic acids and polypeptides according to the invention may be used as targets for the identification of small molecules that modulate or inhibit, *e.g.*, neurogenesis, cell differentiation, cell motility, cell proliferation, hematopoiesis, and angiogenesis.

Additional utilities for the NOVX nucleic acids and polypeptides according to the invention are disclosed herein.

NOV1

A NOV1 sequence according to the invention is a nucleic acid sequence encoding a polypeptide related to the G-antigen (GAGE) family of proteins. A NOV1 nucleic acid is expressed in infant, 8-9 weeks post-partum, and in placenta. A NOV1 nucleic acid and its

encoded polypeptide includes the sequences shown in Table 2. The disclosed nucleic acid (SEQ ID NO:1) is 458 nucleotides in length and contains an open reading frame (ORF) that begins with an ATG initiation codon at nucleotides 61-63 and ends with a TAG stop codon at nucleotides 343-345. The representative ORF encodes a 94 amino acid polypeptide (SEQ ID NO:2) with a predicted molecular weight of 10,366.1 daltons (Da). PSORT analysis of a NOV1 polypeptide predicts a cytoplasmic protein with a certainty of 0.6500. Putative untranslated regions upstream and downstream of the coding sequence are underlined in SEQ ID NO: 1.

TABLE 2.

TCTTCATTCTTTCCGCCATCTTGATTCTTTCTCACTGACCAAGACTCAGCCGTGGGA
AATATGAGTGAGCTTGTAAGAGCAAGATCCCAATCCTCAGAAAGAGGAAATGAC
CAAGAGTCTTCCAGCCGGTTGGATCTGTGATTGTCCAGGAGCCCACTGAGGAAA
AACGTCAAGAAGAGGAACCACCAACTGATAATCAGGGGCCTGACATGGAAGCTTT
TCAACAGGAACTGGCTCTGCTTAAGATAGAGGATGAGCCTGGAGATGGTCCTGAT
GTCAGGGAGGGTATTATGCCCACTTTTGATCTCACTAAAGTGCTGGAAGCAGGTG
ATGCGCAACCATAGGTTTCAAGCAAGACAAATGAAGACTGAAACCAAGAACGTT
ATTCTTAATCTGGAAATTTGACTGATAATATTCTCTTAATAAAGTTTAAAGTTTCT
GCAAAGAATCCTTAAA (SEQ ID NO.: 1)

MSELVRARSQSSERGNQESSQPVGSVIVQEPTEEKRQEEEPPTDNQGPDMEAFQQEL
 ALLKIEDEPGDGPDVREGIMPTFDLTKVLEAGDAQP (SEQ ID NO.: 2)

A NOV1 nucleic acid sequence has a high degree of homology (94% identity) with an uncharacterized region of human chromosome X, including clone RP11-382F24 (CHR X; EMBL Accession No.: 158819), as is shown in Table 3. Also, a NOV1 polypeptide has homology (78% identity, 82% similarity) with a member of the GAGE gene family, human PAGE-2 polypeptide (PAGE2; PatP Accession No.: Y83168), as is shown in Table 4.

TABLE 3

NOV1: 52 gtgggaaatatgagtgagcttgttaagagcaagatcccaatcctcagaaagaggaaatgac 111
 5 CHR X: 102403 gtgggaaatatgagtgagcatgtaagaacaagatcccaatcctcagaaagaggaaatgac 102462
 NOV1: 112 caagagtcttcccagccggttgatctgtgattgt 146 (SEQ ID NO.: 21)
 10 CHR X: 102463 taagagtcttcccagccagttgtatctgtgattgt 102497 (SEQ ID NO.: 22)
 NOV1: 212 tggagccttttcaacaggaactggctctgcttaagatagaggatgagcctggagatggtc 271
 15 CHR X: 43660 tggagccttttcaacaggaactggctctgcttaagatagaggatgcacctggagatggtc 43719
 NOV1: 272 ctgatgtcagggaggggtattatgccacttttgatctcactaaagtgtggaagcaggt 330 (SEQ ID
 20 NO.: 23)
 25 CHR X: 43720 ctgatgtcagggaggggactctgccacttttgatccactaaagtgtggaagcaggt 43778 (SEQ
 ID NO.: 24)
 NOV1: 343 taggtttcaagcaagacaaatgaagactgaaaccaagaacgttattcttaatctggaaat 402
 30 CHR X: 105143 taggtttaaccaagacaaatgaggactgaaaccaagaatcttattcttaatctggaaat 105202
 NOV1: 403 ttgactgataatattctcttaataaagtttta-agttttctgcaaagaatcctt 455 (SEQ ID
 35 NO.: 25)
 40 CHR X: 105203 ttgactgataacattctcctaacaagttttacagttttctgcaaagaatcctt 105256 (SEQ ID
 NO.: 26)

TABLE 4

40 NOV1: 1 MSELVRARSQSSERGNQESSQPVGSVIVQEPTEEKRQEEEPPTDNQG-----PD 50
 *****+*****
 PAGE2: 1 MSELVRARSQSSERGNQESSQPVGSVIVQEPTEEKRQEEEPPTDNQDIEPGQEREGTPP 60
 45 NOV1: 51 MEAFQ-----QELALLKIEDEPGDGPVREGIMPTFDLTKVLEAGDAQP 94 (SEQ ID NO.:
 27)
 +* + **+*****
 PAGE2: 61 IEERKVEGDCQEMALLKIEDEPGDGPVREGIMPTFDLTKVLEAGDAQP 109 (SEQ ID NO.:
 28)
 50 Where * indicates identity and + indicates similarity.

Many human tumors express antigens that are recognized *in vitro* by cytolytic T lymphocytes (CTLs) derived from the tumor-bearing patient. The GAGE gene family members encode such antigens. Family members include GAGE (G antigen), PAGE (Prostate cancer antigen), MAGE (melanoma-specific antigen), XAGE, RAGE, and BAGE. NOV1 represents a new member of the GAGE family, and a NOV1 nucleic acid was identified in placenta and newborn, 8-9 weeks post-partum. NOV1 can be used to detect prostate, placental

and newborn tissue, and is useful in determining changes in expression of genes contained within the GAGE-like protein family. NOV1 satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the treatment of disorders associated with alterations in the expression of members of prostate cancer-associated proteins. NOV1 nucleic acids, polypeptides, antibodies, and other compositions of the present invention are useful in the treatment and/or diagnosis of a variety of diseases and pathologies, including by way of nonlimiting example, those involving prostate cancer, melanoma, and diseases of reproductive health, *e.g.* infertility, sudden infant death syndrome, and newborn failure to thrive.

NOV2

A NOV2 sequence according to the invention is a nucleic acid sequence encoding a polypeptide related to the GAGE family of proteins. A NOV2 nucleic acid is expressed in infant, 8-9 weeks post-partum, and in placenta. A NOV2 nucleic acid and its encoded polypeptide includes the sequences shown in Table 5. The disclosed nucleic acid (SEQ ID NO:3) is 475 nucleotides in length and contains an open reading frame (ORF) that begins with an ATG initiation codon at nucleotides 25-27 and ends with a TAG stop codon at nucleotides 358-360. The representative ORF encodes a 111 amino acid polypeptide (SEQ ID NO:4) with a predicted molecular weight of 12,040.9 daltons (Da). PSORT analysis of a NOV2 polypeptide predicts a cytoplasmic protein with a certainty of 0.6500. Putative untranslated regions upstream and downstream of the coding sequence are underlined in SEQ ID NO: 3.

TABLE 5.

GGGAAGAGACCATGTGTGGGAAATATGAGTGAGCATGTGAGAACAAGATCCCAA
TCCTCAGAAAGAGGAAATGACCAAGAGTCTTCCCAGCCAGTTGGATCTGTGATTG
TCCAGGAGCCCCTGAGGAAAAACGTCAAGAAGAGGAACCACCAACTGATAATC
AGGGTATTGCACCTAGTGGGGAGATCGAAAATGAAGGAGCACCTGCCGTTCAAGG
GCCTGACATGGAAGCTTTTCAACAGGAACTGGCTCTGCTTAAGATAGAGGATGAG
CCTGGAGATGGTCCTGATGTCAGGGAGGGGATTATGCCCACTTTTGATCTCACTAA
AGTGCTGGAAGCAGGTGATGCGCAACCATAGGTTTCAAGCAAGACAAATGAAGA
CTGAAACCAAGAACGTTATTCTTAATCTGGAAATTTGACTGATAATATTCTCTTAA
TAAAGTTTTAAGTTTTCTGCAAAGAAAAAAAAAAAAA (SEQ ID NO.: 3)

MSEHVRTRSQSSERGNDQESSQPVGSVIVQEPTEEKRQEEEPPTDNQGLAPSGEIENEG
 APAVQGPDMFAFQQELALLKIEDEPGDGPVREGIMPTFDLTKVLEAGDAQP (SEQ ID
 NO.: 4)

- 5 A NOV2 nucleic acid sequence has a high degree of homology (95% identity) with an
 uncharacterized region of human chromosome X, including clone RP11-382F24 (CHR X;
 EMBL Accession No.: 158819), as is shown in Table 6. Also, a NOV2 polypeptide has
 homology (81% identity, 86% similarity) with a member of the GAGE gene family, human
 PAGE-2 polypeptide (PAGE2; PatP Accession No.: Y83168), as is shown in Table 7. Further,
 10 a NOV2 polypeptide has homology with another member of the GAGE gene family, PAGE-1
 (PAGE1; GenBank Accession No.: AAC25990.1), as is shown in Table 8.

TABLE 6.

15	NOV2: 16	gtgggaaatatgagtgagcatgtgagaacaagatcccaatcctcagaaaggagaaatgac 75
	CHR X: 102403	gtgggaaatatgagtgagcatgtgagaacaagatcccaatcctcagaaaggagaaatgac 102462
20	NOV2: 76	caagagtcttcccagccagttggatctgtgattgt 110 (SEQ ID NO.: 29)
	CHR X: 102463	taagagtcttcccagccagttgtatctgtgattgt 102497 (SEQ ID NO.: 30)
25	NOV2: 109	gtccaggagccactgaggaaaaacgtcaagaagagggaaccaccaactgataatcagggt 168
	CHR X: 102893	gtccaggagccactgaggaaaaacgtcaagaagagggaaccaccaactgataatcagggt 102952
30	NOV2: 169	attgcacctagtggggagatcgaaaatgaaggagcacctgccgttcaagg 218 (SEQ ID NO.: 31)
	CHR X: 102953	attgcacctactggggagatcgaaaatgaagcgccacctgcccttcaagg 103002 (SEQ ID NO.: 32)
35	NOV2: 227	tggaagcttttcaacaggaactggctctgcttaagatagaggatgagcctggagatggtc 286
	CHR X: 103759	tggaagcttttcaacaggaactggctctgcttaagatagaggatgagcctggagatggtc 103818
40	NOV2: 287	ctgatgtcaggaggaggattatgccacttttgatctcactaaagtgctggaagcagggt 345 (SEQ ID NO.: 33)
	CHR X: 103819	ctgatgtcaggaggaggactctgccactttcgatccactaaagtgctggaagcagggt 103877 (SEQ ID NO.: 34)
45	NOV2: 342	agggtgatgcgcaaccataggtttcaagcaagacaaatgaagactgaaaccaagaacgtta 401
	CHR X: 109697	aggtaatgggcaaccataggttttaaccaagacaaatgaagactgaaaccaagaatgttg 109756
50	NOV2: 402	ttcttaatctggaaatttgactgataatattctcttaataaagtttta-agttttctgca 460
	CHR X: 109757	ttcttatgctggaaatttgactgctaacattctcttaataaagttttacagttttctgca 109816
55		

NOV2: 461 aa 462 (SEQ ID NO.: 35)
 ||
 CHR X: 109817 aa 109818 (SEQ ID NO.: 36)

5

TABLE 7.

NOV2: 1 MSEHVRTRSQSSSERGNDQESSQPVGSVIVQEPTEEKRQEEEPPTDNQGIAPSGEIEENEGA 60
 *** ** *****+***** * * +* * **
 10 PAGE2: 1 MSELVRARSQSSSERGNDQESSQPVGSVIVQEPTEEKRQEEEPPTDNQDIEP-GQ-EREQT 58

NOV2: 61 PAVQGPDMFAFQQELALLKIEDEPGDGPVREGIMPTFDLTKVLEAGDAQP 120 (SEQ ID
 NO.: 37)
 * ++ +* **+*****
 15 PAGE2: 59 PPIEERKVEGDCQEMALLKIEDEPGDGPVREGIMPTFDLTKVLEAGDAQP 109 (SEQ ID
 NO.: 38)

Where * indicates identity and + indicates similarity.

20

TABLE 8.

NOV2: 46 NQGIAPSGEIEENEGAPAVQGPDMFAFQQELALLKIEDEPGDGPVREGIMPTFDLTKVLEAGDAQP 91 (SEQ ID NO.: 39)
 ++ * +* * * * * + * * * * * +
 25 PAGE1: 37 SQDSTPAEEREDEGASAAQGEPEADSQELVQPKTGCEPGDGPDTK 82 (SEQ ID NO.: 40)
 Where * indicates identity and + indicates similarity.

Many human tumors express antigens that are recognized in vitro by cytolytic T lymphocytes (CTLs) derived from the tumor-bearing patient. The GAGE gene family members encode such antigens. Family members include GAGE (G antigen), PAGE (Prostate cancer antigen), MAGE (melanoma-specific antigen), XAGE, RAGE, and BAGE. NOV2 represents a new member of the GAGE family, and a NOV2 nucleic acid was identified in placenta and newborn, 8-9 weeks post-partum. NOV2 can be used to detect prostate, placental and newborn tissue, and is useful in determining changes in expression of genes contained within the GAGE-like protein family. NOV2 satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the treatment of disorders associated with alterations in the expression of members of prostate cancer-associated proteins. NOV2 nucleic acids, polypeptides, antibodies, and other compositions of the present invention are useful in the treatment and/or diagnosis of a variety of diseases and pathologies, including by way of nonlimiting example, those involving prostate cancer, melanoma, and diseases of reproductive health, e.g. infertility, sudden infant death syndrome, and newborn failure to thrive.

NOV3

A NOV3 sequence according to the invention is a nucleic acid sequence encoding a polypeptide related to the GAGE family of proteins. A NOV3 nucleic acid is expressed in infant, 8-9 weeks post-partum, and in placenta. A NOV3 nucleic acid and its encoded polypeptide includes the sequences shown in Table 9. The disclosed nucleic acid (SEQ ID NO:5) is 1,051 nucleotides in length and contains an open reading frame (ORF) that begins with an ATG initiation codon at nucleotides 593-595 and ends with a TAG stop codon at nucleotides 926-928. The representative ORF encodes a 111 amino acid polypeptide (SEQ ID NO:6) with a predicted molecular weight of 12,076 daltons (Da). PSORT analysis of a NOV3 polypeptide predicts a cytoplasmic protein with a certainty of 0.6500. Putative untranslated regions upstream and downstream of the coding sequence are underlined in SEQ ID NO: 5.

TABLE 9.

CGGGAAGGGACCGTGTGGTCAGTAACGAAGGGGCTTGGGAACTGGGAGCGCGGT
 GGGCTGGTGACTGTGGCCCCGAGGTCTGTAGAGTGCCTGGCAGAGGTGTCCTGTG
 AGGAGCATAACGTTCACTCTGTTTCACAATTTCTCACCTCCGCCATGGACATCATA
 GGAAGGAATGGGCGAGGCTGTGCTTTCCAACAAGACTTGATTTTGAGAGGGGTGT
 GGGGGTGAAATGGGCCTAGCAAATCAGAGTGGGACAAAAGCAGTAGTCATTTCA
 GTTTCAATTCTCTGCCCGTTTTTTCCTAAATGTCTTCATGATGGAGAGTCTAATTGT
 GAAACCAAAACGCAGAAATGTCCTCTGTCTTTTGCTATGGCGTTAAGGGGATTTCT
 ATGCCTCTTCGACTATGATACAAACAAATCTGTCTTAGTTTGATTTCGAAAGCATG
 TGTACTTATCATTGCTCTGTGACTTAATTTGAAAATATTTTCAAATTAATAAAGT
 ACAATCACCATTTTGCCGTGGAATGTTTCATATATACTAAGTTCTTACACACT
 TTTTCAAATAACAATATTCTGTTTGCAGTGGGAAATATGAGTGAGCTTGTAAGA
 GCAAGATCCCAATCCTCAGAAAGAGGAAATGACCAAGAGTCTTCCCAGCCGGTTG
 GATCTGTGATTGTCCAGGAGCCCACTGAGGAAAAACGTCAAGAAGAGGAACCAC
 CAACTGATAATCAGGGTATTGCACCTAGTGGGGAGATTGAAAATCAAGCAGTGCC
 TGCTTTTCAAGGGCCTGACATGGAAGCTTTTCAACAGGAACTGGCTCTGCTTAAGA
 TAGAGGATGAGCCTGGAGATGGTCCTGATGTCAGGGAGGGTATTATGCCCACTTT
 TGATCTCACTAAAGTGCTGGAAGCAGGTGATGCGCAACCATAGGTTTCAAGCAAG
 ACAATGAAGACTGAAACCAAGAACGTTATTCTTAATCTGGAAATTTGACTGATA
 ATATTCTCTTAATAAAGTTTAAAGTTTCTGCAAAGAAAAAAAAAAAAAAAAAAAAA
 A (SEQ ID NO.: 5)

MSELVRARSQSSERGNDQESSQPVGSVIVQEPTEEKRQEEEPPTDNQGIAPSGEIENQA
 VPAFQGPDMFAFQQELALLKIEDEPGDGPVREGIMPTFDLTKVLEAGDAQP (SEQ ID
 NO.: 6)

A NOV3 nucleic acid sequence has a high degree of homology (92% identity) with an uncharacterized region of human chromosome X, including clone RP11-382F24 (CHR X; EMBL Accession No.: 158819), as is shown in Table 10. Also, a NOV3 nucleic acid has a high degree of homology (97% identity) with a NOV2 nucleic acid, as shown in Table 11.

- 5 NOV3 polypeptide has homology (40% identity, 49% similarity) with a member of the GAGE gene family, human PAGE-1 polypeptide (PAGE1; EMBL ACCESSION NO.: O60829), as is shown in Table 12.

TABLE 10.

10	NOV3: 243	atcagagtgggacaaaagcagtagtcatttcagtttcaattctctgcccg-ttttttctct	301
	CHR X: 42062	atcagagtgggaggaaagcagcagtcacttcagtttcaatttctgcccgttttttctct	42121
15	NOV3: 302	aaatgtcttcatgatggagagtctaattgtgaacccaaacgcagaaatgtcctctgtct	361
	CHR X: 42122	aaatgtgtaaatgatggagagtctaattgtgaagccaaactcagaaaagtcctctgtct	42181
20	NOV3: 362	tttgctatggcggttaaggggatttctatgcctcttcgactatgatacaaacaaatctgtc	421
	CHR X: 42182	tttgctatggcggttaaggtgttttctgtgcctcttcgactatgatacaaacaaatctgtc	42241
25	NOV3: 422	cttagtttgattcgaaagcatgtgtacttatcattgctctgtgacttaatttgaaaatat	481
	CHR X: 42242	cttagtttgattgaaagcatgcgtacttatcaatgctctgtgacttagtttgaaaatat	42301
30	NOV3: 482	tttcaaaattaaaaaagtacaaatcaccattttgccgtggaatgttcatatatataacta	541
	CHR X: 42302	tttcaaaattaaaaaagtacaaatcaccattttgccatggaatgttcatatatatagcta	42361
35	NOV3: 542	agttcttacacactttttccaataacaatatctgtttgcagtgaggaaatatgagtgag	601
	CHR X: 42362	agttcttacacactttttccaataacaatatctgtttgcagtgagagatatgagtgag	42421
40	NOV3: 602	cttgtaagagcaagatcccaatcctcagaaagggaatgaccaagagtcttcccagccg	661
	CHR X: 42422	catgtaa---caagatcccaatcctcagaaagggaatgaccaagagtcttcccagcca	42478
	NOV3: 662	gttggatctgtgattgt	678 (SEQ ID NO.: 41)
	CHR X: 42479	gttggacctgtgattgt	42495 (SEQ ID NO.: 42)

TABLE 11.

45	NOV3: 584	gtgggaaatatgagtgagcttgtaagagcaagatcccaatcctcagaaaggagaaatgac	643
	NOV2: 16	gtgggaaatatgagtgagcatgtgagaacaagatcccaatcctcagaaaggagaaatgac	75
50	NOV3: 644	caagagtcttcccagccggttgatctgtgattgtccaggagcccactgaggaaaaacgt	703
	NOV2: 76	caagagtcttcccagccagttggatctgtgattgtccaggagcccactgaggaaaaacgt	135
55	NOV3: 704	caagaagaggaaccaccaactgataatcagggattgtcacctagtggggagattgaaaat	763
	NOV2: 136	caagaagaggaaccaccaactgataatcagggattgtcacctagtggggagatcgaaaat	195
	NOV3: 764	caagcagtgctgcttttcaaggcctgacatggaagcttttcaacaggaactggctctg	823

NOV2: 196 gaaggagcacctgccgttcaaggcctgacatggaagcttttcaacaggaactggctctg 255
 NOV3: 824 ctttaagatagaggatgagcctggagatggctcctgatgtcagggaggggtattatgccact 883
 5 NOV2: 256 ctttaagatagaggatgagcctggagatggctcctgatgtcagggaggggattatgccact 315
 NOV3: 884 tttgatctcactaaagtgtctggaagcaggtgatgcgcaaccataggtttcaagcaagaca 943
 10 NOV2: 316 tttgatctcactaaagtgtctggaagcaggtgatgcgcaaccataggtttcaagcaagaca 375
 NOV3: 944 aatgaagactgaaaccaagaacgttattcttaatctggaaatttgactgataatattctc 1003
 NOV2: 376 aatgaagactgaaaccaagaacgttattcttaatctggaaatttgactgataatattctc 435
 15 NOV3: 1004 ttaataaagttttaagttttctgcaaag 1031 (SEQ ID NO.: 43)
 NOV2: 436 ttaataaagttttaagttttctgcaaag 463 (SEQ ID NO.: 44)

TABLE 12.

NOV3: 6 RARSQSSERGNDAQESSQPVGSVIVXXXXXXXXXXXXXXXXXDNQGIAPSGEIQAVPAFQG 65
 * **+* **+ **+ +* *** * * * * * * *
 25 PAGE1: 4 RVRSRSRGRGDGQEAPD-----VVAFVAPGESQQEEPPTDNQDIEPGQEREG--TPPIEE 56
 NOV3: 66 PDMEAFQQELALLKIEDEPGDGPVREGIMPTFDLTKVLEAGDAQP 111 (SEQ ID NO.: 45)
 +* **+ * * * * * * * * * * * * * * * *
 PAGE1: 57 RKVEGDCEMDLEKTRSERGDGSDVKEKTPPNPKHAKTKEAGDGQP 102 (SEQ ID NO.: 46)
 30 Where * indicates identity and + indicates similarity.

Many human tumors express antigens that are recognized in vitro by cytolytic T lymphocytes (CTLs) derived from the tumor-bearing patient. The GAGE gene family members encode such antigens. Family members include GAGE (G antigen), PAGE (Prostate cancer antigen), MAGE (melanoma-specific antigen), XAGE, RAGE, and BAGE. NOV3 represents a new member of the GAGE family, and a NOV3 nucleic acid was identified in placenta and newborn, 8-9 weeks post-partum. NOV3 can be used to detect prostate, placental and newborn tissue, and is useful in determining changes in expression of genes contained within the GAGE-like protein family. NOV3 satisfies a need in the art by providing new

40 diagnostic or therapeutic compositions useful in the treatment of disorders associated with alterations in the expression of members of prostate cancer-associated proteins. NOV3 nucleic acids, polypeptides, antibodies, and other compositions of the present invention are useful in the treatment and/or diagnosis of a variety of diseases and pathologies, including by way of nonlimiting example, those involving prostate cancer, melanoma, and diseases of reproductive

45 health, e.g. infertility, sudden infant death syndrome, and newborn failure to thrive.

NOV4

A NOV4 sequence according to the invention is a nucleic acid sequence encoding a polypeptide related to the GAGE family of proteins. A NOV4 nucleic acid is expressed in adult brain, fetal brain, pregnant uterus, in placenta, and in the cell line JAR. A NOV4 nucleic acid and its encoded polypeptide includes the sequences shown in Table 13. The disclosed nucleic acid (SEQ ID NO:7) is 611 nucleotides in length and contains an open reading frame (ORF) that begins with an ATG initiation codon at nucleotides 174-176 and ends with a TAA stop codon at nucleotides 519-521. The representative ORF encodes a 115 amino acid polypeptide (SEQ ID NO:8) with a predicted molecular weight of 13,656 daltons (Da).

PSORT analysis of a NOV4 polypeptide predicts a nuclear protein with a certainty of 0.8400. Putative untranslated regions upstream and downstream of the coding sequence are underlined in SEQ ID NO: 7.

TABLE 13.

CATGCAGAGTACACTGGGCATCTTCCCTTCGACCCCTTTGCCACGTGGTGACCGC
TGGGGGAGCTGTGAGAGTGTGAGGGACACGTTCCAGCCGTCTGGACTCTTTCTCTC
CTACTGAGACGCAGCCTATAGGTCCGCAGGCCAGTCCTCCCAGGAACTGAAATAG
TGAAATATGAGTTGGCGAGGAAGATCAACATATAGGCCTAGGCCAAGAAGAAGT
TTACAGCCTCCTGAGCTGATTGGGGCTATGCTTGAACCCACTGATGAAGAGCCTA
AAGAAGAGAAACCACCCACTAAAAGTCGGAATCCTACACCTGACTCNAGAAGAG
AGAAGATGATCAGGGTGCAGCTGAGATTCAAGTGCCTGACCTGGAAGCCGATCTC
CAGGAGCTATGTCAGACAAAGACTGGGGATGGATGTGAAGGTGGTACTGATGTCA
AGGGGAAGATTCTACCAAAAGCAGAGCACTTTAAAATGCCAGAAGCAGGTGAAG
GGAAATCACAGGTTTAAAGGAAGATAAGCTGAAACAACACAAACTGTTTTTATAT
TAGATATTTTACTTTTAAAATATCTTAATAAAGTTTTAAGCTTTTCTCCAAAAAAA
AAAAAA (SEQ ID NO.: 7)

MSWRGRSTYRPRPRRSLQPPELIGAMLEPTDEEPKEEKPPTKSRNPTPDSRREKMIRVQ
 LRFKCLTWKPISRSYVRQRLGMDVKVVLMSRGRFYQKQSTLKCQKQVKGNHRFKGR
 (SEQ ID NO.: 8)

A NOV4 nucleic acid sequence has a high degree of homology (92% identity) with a region of the GAGE-2 protein mRNA (GAGE2; GenBank Accession No.: HSU19143), as is shown in Table 14. Also, a NOV4 polypeptide has homology (48% identity, 62% similarity) with a member of the GAGE gene family, human GAGE-2 polypeptide (GAGE2; EMBL Accession No.: AAC33676), as is shown in Table 15.

TABLE 14.

NOV4:	167	gtgaaatatgagttggcgaggaagatcaacatataggcctaggccaaga	215	(SEQ ID NO.: 47)
GAGE2	77	gtgaaatatgagttggcgaggaagatcgacctatcgccctagaccaaga	125	(SEQ ID NO.: 48)

TABLE 15.

NOV4:	1	MSWRGRSTYRPRRRSLQPPELIGAMLEP--TDE----EPKEEKPPTKSRNP	46
		*****+*****+***+**++**++	
GAGE2:	1	MSWRGRSTYRPRRRYVEPPEMIGPMRPEQFSDEVEPATPEEGEPATQRQDP	52
NOV4:	47	DDQGAAEIQVPDLEADLQELCQTKTGDCGGTDVKGKILPKAEHFKMPEAGEGKSQ	104
		(SEQ ID NO.: 49)	
		++++* * *++* * +* * * * + * * * * *++	
GAGE2:	59	EDEGASAGQGPKEAESQEQGHPQTGCECEDGPDGQEMDPPNPPEEVKTPEEGEKQSQ	115
		(SEQ ID NO.: 50)	
Where * indicates identity and + indicates similarity.			

Many human tumors express antigens that are recognized in vitro by cytolytic T lymphocytes (CTLs) derived from the tumor-bearing patient. The GAGE gene family members encode such antigens. Family members include GAGE (G antigen), PAGE (Prostate cancer antigen), MAGE (melanoma-specific antigen), XAGE, RAGE, and BAGE. NOV4 represents a new member of the GAGE family, and a NOV4 nucleic acid was identified in brain, fetal brain, placenta and pregnant uterus. NOV4 can be used to detect brain, prostate, placental and uterine tissue, and is useful in determining changes in expression of genes contained within the GAGE-like protein family. NOV4 satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the treatment of disorders associated with alterations in the expression of members of prostate cancer-associated proteins. NOV4 nucleic acids, polypeptides, antibodies, and other compositions of the present invention are useful in the treatment and/or diagnosis of a variety of diseases and pathologies, including by way of nonlimiting example, those involving prostate cancer, melanoma, and diseases of reproductive health, e.g. infertility and placental insufficiency.

NOV5

A NOV5 sequence according to the invention is a nucleic acid sequence encoding a polypeptide related to the interferon family of proteins. A NOV5 nucleic acid and its encoded polypeptide includes the sequences shown in Table 16. The disclosed nucleic acid (SEQ ID NO:9) is 673 nucleotides in length and contains an open reading frame (ORF) that begins with an ATG initiation codon at nucleotides 34-36 and ends with a TAA stop codon at nucleotides 637-639. The representative ORF encodes a 207 amino acid polypeptide (SEQ ID NO:10) with a predicted molecular weight of 25,218 daltons (Da). PSORT analysis of a NOV5 polypeptide predicts a plasma membrane protein with a certainty of 0.8110. SIGNALP analysis suggests a signal peptide with the likely cleavage site between positions 27 and 28 of SEQ ID NO.: 10. Putative untranslated regions upstream and downstream of the coding sequence are underlined in SEQ ID NO: 9.

TABLE 16.

TAGCTTGCAAAAAAAAAATGAGCACCAAACCTGATATGATTCAAAAGTGTGTTGTGGC
TTGAGATCCTTATGGGTATATTCATTGCTGGCACCCCTATCCCTGGACTGTAACCTA
CTGAACGTTACCTGAGAAGAGTCACCTGGCAAAATCTGAGACATCTGAGTAGTA
TGAGCAATTCATTTCTGTAGAATGTCTACGAGAAAACATAGCTTTTGAGTTGCCC
CAAGAGTTTCTGCAATACACCCAACCTATGAAGAGGGACATCAAGAAGGCCTTCT
ATGAAATGTCCCTACAGGCCTTCAACATCTTCAGCCAACACACCTTCAAATATTGG
AAAGAGAGACACCTCAAACAAATCCAAATAGGACTTGATCAGCAAGCAGAGTAC
CTGAACCAATGCTTGGAGGAAGACAAGAATGAAAATGAAGACATGAAAGAAATG
AAAGAGAATGAGATGAAACCCTCAGAAGCCAGGGTCCCCCAGCTGAGCAGCCTG
GAACTGAGGAGATATTTCCACAGGATAGACAATTCCTGAAAGAAAAGAAATACA
GTGACTGTGCCTGGGAGATTGTCCGAGTGGAATCAGAAGATGTTTGTATTACTTT
TACAAATTTACAGCTCTATTCAGGAGGAAATAAGGTATATTTTTTGAATTAAAT
CCTTTCCCTC (SEQ ID NO.: 9)

MSTKPDMIQKCLWLEILMGIFLAGTSLDCNLLNVHLRRVTWQNLRLHSSMSNSFPVE
 CLRENI AFELPQEFLQYTQPMKRDIKAFYEMSLQAFNIFSQHTFKYWKERHLKQIQIG
 LDQQA EYLNQCLEEDKNENEDMKEMKENEMKPSEARVPQLSSLELRRYFHRIDNFLK
 EKKYSDCAWEIVRVEIRRCLYFYKFTALFRRK (SEQ ID NO.: 10)

A NOV5 nucleic acid sequence has a high degree of homology (100% identity) with a region of an interferon-like protein precursor mRNA, (ILP-P; Genbank Accession No.: AF146759), as is shown in Table 17. A NOV5 polypeptide has a high degree of homology

(99% identity, 100% similarity) with a member of the human keratinocyte-derived interferon (KDI) family (KDI; PatP Accession No.: Y68800), as is shown in Table 18. Also, a NOV5 polypeptide has homology (36% identity, 53% similarity) with a trophoblast protein-1 protein, also known as interferon tau-1 precursor, (INT-T; SwissEmbl Accession No.: P15696), as is shown in Table 19.

TABLE 17.

NOV5: 17	tgagcaccaaacctgatatgattcaaaagtgtttgtggcttgagatccttatgggtatat	76
10 ILP-P: 47	tgagcaccaaacctgatatgattcaaaagtgtttgtggcttgagatccttatgggtatat	106
NOV5: 77	tcattgctggcaccctatccctggactgtaacttactgaacgttcacctgagaagagtca	136
15 ILP-P: 107	tcattgctggcaccctatccctggactgtaacttactgaacgttcacctgagaagagtca	166
NOV5: 137	cctggcaaaatctgagacatctgagtagtatgagcaattcatttcctgtagaatgtctac	196
ILP-P: 167	cctggcaaaatctgagacatctgagtagtatgagcaattcatttcctgtagaatgtctac	226
20 NOV5: 197	gagaaaacatagcttttgagttgccccagaggtttctgcaatacacccaacctatgaaga	256
ILP-P: 227	gagaaaacatagcttttgagttgccccagaggtttctgcaatacacccaacctatgaaga	286
25 NOV5: 257	gggacatcaagaaggccttctatgaaatgtccctacaggccttcaacatcttcagccaac	316
ILP-P: 287	gggacatcaagaaggccttctatgaaatgtccctacaggccttcaacatcttcagccaac	346
NOV5: 317	acaccttcaaataattggaaagagagacacctcaaacaatccaaataggacttgatcagc	376
30 ILP-P: 347	acaccttcaaataattggaaagagagacacctcaaacaatccaaataggacttgatcagc	406
NOV5: 377	aagcagagtacctgaaccaatgcttgga	404 (SEQ ID NO.: 51)
35 ILP-P: 407	aagcagagtacctgaaccaatgcttgga	434 (SEQ ID NO.: 52)
NOV5: 460	ccctcagaagccagggtccccagctgagcagcctggaactgaggagatatccacagg	519
ILP-P: 490	ccctcagaagccagggtccccagctgagcagcctggaactgaggagatatccacagg	549
40 NOV5: 520	atagacaatttcctgaaagaaaagaaatacagtgactgtgcctgggagattgtccgagtg	579
ILP-P: 550	atagacaatttcctgaaagaaaagaaatacagtgactgtgcctgggagattgtccgagtg	609
45 NOV5: 580	gaaatcagaagatgtttgtattacttttacaattttacagctctattcaggaggaaataa	639
ILP-P: 610	gaaatcagaagatgtttgtattacttttacaattttacagctctattcaggaggaaataa	669
50 NOV5: 640	g	640 (SEQ ID NO.: 53)
ILP-P: 670	g	670 (SEQ ID NO.: 54)

TABLE 18.

55 NOV5: 1	MSTKPDMIQKCLWLEILMGIFIAGTSLDCNLLNVHLRRVTWQNLRLHSSMSNSFPVECL	60

15

```

NOV5:      14 LEILMGIFIAGT---LSLDCNLLNVHLRRVTWQNLRLHSSMSNSFPVECLRENIAFELPQ 70
      *  + + + +      * * * *      + +      + + + +      *  + + + +      *  + + +
20  INF-T:      5 LSLMLALVLVSYGPGRSLGACYLSEDHMLGAR-ENLRLLARMNRLSPHPCLQDRKDFGLPQ 63
NOV5:     71 EFLQYTQPMKRDIKKAIFYEMSLQAFNIF-SQHTFKYWKERHLKQIQIGLDQQA EYLNQCL 129
      *  + +      *  *      + + +      *  + + + +      + + + +      *  + + + +      *  + + + +
25  INF-T:     64 EMVEGNQLQKDQAI SVLHEMLQQCFNL FYTEHSSAAWNTTLEQLCTGLQQQLEDLDA CL 123
NOV5:    130 EEDKNENEDMKEMKENEMKPSEARVPQLSSLELRRYFHRIDNFLKEKKYSDCAWEIVRV E 189
      *      *      + + + +      *  +      *  + + + +      *  + + + + + + + + + + + + + + + +
30  INF-T:    124 GPVMGE-----KDSDM----GRMGPI--LTVKKYFQGIHVYLKEKEYSDCAWEIIRVE 170
NOV5:    190 IRRCL 194 (SEQ ID NO.: 57)
      + * *
INF-T:    171 MMRAL 175 (SEQ ID NO.: 58)
Where * indicates identity and + indicates similarity.

```

45

NOV6

A NOV6 sequence according to the invention is a nucleic acid sequence encoding a polypeptide related to the interferon family of proteins. A NOV6 nucleic acid was derived by an exon linking process using a NOV5 nucleic acid (BA403c19_A). A NOV6 nucleic acid and
 5 its encoded polypeptide includes the sequences shown in Table 20. The disclosed nucleic acid (SEQ ID NO:11) is 631 nucleotides in length and contains an open reading frame (ORF) that begins with an ATG initiation codon at nucleotides 1-3 and ends with a TAA stop codon at nucleotides 622-624. The representative ORF encodes a 207 amino acid polypeptide (SEQ ID
 10 NO:12) with a predicted molecular weight of 25,218 daltons (Da). PSORT analysis of a NOV6 polypeptide predicts a plasma membrane protein with a certainty of 0.8110. A putative untranslated region downstream of the coding sequence is underlined in SEQ ID NO: 11.

TABLE 20.

15 ATGAGCACCAAACCTGATATGATTCAAAAGTGTTTGTGGCTTGAGATCCTTATGG
 GTATATTCATTGCTGGCACCCTATCCCTGGACTGTAACCTTACTGAACGTTACCTG
 AGAAGAGTCACCTGGCAAAATCTGAGACATCTGAGTAGTATGAGCAATTCATTTCTG
 CTGTAGAATGTCTACGAGAAAACATAGCTTTTGAGTTGCCCAAGAGTTTCTGCAA
 TACACCCAACCTATGAAGAGGGACATCAAGAAGGCCTTCTATGAAATGTCCCTAC
 20 AGGCCTTCAACATCTTCAGCCAACACACCTTCAAATATTGGAAAGAGAGACACCT
 CAAACAAATCCAAATAGGACTTGATCAGCAAGCAGAGTACCTGAACCAATGCTTG
 GAGGAAGACGAGAATGAAAATGAAGACATGAAAGAAATGAAAGAGAATGAGAT
 GAAACCCTCAGAAGCCAGGGTCCCCCAGCTGAGCAGCCTGGAAGTGAAGGAGATAT
 TTCCACAGGATAGACAATTTCTGAAAGAAAAGAAATACAGTGAAGTGTGCCTGGG
 25 AGATTGTCCGAGTGGAAATCAGAAGATGTTTGTATTACTTTTACAAATTTACAGCT
 CTATTCAGGAGGAAATAAGGTATAT (SEQ ID NO.: 11)

MSTKPDMIQKCLWLEILMGIFLAGTSLDCNLLNVHLRRVTWQNLRLHSSMSNSFPVE
 CLRENIAFELPQEFLLQYTQPMKRDIKKAFYEMSLQAFNIFSQHTFKYWKERHLKQIQIG
 30 LDQQA EYLNQCLEEDENENEDMKEMKENEMKPSEARVPQLSSLELRRYFHRIDNFLK
 EKKYSDCAWEIVRVEIRRCLYFYKFTALFRRK (SEQ ID NO.: 12)

A NOV6 nucleic acid has a high degree of homology (100% identity) with a human interferon like-protein precursor, (ILP-P; Genbank Accession No.: AF146759), as is shown in
 35 Table 21. A NOV6 polypeptide has a high degree of homology (100% identity) with a human interferon-like protein precursor (ILP-P; EMBL Accession No.: AAF67468), as is shown in Table 22.

TABLE 21.

5	NOV6: 1	atgagcaccaaacctgatatgattcaaaagtgtttgtggcttgagatccttatgggtata	60
	ILP-P: 46	atgagcaccaaacctgatatgattcaaaagtgtttgtggcttgagatccttatgggtata	105
10	NOV6: 61	ttcattgctggcaccctatccctggactgtaacttactgaacgttcacctgagaagagtc	120
	ILP-P: 106	ttcattgctggcaccctatccctggactgtaacttactgaacgttcacctgagaagagtc	165
15	NOV6: 121	acctggcaaaatctgagacatctgagtagtatgagcaattcatttcctgtagaatgtcta	180
	ILP-P: 166	acctggcaaaatctgagacatctgagtagtatgagcaattcatttcctgtagaatgtcta	225
20	NOV6: 181	cgagaaaacatagctttttgagttgccccagagtttctgcaatacacccaacctatgaag	240
	ILP-P: 226	cgagaaaacatagctttttgagttgccccagagtttctgcaatacacccaacctatgaag	285
25	NOV6: 241	agggacatcaagaaggccttctatgaaatgtccctacaggccttcaacatcttcagccaa	300
	ILP-P: 286	agggacatcaagaaggccttctatgaaatgtccctacaggccttcaacatcttcagccaa	345
30	NOV6: 301	cacaccttcaaataattggaaagagagacacctcaaacaaatccaaataggacttgatcag	360
	ILP-P: 346	cacaccttcaaataattggaaagagagacacctcaaacaaatccaaataggacttgatcag	405
35	NOV6: 361	caagcagagtacctgaaccaatgcttggag	390 (SEQ ID NO.: 59)
	ILP-P: 406	caagcagagtacctgaaccaatgcttggag	435 (SEQ ID NO.: 60)
40	NOV6: 445	ccctcagaagccagggtccccagctgagcagcctggaactgaggagatatttccacagg	504
	ILP-P: 490	ccctcagaagccagggtccccagctgagcagcctggaactgaggagatatttccacagg	549
45	NOV6: 505	atagacaatttcctgaaagaaaagaaatacagtgactgtgcctgggagattgtccgagt	564
	ILP-P: 550	atagacaatttcctgaaagaaaagaaatacagtgactgtgcctgggagattgtccgagt	609
50	NOV6: 565	gaaatcagaagatgtttgtattacttttacaaatttacagctctattcaggaggaaataa	624
	ILP-P: 610	gaaatcagaagatgtttgtattacttttacaaatttacagctctattcaggaggaaataa	669
55	NOV6: 625	g	625 (SEQ ID NO.: 61)
	ILP-P: 670	g	670 (SEQ ID NO.: 62)

TABLE 22.

50	NOV6:	1	MSTKPDMIQKCLWLEILMGIFIAGTSLDCNLLNVHLRRVTWQNLRLHSSMSNSFPVECL	60
	ILP-P:	1	MSTKPDMIQKCLWLEILMGIFIAGTSLDCNLLNVHLRRVTWQNLRLHSSMSNSFPVECL	60
55	NOV6:	61	RENIAFELPQEFLQYTQPMKRDIKAFYEMSLQAFNIFSQHTFKYWKERHLKQIQIGLDQ	120
	ILP-P:	61	RENIAFELPQEFLQYTQPMKRDIKAFYEMSLQAFNIFSQHTFKYWKERHLKQIQIGLDQ	120
60	NOV6:	121	QAEYLNQCLEEDENENEDMKEMKENEMKPSEARVPQLSSLELRRYFHRIDNFLKEKKYS	180

ILP-P: 121 QAEYLNQCLEEDENENEDMKEMKENEMKPSEARVPQLSSLELRRYFHRIDNFLKEKKYSD
180

NOV6: 181 CAWEIVRVEIRRCLYYFYKFTALFRRK 207 (SEQ ID NO.: 63)

ILP-P: 181 CAWEIVRVEIRRCLYYFYKFTALFRRK 207 (SEQ ID NO.: 64)

Where * indicates identity and + indicates similarity.

A NOV6 polypeptide shares sequence homology with many members of the interferon family, including an interferon like-protein precursor. As such, NOV6 represents a new member of the interferon family, and is useful for detecting novel members of the interferon-like family of proteins. NOV6 is useful in determining changes in expression of genes contained within or controlled by the interferon-like protein family. NOV6 satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the treatment of disorders associated with alterations in the expression of members of interferon-like proteins. NOV6 nucleic acids, polypeptides, antibodies, and other compositions of the present invention are useful in the treatment and/or diagnosis of a variety of diseases and pathologies, including by way of nonlimiting example, those involving viral infections, *e.g.* AIDS, viral hepatitis and viral encephalitis. NOV6 is useful for treating cancer, autoimmune diseases, arthritis, multiple sclerosis, diabetes and allergies.

NOV7

A NOV7 sequence according to the invention is a nucleic acid sequence encoding a polypeptide related to the human odorant receptor (OR) family of the G-protein coupled receptor (GPCR) superfamily of proteins. A NOV7 nucleic acid and its encoded polypeptide includes the sequences shown in Table 23. The disclosed nucleic acid (SEQ ID NO:13) is 9,087 nucleotides in length and contains an open reading frame (ORF) that begins with an ATG initiation codon at nucleotides 1-3 and ends with a TGA stop codon at nucleotides 9,085-9,087. The representative ORF encodes a 3,028 amino acid polypeptide (SEQ ID NO:14). The predicted molecular weight of a NOV7 polypeptide is 330,865.9 Da. PSORT analysis predicts a plasma membrane protein with a certainty of 0.6400. SIGNALP analysis predicts a signal peptide cleavage site between positions 20 and 21 of SEQ ID NO: 14.

TABLE 23.

ATGGCGCCGCCGCCGCCGCCGTCGTCGCCGTCGTCGTCGTCCTGGCCGCCGCCGCCGCCCTGCCGGCGA
TGGGGCTGCGAGCGGCCGCCCTGGGAGCCGCGCGTACCCGGCGGGACCCGCGCCTTCGCCCTCCGGCCCGG
CTGTACCTACGCGGTGGGCGCCGCTTGACGCCCCGGCGCCGGGAGCTGCTGGACGTGGGCCGCGAT

GGGCGGCTGGCAGGACGTGCGGCGCTCTCGGGCGGGGGCGCCGCTGCCGCTGCAAGTCCGCTTGGTGG
CCCCGAGTGCCCCGACGGCGTGAGCCGCCGCTGCGGGCGCGCACGACCTTCCCGGCTGCCGAGCCCG
TGCCCGGCTCTGCGGAACCGGTGCCCGGCTCTGCGGGGCGCTCTGCTTCCCGTCCCGGGCGGCTGCGCG
5 GCGCGCAGCATTGGCGCTCGCAGCTCCGACCACCTTACCCGCTGCCGCTGCCCGCGCGCCCCAGGC
CCCGTGTCCCGGCCGTCCCATCTGCTGCCCGCGGGCGGCTCGGTCCGCTGCGTCTGCTGTGCGCCCT
GCGGCGCGCGGCTGGCGCCGTCCGGGTGGGACTGGCGCTGGAGGCCGCCACCGCGGGGACGCCCTCCGCG
TCGCCATCCCCATCGCCGCCCTGCCGCCGAACCTGCCCGAAGCCCGGGCGGGGCCGCGCGACGGGCC
GGCGGGGACAGAGCGGCGAGGGAGCCTGAAGTTCCGATGCCCAACTACCAGGTGGCGTTGTTGAGAA
CGAACCGGCGGGCACCTCATCTCCAGCTGCACGCGCACTACACCATCGAGGGCGAGGAGGAGCGCGTG
10 AGCTATTACATGGAGGGGCTGTTGACGAGCGTCCCAGGGCTACTTCCGAATCGACTCTGCCACGGGCG
CCGTGAGCACGGACAGCGTACTGGACCGCGAGACCAAGGAGACGCACGTCTCAGGGTGAAAGCCGTGGA
CTACAGTACGCGCGCGGCTCGGCCACCACCTACATCACTGTCTTGGTCAAAGACACCAACGACCACAGC
CCGTCTTCGAGCAGTCGGAGTACCGCGAGCGCGTGGCGGAGAACCTGGAGGTGGGCTACGAGGTGCTGA
CCATCCGCGCCAGCGACCGCGACTCGCCCATCAACGCCAACTTGCGTTACCGCGTGTGGGGGCGCGTG
15 GGACGTCTTCCAGCTCAACGAGAGCTCTGGCGTGGTGAGCACACGGGCGGTGCTGGACGGGAGGAGGCG
GGCAGTACCGGCTCTGGTGAGGCGCAACGACCGGGCGCAATCCGGGCCGCTCAGTGCCACGGCCA
CCGTGTACATCGAGGTGGAGGACGAGAACGACAACTACCCCAAGTTCAGCGAGCAGAACTACGTGGTCCA
GGTGCCCGAGGACGTGGGGCTCAACACGGCTGTGCTGCGAGTGCAGGCCACGGACCGGGACCGGGCCAG
AACGCGGCCATTCACTACAGCATCTCAGCGGGAACGTGGCCGGCCAGTCTACCTGCACCTCGTGAGCG
20 GGATCTTGATGTGATCAACCCCTTGGATTTCGAGGATGTCCAGAAATACTCGCTGAGCATTAAAGGCCCA
GGATGGGGGCGGGCCCCGCTCATATTCTCAGGGGTGGTGTCTGTGAGGTGCTGGATGTCAACGAC
AACGAGCCTATCTTTGTGAGCAGCCCCCTTCCAGGCCACGGTGCTGGAGAATGTGCCCTGGGCTACCCCG
TGGTGACATTACGGCGGTGGACGCGGACTCTGGAGAGAACGCCCGGCTGCACTATCGCCTGGTGACAC
GGCTCCACCTTTCTGGGGGCGCGCAGCGTGGGCTAAGAATCCTGCCCCACCCCTGACTTCCCTTC
25 CAGATCCACAACAGCTCCGGTGGATCACAGTGTGTGCCGAGCTGGACCGCGAGGAGGTGGAGCACTACA
GCTTCGGGGTGGAGGCGGTGGACCAGCGCTCGCCCCCATGAGCTCCTCCACAGCGTGTCCATCACGGT
GCTGGACGTGAATGACAACGACCGGTGTTACGCGAGCCCACTACGAGCTTCGTCTGAATGAGGATGCG
GCCGTGGGGAGCAGCGTGTGACCTGACGGCCCGCAGCGTGACGCCAACAGTGTGATTACCTACCAGC
TCACAGGCGGCAACACCCGGAACCGCTTTCAGCTCAGCAGCCAGAGAGGGGGCGGCCTCATCACCTGGC
30 GCTACCTCTGGACTACAAGCAGGAGCAGCAGTACGTGCTGGCGGTGACAGCATCCGACGGCAGACGGTCG
CACACTGCGCATGTCTAATCAACGTCACTGATGCCAACACCCACAGGCCTGTCTTTCAGAGCTCCCAAT
ACACAGTGAGTGTGAGTGGAGCAGGCTGTGGGACCTCCATTGCTACCTCAGTGCCCAACGATGAGGA
CACAGGAGAGTAATGCCCGCATCACCTACGTGATTGAGGACCCCGTGCCGAGTTCCGCTATGACCCGAC
AGTGGCACCATGTACACCATGATGGAGCTGGACTATGAGAACCAGGTGCGCTACACGCTGACCATCATGG
35 CCCAGGACAACGGCATCCCGCAGAAATCAGACACCACCCCTAGAGATCTCATCTCGATGCCAATGA
CAATGCACCCAGTTCTGTGGGATTCTACCAGGGTTCATCTTTGAGGATGCTCCACCTCGACCCAGC
ATCTCCAGGTCTCTGCCACGGACCGGGACTCAGGTCCCAATGGGCGTCTGCTGTACACCTTCCAGGGTG
GGGACGACCGGTGAGGGGACTTCTACATCGAGCCACGCTCCGGTGTGATTGCGACCCAGCGCCGCTGGA
CCGGGAGAATGTGGCGGTGTACAACCTTTGGGCTCTGGCTGTGGATCGGGGCGAGTCCCACTCCCTTAGC
40 GCCTCGGTAGAAATCCAGGTGACCATCTTGGACATTAATGACAATGCCCCATGTTTGAGAAGGACGAAC
TGGAGCTGTTTGTGAGGAGAAACCCAGTGGGTGCGGTGGTGGCAAAGATTCTGTGCTAACGACCTGA
TGAAGGCCCTAATGCCAGATCATGTATCAGATTGTGGAAGGGGACATGCGGCATTTCTTCCAGCTGGAC
CTGCTCAACGGGGACCTGCGTGCCATGGTGGAGCTGGACTTTGAGGTCCGCGGGAGTATGCTGGTGG
TGCAGGCCACGTCCGCTCCGCTGGTGAGCCGAGCCAGGTGCACATCTTCTCGTGGACAGAAATGACAA
45 CCCGCTGTGCTGCCGACTTCCAGATCTCTTCAACAATATGTACCAACAAGTCCAACAGTTTCCCC
ACCGCGGTGATCGGCTGCATCCCGGCCATGACCCGACGTGTGAGACAGCTCAACTACACCTTCGTGC
AGGGCAACGAGCTGCGCTGTTGCTGCTGGACCCCGCCACGGGCGAACTGCAGCTCAGCCGCGACCTGGA
CAACAACCGGCGCGTGGAGGCGCTCATGGAGGTGTCTGTGCTGAGATGGCATCCACAGCGTACGGCC
TTCTGCACCTTGCCTGTACCATCATCAGGACGACATGCTGACCAACAGCATCACTGTCCGCTGGAGA
50 ACATGTCCAGGAGAAGTTCTGTCCCCGCTGCTGGCCCTCTTCTGTGGAGGGGGTGGCCGCGTGTGTC
CACCACCAAGGACGACGTCTTCTGCTTCAACGTCCAGAACGACACCGACGTGAGCTCCAACATCTGAAC
GTGACCTTCTCGGCGCTGCTGCTGGCGGCGTCCGCGGCCAGTTCTTCCCGTGGGAGGACCTGCAGGAGC
AGATCTACCTGAATCGGACGTGCTGACACCATCTCCACGACGCGGTGCTGCCCTTCGACGACAACAT
55 CTGCTGCGCGAGCCCTGCGAGAACTACATGAAGTGCGTGTCCGTTCTGCGATTGACAGCTCCGCGCCC
TTCTCAGCTCCACACCGTGTCTTCCGGGCCATCCACCCATCAACGGCTGCGCTGCCGCTGCCCGC
CCGGCTTACCGGCGACTACTGCGAGACGGAGATCGACCTCTGCTACTCCGACCCGTGCGGCGCCAACGG
CCGCTGCCGCGAGCCGCGAGGGCGGCTACACCTGCGAGTGTCTCGAGGACTTCACTGGAGAGCACTGTGAG
GTGGATGCCCGCTCAGGCCGCTGTGCCAACGGGGTGTGCAAGAACGGGGGACCTGCGTGAACCTGCTCA
TCGGCGGCTTCCACTGCGTGTGCTCTCTGGCGAGTATGAGAGGCCCTACTGTGAGGTGACCACAGGAG
60 TTTCCCGCCCCAGTCTTCTGTCACCTTCCGGGGCTGAGACAGCGCTTCCACTTCACTTCCCTCAGC
TTTGCCACTCAGGAAGGAACGGCTTGCTTCTACAACGGCCGCTTCAATGAGAAGCAGACTTCATCG
CCCTGGAGATCGTGGACGAGCAGGTGCAGCTCACCTTCTGTCAGGTGCAGGCGAGACAACAACGACCGT
GGCACCGAAGGTTCCAGTGTGTGAGTGACGGGCGGTGGCACTCTGTGAGGTGCAGTACTACAACAAG
GTAAGATGGGCCCCACCACTTCCCCCTGGCCCCAGCCCAATATTGGCCACCTGGGCTGCCCATGGGC
65 CGTCCGGGGAAAAGATGGCGTGGTGACAGTGGATGATTGTGACACAACCATGGCTGTGCGCTTTGAA
GGACATCGGGAACTACAGCTGCGCTGCCAGGCGACTCAGACCGGCTCCAAGAAGTCCCTGGATCTGACC

GGCCCTCTACTCCTGGGGGGTGTCCCAACCTGCCAGAAGACTTCCAGTGCACAACCGGCAGTTTCGTGG
GCTGCATGCGGAACCTGTCAGTCGACGGCAAAATGTGGACATGCGCGGATTTCATCGCCAACATGGGCAC
CCGGGAAGGCTGCGCTGCTCGGAGGAACCTTCTGCGATGGGAGGCGGTGTGAGAATGGAGGCACCTGTGTG
AACAGGTGGAATATGTATCTGTGTGAGTGTCCACTCCGATTTCGGCGGGAAGAACTGTGAGCAAGCCATGC
5 CTCACCCCCAGCTCTTCAGCGGTGAGAGCGTCTGTCTGGAGTGACCTGAACATCATCATCTCTGTGCC
CTGGTACCTGGGGCTCATGTTCCGGACCCGGAAGGAGGACAGCGTTCTGATGGAGGCCACCAGTGGTGGG
CCCACCAGCTTTCGCTCCAGATCCTGAACAACCTACCTCCAGTTTGGAGTGTCCCACGGCCCCCTCCGATG
TGGAGTCCGTGATGCTGTCCGGGTTGCGGGTGACCGACGGGGAGTGGCACCACCTGCTGATCGAGCTGAA
GAATGTTAAGGAGGACAGTGAGATGAAGCACCTGGTCACCATGACCTTGGACTATGGGATGGACCAGAAC
10 AAGGCAGATATCGGGGGCATGCTTCCCGGGCTGACGGTAAGGAGCGTGGTGGTTCGGAGGCCCTCTGAAG
ACAAGGTCTCCGTGCGCGTGGATTCCGAGGCTGTCATGACGGGAGTGAGGATGGGGGGGACCCCAACAA
CGTCGCCACCCTGAACATGAACAACGCACTCAAGGTGAGGGTGAAGGACGGCTGTGATGTGGACGACCCC
TGTACCTCGAGCCCCGTGTCACCAATAGCCGCTGCCACGACGCTGGGAGGACTACAGTGCCTGTGTG
ACAAAGGGTACCTTGGATAAACTGTGTGGATGCCTGTACCTGAACCCCTGCGAGAATCGGGGGCCTG
15 CGTGCCTCCCCGGCTCCCCGACGGGCTACGTGTGCGAGTGTGGGCCAGTCACTACGGGCCGTACTGT
GAGAACAACTCGACCTTCCGTGCCCCAGAGGCTGGTGGGGGAACCCGCTGTGTGGACCTGCCACTGTG
CCGTGAGCAAGGCTTTGATCCCGAGTCTTAATAAGAACGCGCAGTGCCAATGCAAGGAGAATTACTA
CAAGCTCCTAGCCCAGGACACCTGTCTGCCCTGCGACTGCTTCCCCCATGGCTCCCACAGCCGCACTTGC
GACATGGCCACCGGGCAGTGTGCCTGCAAGCCGGCGTCATCGGCCGCCAGTGCAACCGCTGCGACAACC
20 CGTTTGGCGAGGTACACACGCTCGGCTGTGAAGTGATCTACATGGCTGTCCCAAAGCATTTGAGGCCGG
CATCTGGTGGCCACAGACCAAGTTCGGGCAGCGGCTGCGGTGCCATGCCCTAAGGGATCCGTTGGAAAT
GCGGTCCGACACTGCGAGCGGGAGAGGCTGCTGCCCCAGAGCTCTTAACTGTACCACTTCCCTCCT
TCGTGGACCTCAGGGCCATGAATGAGAAGCTGAGCCGCAATGAGACGAGGTGGACGGCGCCAGGGCCCT
GCAGCTGGTGAGGGCGCTGCGCAGTGTACACAGCACACGGGCACGCTCTTTGGCAATGACGTGCGCACG
25 GCTTACAGCTGCTGGGCCACGTCTTTCAGCAGGAGAGTGGCAGCAGGGCTTCGACCTGGCAGCCACGC
AGGACGCGGACTTTCACGAGGACGTATCCACTCGGGCAGCGCCCTCCTGGCCCCAGCCACCAGGGCGGC
GTGGGAGCAGATCCAGCGGAGCGAGGGCGGCACGGCAGCTGCTCCGGCGCCTCGAGGGCTACTTCAGC
AACGTGGCAGCAACGTGCGGCGGACGTACCTGCGGCCCTTCGTGATCGTCACCGCCAACATGGTTCTTG
CTGTGACATCTTTGACAAGTTCAACTTTACGGGAGCCAGGGTCCCGGATTTCGACACCATCCATGAAGA
30 GTTCCCCAGGGAGCTGGAGTCTCCTGCTCTCTTCCAGCCGACTTCTTCAGACCACCTGAAGAAAAGAA
GGCCCCCTGCTGAGGGCCGCTGGCCGGAGGACCACCCCGCAGACCACGCGCCCGGGGCTGGCACCGAGA
GGGAGGCCCCGATCAGCAGGCGGAGGCGACACCTGATGACGCTGGCCAGTTCGCCGTCGCTCTGGTCA
CATTTACCGACACCTTGGGCGAGCTCCTGCCGAGCGCTACGACCCCGACCGTTCGACGCTTCCGCT
CACCGGCCCATCATTAATACCCCGATGGTGAGCAGCTGGTGTACAGCGAGGGGGCTCCGCTCCCGAGAC
35 CCTGGAGAGGCCCGTCTGCTGGAGTTTCGCCCTGCTGGAGGTGGAGGAGCGAACAAGCCTGTCTGCGT
GTTCTGGAACCACTCCCTGGCCGTGGTGGGACGGGAGGGTGGTGTGCCCCGGGGCTGCGAGCTCCTGTCC
AGGAACCGGACACATGTCGCTGCCAGTGACGCCACAGCCAGCTTTGCGGTGCTCATGGATATCTCCA
GGCGTGAGAAGCGGGAGGTCTGCTCTGAAGATTGTCACTATGCGCTGTGTCTTGTCACTGGCAGC
CCTGCTGGCTGGCCCTTCGTCTCTGAGCCTGGTCCGATGCTGCGCTCCAACCTGCACAGCATTACAAG
40 CACCTCGCCGTGGCGCTCTTCTCTCTCAGCTGGTGTTCGTGATTGGGATCAACCAGACGGAAAACCGT
TTCTGTGCACAGTGGTTGCCATCCTCCTCCACTACATCTACATGAGCACCTTTGCCTGGACCCCTCGTGGA
GAGCCTGCATGTCTACCGCATGCTGACCGAGGTGCGCAACATCGACACGGGGCCCATGCGGTTCTACTAC
GTCGTGGGCTGGGGCATCCCGCCATTGTACAGGACTGGCGGTGCGCCTGGACCCCGAGGGCTACGGGA
ACCCGACTTCTGCTGGTGTGCTTCAAGACACCCCTGATTGGAGCTTTGCGGGGCCCATCGGAGCTGT
45 TATAATCATCAACACAGTCACTTCTGTCTATCTGCAAAGTTTCTGCAAAGAAAGCACCATTATTAT
GGGAAAAAAGGGATCGTCTCCCTGCTGAGGACCGCATTCCTCCTGTGCTGCTCATCAGCGCCACCTGGC
TGCTGGGGCTGCTGGCTGTGAACCGCATGCACTGAGCTTTCCTACCTCTTCGCCATCTTCAGCGGCTT
ACAGGGCCCCCTTCGTCTCTTTCCTACTGCGTGCTCAACCAGGAGGTCCGGAAGCACCTGAAGGGCGTG
CTCGGCGGGAGGAAGCTGCACCTGGAGGACTCCGCCACACAGGGCCACCTGCTGACGCGCTCCCTCA
50 ACTGCAACACCACCTTCGGTGACGGGCTGACATGCTGCGCACAGACTTGGGCGAGTCCACCGCTCGCT
GGACAGCATCGTCAGGGATGAAGGGATCCAGAAGCTCGGCGTGTCTCTGGGCTGGTGGGGGACGCCAC
GGAGAGCCAGACGCGTCCCTCATGCCAGGAGTGCAAGGATCCCCCTGGCCACGATTCCGACTCAGATA
GCGAGCTGTCCCTGGATGAGCAGAGCAGCTCTACGCTCTCACACTCGTCAGACAGCGAGGACGATGG
GGTGGGAGCTGAGGAAAAATGGGACCCGGCCAGGGCGCGCTCCACAGCACCCCAAGGGGACGCTGTG
55 GCCAACCACTTCCGGCCGGCTGGCCCCAGCAGCCTGGCTGAGAGTGACAGTGAGGACCCCGAGCGGCA
AGCCCCGCTGAAGGTGGAGACCAAGGTGACGCTGGAGCTGCACCGCAGGAGCAGGGCAGTACCGTGG
AGAGTACCCCGCGGACAGGAGAGCGGGGGCGCAGCCAGGCTTGCTAGCAGCCAGCCCCAGAGCAGAGG
AGCATCTTGAATAAAGTCACTACCCGCCCGCTGACGCTGACGGAGCAGACGCTGAAGGGCCGGC
TCCGGGAGAAAGCTGGCCGACTGTGAGCAGAGCCCACTCTCGCGCACGCTCTTCCCTGGGCTCTGGCGG
60 CCGGACTGCGCCATCAAGTCAAGAGCCCTGGAGGGAGCGGGGCGTGACCACTCAACGGGGTGGCC
ATGAATGTGCGCACTGGGAGCGCCAGGCCGATGGCTCCGACTCTGAGAAACCGTGA (SEQ ID NO.: 13)

MAPPPPPVLPVLLLLAAAAALPAMGLRAAAWEPRVPGGTRAFALRPGCTYAVGAACTPRAPRELLDVGRDGLRAGR
RRVSGAGRPLPLQVRLVARSAPTALSRRLRARTHLPGCGARARLCGTGARLCGALCFVPVGGCAAQHSALAAPTT

LPACRCPPRRPRCPGRPICLPFGGSSVRLRLCALRRAAGAVRVGLALEAATAGTPSASPSPSPPLPPNLPEARAG
 PARRARRGTSGRGLSKFPMQYVALFENEPAGTLILQLHAHYTTIEGEEERSYMEGLFDERSGYFRIDSATGA
 VSTDVLDRETKETHVLRVKAVDYSTPPRSATTYITVLVKDNDHSPVFEQSEYRERVRENLEVGVEVLTIRASDR
 DSPINANLRYRVLGGAWDVFQINNESSGVVSTRAVLDRREEAAEYQLLVEANDQGRNPGPLSATATVYIEVEDENDNY
 5 PQFSEQNYVVQVPEDVGLNTAVLRVQATDRDQGGQNAIHYISILSGNVAGQFYLHSLSGILDVINPLDFEDVQKYSL
 SIKAQDGGRRPPLINSSGVVSQVLDVNDNEPIFVSSPFQATVLENVPLGYFVVHIIQAVDADSDGENARLHYRLVDTA
 STFLGGGSAGPKNPAPTDPDFPFIHNSSGWITVCAELDREEVEHYSGVEAVDHGSPPMSSSTSISITVLVDVNDND
 PVFTQPTYELRLNEDAAGSSVLTQLQARDRDANSVITYQLTGGNTRNRFALSSQSGGLITLALPLDYKQEQQYVL
 10 AVTASDGRSTRHTAHVLINVTANTRHPVFQSSHYTVSVSEDRPVGTSIATLSANDEDTGENARITYVIQDPVPQFR
 IDPDSGTMYTMMELDYENQVAYTLTITMAQDNGIPQKSDTTTLEILILDANDNAPQFLWDFYQGSIFEDAPPSTISIL
 QVSATDRDSGPNRLLYTFQGGDDGDGDFYIEPTSGVIRTQRRLDRENVAVYNLWALAVDRGSPPLSASVEIQVT
 ILDINDNAPMFEKDELELFVEENNPVGSVVAKIRANDPDEGPNQAQIMYQIVEGDMRHHFFQLDLLNGDLRAMVELDF
 EVRREYVLVVQATSAPLVSRATVHILLVDQNDNPPVLPDFQILFNNYVTNKSNSFPTGVIGCIPAHDPDVSDSLNY
 15 TFVQGNELRLLLDPATGELQLSRDLNDRPLEALMEVSVSADGIHSVTAFTCLRVTIITDDMLTNSITVRLNMS
 QEKFLSPLLLALFVEGVA AVLSTTKDDVFVFNVDNDTVSSNILNVTFSSALLPGGVVRGQFFPSEDLEQEIYLNRTLL
 TTISTQRVLPFDNICLREPCENYMKCVSVLRFDSAPFLSSITVLRPIHPINGLRRCRCPGFTGDYCEIDEIDLC
 YSDPCGANGRCRSREGGYTCECFEDFTGEHCEVDARSGRCANGVCKNGGTCVNLLIGGFHCVCPPGEYERPYCEVT
 TRSFPPQSFVTFRGLRQRFHFTISLTFATQERNGLLYNGRFNEKHDFIALEIVDEQVQLTFSAGAGETTTTVPAPK
 VPSSGVS DGRWHSVQVQYYNKVRWAPPLPPGPQPNIGHLGLPHGPSGEKMAVVTVDDCDTTMAVRFKDIGNYSCAA
 20 QGTQTGSKKSLDLTGPLLLGGVFNLPEDFPVHNRQFVGCNRNLSVDGKNVDMAGFIANNGTREGCAARRNFCDGRR
 CQNGGTCVNRWNMYLCECPLRFGGKNCQAMPHPQLFSGESVSVWSDLNIIISVPWYLGMLFRTRKEDSVLMEATS
 GGPTSFRLQILNNYLQFEVSHGPSDVESVMSGLRVTDGEGWHLLIELKNVKEDSEMHLVTMTLDYGMQDNKADI
 GGMPLGLTVRSVVVGASEDKVSVRRGFRGCMQGVMMGGTPTNVATLNMNNAKVRVKDGDVDDPCTSSPCPPNS
 RCHDAWEDYSCVCDKGYLGINCVDACHLNPENMGACVRS PGSPQGYVCEGPHSHYGPYCENKLDLPCPRGWGNGP
 25 VCGPCHCAVSKGFDPDCNKTNGQCCKENYKLLAQDTCLPCDCFPHGSHSRTCDMATGQCACKPGVIGRQCNRCD
 NPFAEVTTLGCEVIYNGCPKAFAEAGIWWPQTKFGQPAAVPCPKGSVGNVHRHCSGEKGLWLPPELFNCTTISFVDLR
 AMNEKLSRNETQVDGARALQLVRALRSATQHTGTLEGNDRVTAQYLLGHVQLQESWQQGFDLAATQADDFHEDVH
 SGAALLAPATRAAWEQIQRSEGGTAQLLRLEGYFSNVARNVRRTYLRPFVIVTANMVLAVDIFDKENFTGARVPR
 FDTIHEEFPRELESSVSFPADFFRPPEEKEGPLL RPAGRRTTPQTRPGPGTEREAPISRRRRHPDDAGQFAVALV
 30 IIRYTLGQLLPERYDPDRSLRLPHRPIINTPMVSTLVYSEGAPLPRPLERPVLEFALLEVEERTKPVCFWNHS
 LAVGGTGGWSARGCELLSRNRTHVACQCSHTASFAVLMDISRRENGEVLPLKIVTYAAVSLSLAALLVAFVLLSLV
 RMLRSNLHSIHKHLAVALFSLQVLFVIGINQTNPFCLTVVAILLHYIYMSTFAWTLVESLHVYRMLTEVRNIDTG
 PMRFYVVVGWGIPIAVTGLAVGLDPQGYGNPDEFWLSLQDTLIWSFAGPIGAVIIINTVTSVLAKVSCQRKHYY
 35 GKKGI VSLRLTAFLLLLLISATWLLGLLAVNRDALSFHYLFAIFSGLQGPVLLFHCVLNQEVKRLKGLVGGGRKL
 HLED SATTRATLLTRSLNCNTTFGDGPDMLRTDLGESTASLDSIVRDEGIQKLGVS SGLVRGSHGEPDASLMPRSC
 KDPPGHDS DSDSELSLDEQSSSYASSHSSDSEDDGVGAEEKWDPAARGAVHSTPKGDAVANHVPAAGWPDQSLAESDS
 EDPSGKPRKLVETKVSVELHREEQSGHRGEYPPDQESGGAARLASSQPPEQRSILKNKVTPPPLTLTEQTLKGR
 REKLADCEQSPTSSRTSSLSGSGGPDCAITVKSPPGREPRDHLNGVAMNVRTGSAQADGSDSEKP (SEQ ID
 NO.: 14)

40

The OR family of the GPCR superfamily is a group of related proteins specifically
 located at the ciliated surface of olfactory sensory neurons in the nasal epithelium and are
 involved in the initial steps of the olfactory signal transduction cascade. NOV7 nucleic acids,
 polypeptides, antibodies, and other compositions of the present invention can be used to detect
 45 nasal epithelial neuronal tissue.

The NOV7 nucleic acid has a high degree of homology (99% identity) with human
 chromosome 22q13.2-13.33, including the uncharacterized genomic clone RP5-1163J1 (CHR
 22; GenBank Accession No.: HS1163J1), as shown in Table 24. The NOV7 nucleic acid also
 has a high degree of homology (99% identity) with human chromosome 22q13.31-13.33,
 50 including the uncharacterized genomic clone RP3-439F8 (CHR 22; GenBank Accession No.:
 HS439F8), as shown in Table 25. The NOV7 polypeptide has homology (approximately 80%
 identity, 87% similarity) to a member of the mouse Celsr family of evolutionarily conserved

seven-pass transmembrane receptors expressed during embryogenesis (Celsr; EMBL Accession No.:T14119), as is shown in Table 26. Overall amino acid sequence identity within the mammalian OR family ranges from 45% to >80%. OR genes that are 80% or more identical to each other at the amino acid level are considered by convention to belong to the same subfamily. See *Dryer and Berghard, Trends in Pharmacological Sciences*, 1999, 20:413. Therefore, NOV7 and the mouse Celsr protein are in the same subfamily. OR proteins have seven transmembrane α -helices separated by three extracellular and three cytoplasmic loops, with an extracellular amino-terminus and a cytoplasmic carboxy-terminus. Multiple sequence alignment suggests that the ligand-binding domain of the ORs is between the second and sixth transmembrane domains.

TABLE 24.

15	NOV7:	8594	CCAAAGGGGACGCTGTGGCCAACCACGTTCCGGCCGGCTGGCCCGACCAGAGCCTGGCTG	8653
	CHR22:	65389	CCTCAGGGGACGCTGTGGCCAACCACGTTCCGGCCGGCTGGCCCGACCAGAGCCTGGCTG	65448
20	NOV7:	8654	AGAGTGACAGTGAGGACCCAGCGGCAAGCCCGCCTGAAGGTGGAGACCAAGGTCAGCG	8713
	CHR22:	65449	AGAGTGACAGTGAGGACCCAGCGGCAAGCCCGCCTGAAGGTGGAGACCAAGGTCAGCG	65508
25	NOV7:	8714	TGGAGCTGCACCGCGAGGAGCAGGGCAGTCACCGTGGAGAGTACCCCGGACCAGGAGA	8773
	CHR22:	65509	TGGAGCTGCACCGCGAGGAGCAGGGCAGTCACCGTGGAGAGTACCCCGGACCAGGAGA	65568
30	NOV7:	8774	GCGGGGGCGCAGCCAGGCTTGCTAGCAGCCAGCCCCAGAGCAGAGGA	8821 (SEQ ID NO.: 65)
	CHR22:	65569	GCGGGGGCGCAGCCAGGCTTGCTAGCAGCCAGCCCCAGAGCAGAGGA	65616 (SEQ ID NO.: 66)

TABLE 25.

35	NOV7:	1	ATGGCGCGCGCGCGCGCGCGCTGCTGCCCCTGCTGCTGCTCCTGGCCGCGCGCGCGCC	60
	CHR22:	111063	ATGGCGCGCGCGCGCGCGCGCTGCTGCCCCTGCTGCTGCTCCTGGCCGCGCGCGCGCC	111122
40	NOV7:	61	CTGCCGCGATGGGGCTGCGAGCGGCGCGCTGGGAGCCGCGCTACCCGGCGGGACCCGC	120
	CHR22:	111123	CTGCCGCGATGGGGCTGCGAGCGGCGCGCTGGGAGCCGCGCTACCCGGCGGGACCCGC	111182
45	NOV7:	121	GCCTTCGCCCTCCGGCCCGGCTGTACCTACGCGGTGGGCGCGCTTGACGCCCCGGGCG	180
	CHR22:	111183	GCCTTCGCCCTCCGGCCCGGCTGTACCTACGCGGTGGGCGCGCTTGACGCCCCGGGCG	111242
50	NOV7:	181	CCGCGGGAGCTGCTGGACGTGGGCGCGATGGGCGGCTGGCAGGACGTCGGCGGCTCTCG	240
	CHR22:	111243	CCGCGGGAGCTGCTGGACGTGGGCGCGATGGGCGGCTGGCAGGACGTCGGCGGCTCTCG	111302
55	NOV7:	241	GGCGCGGGCGCCCGCTGCCGCTGCAAGTCCGCTTGGTGGCCCGCAGTGCCCCGACGGCG	300
	CHR22:	111303	GGCGCGGGCGCCCGCTGCCGCTGCAAGTCCGCTTGGTGGCCCGCAGTGCCCCGACGGCG	111362
	NOV7:	301	CTGAGCCGCGCGCTGCGGGCGCGCACGACCTTCCCGGCTGCGGAGCCCGTGCCCGGCTC	360
	CHR22:	111363	CTGAGCCGCGCGCTGCGGGCGCGCACGACCTTCCCGGCTGCGGAGCCCGTGCCCGGCTC	111422

26

|||||
CHR22: 112443 GAGCAGAACTACGTGGTCCAGGTGCCGAGGACGTGGGGCTCAACACGGCTGTGCTGCGA 112502
5 NOV7: 1441 GTGCAGGCCACGGACCGGGACCAGGGCCAGAACGCGGCCATTCACTACAGCATCCTCAGC 1500
|||||
CHR22: 112503 GTGCAGGCCACGGACCGGGACCAGGGCCAGAACGCGGCCATTCACTACAGCATCCTCAGC 112562
NOV7: 1501 GGGAACTGGCCGGCCAGTTCTACCTGCACTCGCTGAGCGGGATCCTGGATGTGATCAAC 1560
|||||
10 CHR22: 112563 GGGAACTGGCCGGCCAGTTCTACCTGCACTCGCTGAGCGGGATCCTGGATGTGATCAAC 112622
NOV7: 1561 CCCTTGATTTTCGAGGATGTCCAGAAATACTCGCTGAGCATTAAAGGCCAGGATGGGGC 1620
|||||
15 CHR22: 112623 CCCTTGATTTTCGAGGATGTCCAGAAATACTCGCTGAGCATTAAAGGCCAGGATGGGGC 112682
NOV7: 1621 CGGCCCCGCTCATCAATTCTTCAGGGGTGGTGTCTGTGCAGGTGCTGGATGTCAACGAC 1680
|||||
CHR22: 112683 CGGCCCCGCTCATCAATTCTTCAGGGGTGGTGTCTGTGCAGGTGCTGGATGTCAACGAC 112742
20 NOV7: 1681 AACGAGCCTATCTTTGTGAGCAGCCCTTCCAGGCCACGGTGTGGAGAATGTGCCCTG 1740
|||||
CHR22: 112743 AACGAGCCTATCTTTGTGAGCAGCCCTTCCAGGCCACGGTGTGGAGAATGTGCCCTG 112802
NOV7: 1741 GGCTACCCCGTGGTGACATTTCAGCGGTGGACGCGGACTCTGGAGAGAACGCCCCGCTG 1800
|||||
25 CHR22: 112803 GGCTACCCCGTGGTGACATTTCAGCGGTGGACGCGGACTCTGGAGAGAACGCCCCGCTG 112862
NOV7: 1801 CACTATCGCCTGGTGGACACGGCCTCCACCTTTCTGGGGGGCGGCAGCGCTGGGCCTAAG 1860
|||||
30 CHR22: 112863 CACTATCGCCTGGTGGACACGGCCTCCACCTTTCTGGGGGGCGGCAGCGCTGGGCCTAAG 112922
NOV7: 1861 AATCCTGCCCCACCCCTGACTTCCCTTCCAGATCCACAACAGCTCCGGTTGGATCACA 1920
|||||
35 CHR22: 112923 AATCCTGCCCCACCCCTGACTTCCCTTCCAGATCCACAACAGCTCCGGTTGGATCACA 112982
NOV7: 1921 GTGTGTGCCGAGCTGGACCGCGAGGAGGTGGAGCACTACAGCTTCGGGGTGGAGGCGGTG 1980
|||||
CHR22: 112983 GTGTGTGCCGAGCTGGACCGCGAGGAGGTGGAGCACTACAGCTTCGGGGTGGAGGCGGTG 113042
40 NOV7: 1981 GACCACGGCTCGCCCCCATGAGCTCCTCCACCAGCGTGTCCATCACGGTGTGGACGTG 2040
|||||
CHR22: 113043 GACCACGGCTCGCCCCCATGAGCTCCTCCACCAGCGTGTCCATCACGGTGTGGACGTG 113102
NOV7: 2041 AATGACAACGACCCGGTGTTCACGCAGCCACCTACGAGCTTCGTCTGAATGAGGATGCG 2100
|||||
45 CHR22: 113103 AATGACAACGACCCGGTGTTCACGCAGCCACCTACGAGCTTCGTCTGAATGAGGATGCG 113162
NOV7: 2101 GCCGTGGGGAGCAGCGTGTGACCTGCAGGCCGCGACCGTGACGCCAACAGTGTGATT 2160
|||||
50 CHR22: 113163 GCCGTGGGGAGCAGCGTGTGACCTGCAGGCCGCGACCGTGACGCCAACAGTGTGATT 113222
NOV7: 2161 ACCTACCAGCTCACAGGCGGCAACACCCGGAACCGCTTTGCACTCAGCAGCCAGAGAGGG 2220
|||||
55 CHR22: 113223 ACCTACCAGCTCACAGGCGGCAACACCCGGAACCGCTTTGCACTCAGCAGCCAGAGAGGG 113282
NOV7: 2221 GGCGGCCTCATCACCTGGCGCTACCTCTGGACTACAAGCAGGAGCAGCAGTACGTGCTG 2280
|||||
CHR22: 113283 GGCGGCCTCATCACCTGGCGCTACCTCTGGACTACAAGCAGGAGCAGCAGTACGTGCTG 113342
60 NOV7: 2281 GCGGTGACAGCATCCGACGGCACACGGTCGCACACTGCGCATGTCTAATCAACGTCACT 2340
|||||
CHR22: 113343 GCGGTGACAGCATCCGACGGCACACGGTCGCACACTGCGCATGTCTAATCAACGTCACT 113402
NOV7: 2341 GATGCCAACACCCACAGGCCTGTCTTTAGAGCTCCCATACACAGTGAGTGTCACTGAG 2400
|||||
65 CHR22: 113403 GATGCCAACACCCACAGGCCTGTCTTTAGAGCTCCCATACACAGTGAGTGTCACTGAG 113462
NOV7: 2401 GACAGGCCTGTGGGCACCTCCATTGCTACCTCAGTGCCAACGATGAGGACACAGGAGAG 2460
|||||
70 CHR22: 113463 GACAGGCCTGTGGGCACCTCCATTGCTACCTCAGTGCCAACGATGAGGACACAGGAGAG 113522
NOV7: 2461 AATGCCCGCATCACCTACGTGATTAGGACCCCGTGCCGAGTTCCGATGACCCCGAC 2520

|||||
CHR22: 113523 AATGCCCGCATCACCTACGTGATTGAGGACCCCGTGCCGCGAGTTCCGCATTGACCCCGAC 113582
5 NOV7: 2521 AGTGGCACCATGTACACCATGATGGAGCTGGACTATGAGAACCAGGTCGCCTACACGCTG 2580
|||||
CHR22: 113583 AGTGGCACCATGTACACCATGATGGAGCTGGACTATGAGAACCAGGTCGCCTACACGCTG 113642
NOV7: 2581 ACCATCATGGCCAGGACAAACGGCATCCCGCAGAAATCAGACACCACCACCTAGAGATC 2640
|||||
10 CHR22: 113643 ACCATCATGGCCAGGACAAACGGCATCCCGCAGAAATCAGACACCACCACCTAGAGATC 113702
NOV7: 2641 CTCATCCTCGATGCCAATGACAATGCACCCAGTTCCTGTGGGATTTCTACCAGGGTTCC 2700
|||||
15 CHR22: 113703 CTCATCCTCGATGCCAATGACAATGCACCCAGTTCCTGTGGGATTTCTACCAGGGTTCC 113762
NOV7: 2701 ATCTTTGAGGATGCTCCACCTCGACCAGCATCCTCCAGGTCTCTGCCACGGACCGGGAC 2760
|||||
CHR22: 113763 ATCTTTGAGGATGCTCCACCTCGACCAGCATCCTCCAGGTCTCTGCCACGGACCGGGAC 113822
20 NOV7: 2761 TCAGGTCCCAATGGGCGTCTGCTGTACACCTTCAGGGTGGGGACGACGGCGATGGGGAC 2820
|||||
CHR22: 113823 TCAGGTCCCAATGGGCGTCTGCTGTACACCTTCAGGGTGGGGACGACGGCGATGGGGAC 113882
NOV7: 2821 TTCTACATCGAGCCACGTCGGGTGTGATTGCGACCCAGCGCCGGCTGGACCGGGAGAAT 2880
|||||
25 CHR22: 113883 TTCTACATCGAGCCACGTCGGGTGTGATTGCGACCCAGCGCCGGCTGGACCGGGAGAAT 113942
NOV7: 2881 GTGGCCGTGTACAACTTTGGGCTCTGGCTGTGGATCGGGGCAGTCCCACTCCCCTTAGC 2940
|||||
30 CHR22: 113943 GTGGCCGTGTACAACTTTGGGCTCTGGCTGTGGATCGGGGCAGTCCCACTCCCCTTAGC 114002
NOV7: 2941 GCCTCGGTAGAAATCCAGGTGACCATCTTGGACATTAATGACAATGCCCCCATGTTTGAG 3000
|||||
35 CHR22: 114003 GCCTCGGTAGAAATCCAGGTGACCATCTTGGACATTAATGACAATGCCCCCATGTTTGAG 114062
NOV7: 3001 AAGGACGAACTGGAGCTGTTTGTGAGGAGAAACAACCCAGTGGGGTCCGGTGGTGGCAAAG 3060
|||||
CHR22: 114063 AAGGACGAACTGGAGCTGTTTGTGAGGAGAAACAACCCAGTGGGGTCCGGTGGTGGCAAAG 114122
40 NOV7: 3061 ATTCTGTCTAACGACCCTGATGAAGGCCCTAATGCCAGATCATGTATCAGATTGTGGAA 3120
|||||
CHR22: 114123 ATTCTGTCTAACGACCCTGATGAAGGCCCTAATGCCAGATCATGTATCAGATTGTGGAA 114182
NOV7: 3121 GGGGACATGCGGCATTTCTTCCAGCTGGACCTGCTCAACGGGGACCTGCGTGCCATGGTG 3180
|||||
45 CHR22: 114183 GGGGACATGCGGCATTTCTTCCAGCTGGACCTGCTCAACGGGGACCTGCGTGCCATGGTG 114242
NOV7: 3181 GAGCTGGACTTTGAGGTCCGGCGGGAGTATGTGCTGGTGGTGCAGGCCACGTCGGCTCCG 3240
|||||
50 CHR22: 114243 GAGCTGGACTTTGAGGTCCGGCGGGAGTATGTGCTGGTGGTGCAGGCCACGTCGGCTCCG 114302
NOV7: 3241 CTGGTGAGCCGAGCCACGGTGACATCCTTCTCGTGGACCAGAATGACAACCCGCTGTG 3300
|||||
55 CHR22: 114303 CTGGTGAGCCGAGCCACGGTGACATCCTTCTCGTGGACCAGAATGACAACCCGCTGTG 114362
NOV7: 3301 CTGCCCAGCTTCCAGATCCTCTTCAACAATATGTACCAACAAGTCCAACAGTTTCCCC 3360
|||||
CHR22: 114363 CTGCCCAGCTTCCAGATCCTCTTCAACAATATGTACCAACAAGTCCAACAGTTTCCCC 114422
60 NOV7: 3361 ACCGGCGTGATCGGCTGCATCCCGGCCATGACCCGACGTGTCAGACAGCTCAACTAC 3420
|||||
CHR22: 114423 ACCGGCGTGATCGGCTGCATCCCGGCCATGACCCGACGTGTCAGACAGCTCAACTAC 114482
NOV7: 3421 ACCTTCGTGCAGGGCAACGAGCTGCGCCTGTTGCTGCTGGACCCGCCACGGGCGAACTG 3480
|||||
65 CHR22: 114483 ACCTTCGTGCAGGGCAACGAGCTGCGCCTGTTGCTGCTGGACCCGCCACGGGCGAACTG 114542
NOV7: 3481 CAGCTCAGCCGCGACCTGGACAACAACCGGCCGCTGGAGGCGCTCATGGAGGTGTCTGTG 3540
|||||
70 CHR22: 114543 CAGCTCAGCCGCGACCTGGACAACAACCGGCCGCTGGAGGCGCTCATGGAGGTGTCTGTG 114602
NOV7: 3541 TCTGCAGA-TGGC 3552 (SEQ ID NO.: 67)

1111 11 1111
 CHR22: 114603 TCTGGTGAGTGGC 114615 (SEQ ID NO.: 68)

5

TABLE 26.

10	NOV7:	1	MAPPPPPVLPVLLLLLAAAAALPAMGLRAAAWEPRVPGGTRAFALRPGCTYAVGAACTPRA	60
	Celsr:	1	MAPSSPRVLPALVLLAAAA-LPALELGAAAWELRVPGGARAFALGPGWSYRLD---TTRT	56
15	NOV7:	61	PRELLDVGRDGRLAGRRRVSGAG-----RPLPLQVRLVARSAPTALSRRLRARHL	111
	Celsr:	57	PRELLDVSREGPAAGRRLGLGACTLGCARLAGRLLPLQVRLVARGAPTAPSLVLRARAYG	116
20	NOV7:	112	PGCGARA-RLCGTGARLCGALCFVPG-GCAAQHSALAAPTTLACRCPPRPRPRCPGR	169
	Celsr:	117	ARCGVRLRRSARGAELRSPAVRSVPGLGDALCFPAAGGGAASLTSVLEAITNFPACSCP	176
25	NOV7:	170	PICLPPGGSVRLRLLCALRRAAGAVRV---GLALEA-----ATAGTPSASPSPP	217
	Celsr:	177	PVA---GTGCRGPICLRPGGSAELRLVCALGRAAGAVWVELVIQATSGTPSESPSVSPS	233
30	NOV7:	218	LPPNLPEARAGPARRARRGTSGRGLKFPMPNYQVALFENEPAGTLILQLHAHYTIEGEE	277
	Celsr:	234	LL-NLSQPRAGVVRRSRRGTGSSTSPQFPLPSYQVSVPENEPAGTAVIELRAHDPDEGDA	292
35	NOV7:	278	ERVSYYMEGLFDESRGYFRIDSATGAVSTDSVLDRETKETHVLRVKAVDYSTPPRSATT	337
	Celsr:	293	GRLSYQMEALFDESRNGYFLIDAATGAVTTARSLDRETKDTHVLKVSVDHGSPPRSAAT	352
40	NOV7:	338	YITVLVKDNDHSPVFEQSEYRERVRENLEVGYEVLTIASDRDSPINANLRYRVLGGA	397
	Celsr:	353	YLTVTVSDNDHSPVFEQSEYRERIRENLEVGYEVLTIATDGDAPSANMRYRLLEGAG	412
45	NOV7:	398	DVFQLNESSGVVSTRAVLDRREEAAEYQLLVEANDQGRNPGPLSATATVYIEVEDENDNYP	457
	Celsr:	413	GVFEIDARSGVVTRAVVDREEAAEYQLLVEANDQGRNPGPLSASATVHIVVEDENDNYP	472
50	NOV7:	458	QFSEQNYVVQVPEDVGLNTAVLRVQATDRDQGQNAAIHYSILSGNVAGQFYLHSLSGILD	517
	Celsr:	473	QFSEKRYVVQVPEDVAVNTAVLRVQATDRDQGQNAAIHYSIVSGNLKGQFYLHSLSGSLD	532
55	NOV7:	518	VINPLDFEDVQKYSLSIKAQDGRPPLINSSGVSVQVLDVNDNEPIFVSSPFQATVLEN	577
	Celsr:	533	VINPLDFEAIREYTLRIKAQDGRPPLINSSGLVSVQVLDVNDNAPIFVSSPFQAAVLEN	592

PCT/US01/04828

30

PCT/US01/04828

31

NOV7: 1838 VGGASEDKVSVRRGFRGCMQGVRMGGTPTNVATLNMNNAKVRVKDGCDDPCTSSPCP
1897
+** +***** ***** * **+*****+*****+*****+*****
5 Celsr: 1838 IGGVTEDKVSVRHGFRGCMQGVRMGETSTNIATLNMNDALKVRVKDGCDDVEDPCASSPCP
1897
NOV7: 1898 PNSRCHDAWEDYSCVCDKGYLGINCVDACHLNPENMGACVRSPPGSPQGYVCECGPSHYG
1957
*+ * * *+ ***+*** * ***** **++ ***** +*** *****
10 Celsr: 1898 PHRPCRDTWDSYSCICDRGYFGKKCVDACLLNPCKHVAACVRSNPTRGYSCECGPGHYG
1957
NOV7: 1958 PYCENKLDLPCPRGWGNPVCGPCHCAVSKGFDPCNKTNGQCQCKENYKLLAQDTCLP
2017
*****+*****+*****+*****+*****+*****+*****+*****
15 Celsr: 1958 QYCENKVDLPCPKGWGNPVCGPCHCAVSKGFDPCNKTNGQCQCKENYKPPAQDACLP
2017
NOV7: 2018 CDCFPHGSHSRTCDMATGQCACKPGVIGRQCNRCNPF AEVTTLGCEVIYNGCPKAFEAG
2077
***** ***** ** *****+*****+*****+*****
Celsr: 2018 CDCFPHGSHSRACDMGTGQCACKPGVIGRQCNRCNPF AEVTSLGCEVIYNGCPRAFEAG
2077
25 NOV7: 2078 IWWPQTKFGQPAAVPCPKGSVGNVHRHCSGEKGWLPPELFNCTTISFVDLRAMNEKLSRN
2137
*****+*****+*****+*****+*****+*****+*****+*****
Celsr: 2078 IWWPQTKFGQPAAVPCPKGSVGNVHRHCSGEKGWLPPELFNCTSGSFVDLKALNEKLNRN
2137
30 NOV7: 2138 ETQVDGARALQLVRALRSATQHTGTLEGNVVRTAYQLLGHVLQHESWQQGFDLAATQDAD
2197
+ *+*+* +***+*** ***** +***** *****+***+
35 Celsr: 2138 ETRMDGNRSLRLAKALRNATQGNSTLEGNVVRTAYQLLARILQHESRQQGFDLAATREAN
2197
NOV7: 2198 FHEDVIHSGSALLAPATRAAWEQIQRSEGGAQLLRRLEGYFSNVARNVRRTYLRPFVIV
2257
*****+*****+*****+*****+*****+*****+*****+*****
40 Celsr: 2198 FHEDVVHTGSALLAPATEASWEQIQRSEAGAAQLLRHFEAYFSNVARNVKRTYLRPFVIV
2257
NOV7: 6772 TANMVLAVDIFDKFNFTGARVPREDTIEEFPRELESSVSFPADFFRPPEEKEGPLLRPA
6951
*****+*****+*****+*****+*****+*****+*****+*****
45 Celsr: 2258 TANMILAVDIFDKLNFTGAQVPRFEDIQEELPRELESSVSFPADTFKPPEKKEGPVVRT
2317
50 NOV7: 2318 GRRTTPQTTRPGPGTEREAPISRRRRHPDDAGQFAVALVVIYRTLGLQLPERYDPDRSL
2377
***** * +* * * *****+*****+*****+*****+*****
55 Celsr: 2318 NRRTTPLTAQPEPRAERETSSRRRRHPDEPGQFAVALVVIYRTLGLQLPEHYDPDRSL
2377
NOV7: 2378 RLPHRPIINTPMVSTLVYSEGAPLPRPLERPVLVEFALLEVEERTKPVCFWNHSLAVGG
2437
++*****+*** +***** ** *+***+*****+*** **+*****+*****
60 Celsr: 2378 RLPNRPVINTPVVSAMVYSEGTPLPSSLQRPILVEFSLETEERSKPVCFWNHSLDTGG
2437
NOV7: 2438 TGGWSARGCELLSRNRTHVACQCSHTASFAVLMDISRRENGVLPLKIVTYAAVSLSLAA
2497
65 *****+*****+*****+*****+*****+*****+*****+*****

Celsr: 2438 TGGWSAKGCELLSRNRTHVTCQCSHSASCAVLMDISRREHGEVLPLKIITYAALSLSLVA
 2497
 5 NOV7: 2498 LLVAFVLLSLVRMLRSNLHSIHKHLAVALFLSQLVFFVIGINQ TENPFLCTVVAILLHYIY
 2557
 *****+* *** **++*****+
 Celsr: 2498 LLVAFVLLSLVRTLRNLHSIHKNLIAALFFSQLIFMVGINQ TENPFLCTVVAILLHYVS
 2557
 10 NOV7: 2558 MSTFAWTLVESLHVYRMLTEVRNIDTGPMRFYVVGWGIPAIVTGLAVGLDPQGYGNPDF
 2617
 * *****+*****+*****+*****+
 Celsr: 2558 MGTFAWTLVENLHVYRMLTEVRNIDTGPMRFYHVVGWGIPAIVTGLAVGLDPQGYGNPDF
 2617
 15 NOV7: 2618 CWLSLQDTLIWSFAGPIGAVIIINTVTSVLSAKVSCQRKHHYYGKKGIVSLRТАFLLLL
 2677
 *****+* ***** ***** ++**+*****
 20 Celsr: 2618 CWLSLQDTLIWSFAGPVGTVIIINTVIFVLSAKVSCQRKHHYYERKGVVSMRТАFLLLL
 2677
 NOV7: 2678 LISATWLLGLLAVNRDALSFHYLFAIFSGLQGPVLLFHCVLNQEV RKHLKGVLGGRKLH
 2737
 ++*** * ***** ** *** ***** ++*****+ ** +**
 25 Celsr: 2678 LVTATWLLGLLAVNSDTLSFHYLFAAFSCLQGI FVLLFHCVAHREVRKHLRAVLAGKKLQ
 2737
 NOV7: 2738 LEDSATTRATLLTRSLNCNTTFGDGPDMLRTDLGESTASLDSIVRDEGIQKLGVS SGLVR
 2797
 30 **++***** ***** *+ ***** ***** *****+*** ***** *
 Celsr: 2738 LDDSATTRATLLTRSLNCNTTYSEGPDMLRTALGESTASLDSTTRDEGVQKLSVSSGP AR
 2797
 NOV7: 2798 GSHGEPDASIMPRSCDKPPGHDSDSSELSLDEQSSSYASSHSSDSEDDGVGAEEKWDPA
 2857
 35 **++***** * +**+ * ***** ***** *****+***** **++**+
 Celsr: 2798 GNHGEPTSFI PRNSKKAHGPDSDSELSLDEHSSSYASSHTSDSEDDGGEAEDKWNPA
 2857
 NOV7: 2858 RGAVHSTPKGD AVANHV PAGWPDQSLAESDSEDPGKPR LKVETKVSVELHREEQGSHRG
 2917
 * ***** **++*****+*** *****+ +* ***** *****+ **++ *
 40 Celsr: 2858 GGAHSTPKADALANHV PAGWPDES LAGSDSEELDTEPHLKVETKVSVELHREQAGNHCG
 2917
 45 NOV7: 2918 EYPPDQESGGAAR---LASSQPPEQRS-ILKNKV TYPPLTLTEQTLKGR LREKLADCEQ
 2975
 + * * * * * + * * * * * ***** * * * * *
 50 Celsr: 2918 DRPSDPESGVLAKPVAVLSSQPQEQRKGILKNKV TYPPLP--EQPLKSRLREKLADCEQ
 2975
 NOV7: 2976 SPTSSRTSSLGSG---GPDCAITVKSPGREPGRDHLNGVAMNVRTGSAQADGSDSEKP
 3034 SEQ ID NO.: 69)
 ***** ***** ** **++** *****+*****+*****
 55 Celsr: 2976 SPTSSRTSSLGSGDGVHATDCVITIKTPRREP GREHLNGVAMNVRTGSAQANGSDSEKP
 3034 (SEQ ID NO.: 70)

The OR family of the GPCR superfamily is a group of related proteins located at the
 ciliated surface of olfactory sensory neurons in the nasal epithelium. The OR family is
 60 involved in the initial steps of the olfactory signal transduction cascade. Accordingly, the

NOV7 nucleic acid, polypeptide, antibodies and other compositions of the present invention can be used to detect nasal epithelial neuronal tissue.

Based on its relatedness to the known members of the OR family of the GPCR superfamily, NOV7 satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the treatment of disorders associated with alterations in the expression of members of OR family-like proteins. NOV7 Nucleic acids, polypeptides, antibodies, and other compositions of the present invention are useful in the treatment and/or diagnosis of a variety of diseases and pathologies, including by way of nonlimiting example, those involving neurogenesis, cancer and wound healing.

NOV8

A NOV8 sequence according to the invention is a nucleic acid sequence encoding a polypeptide related to the mast cell protease family of proteins. A NOV8 nucleic acid and its encoded polypeptide includes the sequences shown in Table 27. The disclosed nucleic acid (SEQ ID NO:15) is 948 nucleotides in length and contains an open reading frame (ORF) that begins with an ATG initiation codon at nucleotides 61-63 and ends with a TAG stop codon at nucleotides 931-934. The representative ORF encodes a 290 amino acid polypeptide (SEQ ID NO:16). PSORT analysis suggests that a NOV8 polypeptide is contained within the mitochondrial matrix, with a certainty of 0.4366. SIGNALP predicts a signal peptide with the most likely cleavage site between positions 16 and 17 of SEQ ID NO.: 16. Putative untranslated regions up- and downstream of the coding sequence are underlined in SEQ ID NO: 15.

TABLE 27.

TGACCTCCCTGCCTGATGGGCTCTGTGCCAGGAACCCAGGCGAGTCCGCCCCACCAATGCCCTGCTGCCA
GCCGGTCTCTCTGGTGGCCCTGAGCTCTGGGAAGACCTCGTCCGTCCCCCTCATGAGCCCGGCACGGGGCGTGA
GCTGGTGGGCATCACTGGGGGCTGCGACGTCTCGGCCAGGAGGCACCCCTGGCAGGTCAGCCTGAGGTTCTACAGC
ATGAAGAAGGGTCTGTGGGAGCCCATCTGTGGGGGCTCCCTCATCCACCCAGAGTGGGTGCTGACCGCCGCCACT
GCCTTGGGCTGAGGAGTTGGAGGCTTGCGCGTTTAGAGTGCAGGTGGGGCAGCTGAGGCTCTATGAGGACGACCA
GCGGACGAAGGTGGTTGAGATCGTCCGTACCCCCAGTACAACGAGAGCCTGTCTGCCAGGGCGGTGCGGACATC
GCCCTGCTGAAGCTGGAGGCCCGGTGCGGCTGTCTGAGCTCATCCACCGGTCTCGCTCCCGTCTGCCTCCCTGG
ACGTGCCCTCGGGGAAGACCTGCTGGGTGACCGGCTGGGGTGTCATTGGACGTGGAGAACTACTGCCCTGGCCCT
CAGCTTGTGGGAGGCGACGGTGAAGGTGAGGAGCAACGTCTCTGTAAACGACCTGTCGCCGCGCTTTCCTTCC
AACCACACTGAGCGGTTTGAGCGGCTCATCAAGGACGACATGCTGTGTGCCGGGGACGAGCGCCATCTCTCCCCAC
AGGGCGACAACGGGGGCCCCCTCCTGTGCAGGCGGAATTGCACCTGGGTCCAGGTGGAGGTGGTGAAGTGGGGCAA
ACTCTGCGGCCTTCGCGGTATCCCGGCATGTACACCGCGTGACGAGTACGTGTCTGGATCCGCCAGTACGTC
CCGCGTTCCCCAGACGCTAGTGGGGTGAGTGGG (SEQ ID NO.: 15)

5 MPLLPSRSLVPLSSGKTLVRPPHEPGTGRELVGITGGCDVSARRHPWQVSLRFYSMKKGLWEPICGGS
 LIHPEWVLTAACHLGPEELEACAFRVQVGQLRLYEDDQRTKVVEIVRHPQYNESLSAQGGADIALLLKLE
 APVPLSELVPLSASLDVPSGKTCWVTGWGVIGRGELLWPPLSLWEATVKVRSNVLCNQTCRRRFP
 SNHTEFERFERLIKDDMLCAGDERHLSPOGDNGGPLLCCRNTWVQVEVVSWSGKLCGLRGYPGMYTRVTSY
 VSWIRQYVPPFP (SEQ ID NO.: 16)

The NOV8 polypeptide has homology (58% identity, 66% similarity) to a canine
 mastocytoma protease precursor (MPP; SwissProt Accession No.: P19236), as is shown in
 Table 28. The NOV8 polypeptide also has homology (48% identity, 61% similarity) to a
 10 human beta tryptase precursor protein (BTPP; SwissProt Accession No.: Q13607), as is shown
 in Table 29.

Table 28.

NOV8:	27	GTGRELVGITGGCDVSARRHPWQVSLRFYSMKKGLWEPICGGS	LIHPEWVLTAACHLGPE	86
		** *** * * * * * * * * * * * * * * *		
15 MPP:	12	GTLSPKVGIVGGCKVPARRYPWQVSLRFHGMGSGQWQHICGGS	LIHPQWVLTAACHVBLE	71
NOV8:	85	ELEACAFRVQVGQLRLYEDDQRTKVVEIVRHPQYNESLSAQGGADIALLLKLEAPVPLSEL	144	
		*** *****+ ** * *+*** +* * *****+ ***		
20 MPP:	72	GLEAATLRVQVGQLRLYDHDQLCNVTEIIRHPNFMNSWYGWDTADIALLLKLEAPLTSED	131	
NOV8:	145	IHPVSLPSASLDVPSGKTCWVTGWGVIGRGELLWPPLSLWEATVKVRSNVLCNQTCRRRF	204	
		++ ***** * * * * * * * * * * * * * *		
MPP:	132	VNLVSLPSPSLIVPPGMLCWTGWGDIADHTPLPPPYHLQVEVEVPIVGNREC	--CHYQ- 188	
25 NOV8:	205	PSNHTEFERFERLIKDDMLCAGDERHLSPOGDNGGPLLCCRNTWVQVEVVSWSGKLCGLRGY	264	
		+ + + ***** * * * * *+***** ***+*** ***** *		
MPP:	189	--TILEQDDEVIKQDMLCAGSEGHDSQMDSGGPLVCRWKCTWIQVGVVSWGYGCGYN-L	245	
NOV8:	265	PGMYTRVTSYVSWIRQYVPPFP	286 (SEQ ID NO.: 71)	
30		***+ ***** *+* *		
MPP:	246	PGVYARVTSYVSWIHQHIPLSP	267 (SEQ ID NO.: 72)	

Where * indicates identity and + indicates similarity.

Table 29.

35 NOV8:	26	PGTGRELVGITGGCDVSARRHPWQVSLRFYSMKKGLWEPICGGS	LIHPEWVLTAACHLGP	85
		** + *** ** + + ***** + * *****+*****+***		
BTPP:	22	PGQALQRVGIVGGQEAPRSKWPQVSLRVHGP---	YWMHFCGGS	LIHPQWVLTAACHVGP 78
NOV8:	86	ELEACAFRVQVGQLRLYEDDQRTKVVEIVRHPQYNESLSAQGGADIALLLKLEAPVPLSE	145	
40		+ + * ***+ + ** ** * *+ ***+ +** *****+*** ** +*		
BTPP:	79	DVKDLAALRVQLREQHLYYQDQLLPVSRIIVHPQF---	YTAQIGADIALLELEEPVKVSS 135	
NOV8:	146	LIHPVSLPSASLDVPSGKTCWVTGWGVIGRGELLWPPLSLWEATVKVRSNVLCNQTCRRR	205	
		+* +*** ** * * ***** + * ** * * + * + * +* *		
45 BTPP:	136	HVHTVTLPASETFPMPGMCWVTGWGDVNDERLPPFPPLKQVKVPIMENHICDA---	KY 192	
NOV8:	206	FPSNHTEFERFERLIKDDMLCAGDERHLSPOGDNGGPLLCCRNTWVQVEVVSWSGKLCGLRG	265	
		+* *+*+*****+ * * *+*****+* * *+* *****+ *		
50 BTPP:	193	HLGAYTGDDVRIVRDDMLCAGNTRRDSQGDGGPLVCKVNGTTLQAGVVSWGEGCAQPN	252	
NOV8:	266	YPGMYTRVTSYVSWIRQYVPPFP	288 (SEQ ID NO.: 73)	

+** *+ ** *** *

BTPP: 253 RPGIYTRVTYYLDWIHHYVPKKP 275 (SEQ ID NO.: 74)

Where * indicates identity and + indicates similarity.

5

The term mastocytosis denotes a heterogenous group of disorders characterized by abnormal growth and accumulation of mast cells in one or more organs. Cutaneous and systemic variants of the disease have been described. Mast cell disorders have also been categorized according to other aspects, such as family history, age, course of disease, or

10 presence of a concomitant myeloid neoplasm. However, so far, generally accepted disease criteria are missing. Recently, a number of diagnostic (disease-related) markers have been identified in mastocytosis research. These include the mast cell enzyme tryptase, CD2, and mast cell growth factor receptor c-kit (CD117). The mast cell enzyme tryptase is increasingly used as a serum- and immunohistochemical marker to estimate the actual spread of disease

15 (burden of neoplastic mast cells). The clinical significance of novel mastocytosis markers is currently under investigation. First results indicate that they may be useful to define reliable criteria for the delineation of the disease.

The NOV8 nucleic acids and proteins of the invention are useful in potential therapeutic applications implicated in disorders characterized by abnormal growth and

20 accumulation of mast cells in one or more organs including, but not limited to skin, ear and brain as well as other pathologies and disorders. The NOV8 nucleic acid and protein of the invention, or fragments thereof, may further be useful in diagnostic applications, wherein the presence or amount of the NOV8 nucleic acid or the protein are to be assessed.

25 NOV9

A NOV9 sequence according to the invention is a nucleic acid sequence encoding a polypeptide related to the hepatocyte nuclear factor-like family of proteins. A NOV9 nucleic acid and its encoded polypeptide includes the sequences shown in Table 30. The disclosed nucleic acid (SEQ ID NO:17) is 542 nucleotides in length and contains an open reading frame

30 (ORF) that begins with an ATG initiation codon at nucleotides 7-9 and ends with a TGA stop codon at nucleotides 514-516. The representative ORF encodes a 169 amino acid polypeptide (SEQ ID NO:18). The predicted molecular weight of a NOV9 polypeptide is 19458.9 Da. PSORT analysis suggests that a NOV9 polypeptide is contained within the microbody

(peroxisome), with a certainty of 0.6400. Putative untranslated regions up- and downstream of the coding sequence are underlined in SEQ ID NO: 17.

TABLE 30.

5 TATGCCATGTATACGAATTCGAGCTCCTACCAGACTGGCCCGAATCATGAGTTCT
 ACAAGAACGCCGACGTCCGGCCCCCTTCACCTACGCCTCCCTCATCCGCCAGGC
 CATCCTGGAAACCCCTGACAGGCAGCTGACCCTGAATGAGATCTATAACTGGTTC
 ACCAGGATGTTTCGCCTATTTCCGCAGAAACACTGCCACCTGGAAGAACGCCGTGC
 10 GCCACAACCTCAGCCTGCACAAGTGCTTCGTCCGCGTGGAGAACGTCAAGGGTGC
 CGTGTGGACTGTGGACGAGCGGGAGTATCAGAAGCGGAGACCGCCAAAGATGAC
 AGGGTATGTGGGTCCAGAGCTGGATGGGCTGTACCTGCCCAGGGGGCAGGAGCCA
 ACTCACCCCCACCCCTACCTCTCCAGGGTACACATGTGCACCAGATCCTTCCTGG
 CTGGGGGAAGGGGTGTGGGGAGAAAGGAGCAGAGGAGACTAGTGCTTGGGGACA
 15 GGGGGCTGGAATCCGGAAGTGATGGATAATCAGAAGGCAGACATTAT (SEQ ID
 NO.: 17)

MYTNSSSYQTGPNHEFYKNADVRRPPFTYASLIRQAILETPDRQLTLNEIYNWFTRMFA
 YFRNTATWKNVVRHNLHLKCFVRVENVKGAVWTVDEREYQKRPPKMTGYVGP
 20 ELDGLYLPRGQEPHTPHPLPLQGTHVHQILPGWVGKGCGEKGAEETSAWGQGAGIRK
 (SEQ ID NO.: 18)

The NOV9 nucleic acid has a high degree of homology (100% identity) with a region of clone RP11-328M4 on chromosome 6 (CHR6; Genbank Accession No.: AL139331), as shown in Table 31. The NOV9 polypeptide has a high degree of homology (approximately 90% identity, 96% similarity) to a glutamine (Q)-rich factor-1 (QRF-1; EMBL Accession No.: G455862), as is shown in Table 32. Also, the NOV9 polypeptide has homology (66% identity, 82% similarity) with a mouse fork-head protein (mFHP; PatP Accession No.: Y77662), as is shown in Table 33.

TABLE 31.

30 NOV9: 209 agaacgccgtgcccacaacctcagcctgcacaagtgttcgtccgcgtggagaacgtca 268
 |||
 Chr6: 166408 agaacgccgtgcccacaacctcagcctgcacaagtgttcgtccgcgtggagaacgtca 166467

35 NOV9: 269 aggggtgccgtgtggactgtggacgagcgggagtagatcagaagcggagaccgccaagatga 328
 |||
 Chr6: 166468 aggggtgccgtgtggactgtggacgagcgggagtagatcagaagcggagaccgccaagatga 166527

40 NOV9: 329 cagggtatgtgggtccagagctggatgggctgtacctgcccagggggcaggagccaactc 388
 |||
 Chr6: 166528 cagggtatgtgggtccagagctggatgggctgtacctgcccagggggcaggagccaactc 166587

45 NOV9: 389 acccccacccctacctctccagggtacacatgtgcaccagatccttcctggctggggga 448
 |||
 Chr6: 166588 acccccacccctacctctccagggtacacatgtgcaccagatccttcctggctggggga 166647

NOV9: 449 aggggtgtggggagaaaggagcagaggagactagtgccttggggacagggggctggaatcc 508
 ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
 Chr6: 166648 aggggtgtggggagaaaggagcagaggagactagtgccttggggacagggggctggaatcc 166707
 NOV9: 509 ggaagtgatggataatcagaaggcagacatttat 542 (SEQ ID NO.: 75)
 ||||||||||||||||||||||||||||||||||||||||
 Chr6: 166708 ggaagtgatggataatcagaaggcagacatttat 166741 (SEQ ID NO.: 76)

TABLE 32.

NOV9: 23 RPPFTYASLIRQAILETPDRQLTLNEIYNWFTRMFAYFRRNTATWKNVHRNLSLHKCFV 82
 *****+*+*****
 QRF-1: 1 RPPFTYASLIRQAILESPEKQLTLNEIYNWFTRMFAYFRRNAATWKNVHRNLSLHKCFV 60
 NOV9: 83 RVENVKGA VWTVDEREYQKRRPPK 106 (SEQ ID NO.: 77)
 *****+*+*****
 QRF-1: 61 RVENVKGA VWTVDDEFQKRRPQK 84 (SEQ ID NO.: 78)
 Where * indicates identity and + indicates similarity.

TABLE 33.

NOV9: 1 YAMYTNSSSYQTGPNHEFYKNADV RPPFTYASLIRQAILETPDRQLTLNEIYNWFTRMFA 60
 + + *** + * +++* +*****+*** *****+*****
 MFHP: 313 WSGHGNSSFPEFFHNMDYFKYHNMRPFTYATLIRWAILEAPERQRTLNEIYHWFTRMFA 372
 NOV9: 61 YFRRNTATWKNVHRNLSLHKCFVRVENVKGA VWTVDEREYQKRR 125 (SEQ ID NO.: 79)
 *** + *****+*****+*****+*+*+*
 MFHP: 373 YFRNHPATWKNAIRHNLSLHKCFVRVESEKGA VWTVDEFEFRKKR 417 (SEQ ID NO.: 80)
 Where * indicates identity and + indicates similarity.

A NOV9 polypeptide is highly related to QRF-1, a B-cell-derived DNA-binding protein, and mFHP, which are members of the hepatocyte nuclear factor 3/fork-head family of proteins. A NOV9 nucleic acid is also useful as a marker for chromosome 6. Based on its relatedness to the known members of the hepatocyte nuclear factor 3/fork-head family, NOV9 satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the treatment of disorders associated with alterations in the expression of members of hepatocyte nuclear factor 3/fork head-like proteins. NOV9 nucleic acids, polypeptides, antibodies, and other compositions of the present invention are useful in the treatment and/or diagnosis of a variety of diseases and pathologies, including by way of nonlimiting example, those involving hepatic disorders, *e.g.* liver cancer, cirrhosis, ischaemia-reperfusion injury, and diabetes.

NOV10

A NOV10 sequence according to the invention is a nucleic acid sequence encoding a polypeptide related to the mast cell protease family of proteins. A NOV10 nucleic acid and its encoded polypeptide includes the sequences shown in Table 34. The disclosed nucleic acid (SEQ ID NO:19) is 870 nucleotides in length and contains an open reading frame (ORF) that begins with an ATG initiation codon at nucleotides 43-45 and ends with a TAA stop codon at nucleotides 868-870. The representative ORF encodes a 275 amino acid polypeptide (SEQ ID NO:20). The predicted molecular weight of a NOV10 polypeptide is 30,467.7 Da. PSORT analysis suggests that a NOV10 polypeptide is contained within the lysosome, with a certainty of 0.8650. A putative untranslated region upstream of the coding sequence is underlined in SEQ ID NO: 19. SIGNALP analysis indicates a probable signal peptide with the most likely cleavage site occurring between positions 19 and 20.

TABLE 34.

15 ATCTGGCCAGAGTGGGCTTGGCCAGTTGTGGTGGGCACCACCATGCTGCTGCTGCTGCTGTTCTGGCT
 G
 TCTCCTCCCTGGGGAGCTGTAGCACTGGGAGTCCAGCTCCCGTCCCCGAGAATGACCTGGTGGGCATTG
 T
 GGGGGGCCACAACACCCAGGGGAAGTGGTCGTGGCAGGTGAGCCTGAGGATCTATAGCTACCACTGGGC
 20 C
 TCCTGGGTGCCCATCTGCGGGGGCTCCCTCATCCACCCCCAGTGGGTGCTGACCGCCGCTCACTGCATT
 T
 TCCGGAAGGACACCGACCCGTCACCTACCGGATTACACCCAGGGATGTGTATCTGTACGGGGGCCGGG
 G
 25 GCTGCTGAATGTCAGCCAGATCGTCGTCCACCCCACTACTCTGTCTTCTTCTGGGGGCAGACATCGC
 C
 CTGCTGAAGCTGGCCACCAGTGTGAGAACAACAACACTCTCGCGGCAGTCGCCCTGCCGTCATTGTCC
 C
 TGGAGTTCACTGACAGTGACAACCTGCTGGAACACAGGCTGGGGCATGGTCGGCTTGTTGGATATGCTGC
 30 C
 GCCTCCTTACCGCCCGCAGCAGGTGAAGTCTCACACTGAGCAATGCAGACTGTGAGCGGCAGACCTA
 C
 GATGCTTTTCTGGTGCTGGAGACAGAAAGTTCATCCAGGATGACATGATCTGTGCCGGCCGCACGGGC
 C
 35 GCCGCACCTGGAAGGGTGACTCAGGCGGCCCCCTGGTCTGCAAGAAGAAGGGTACCTGGCTCCAGGCGG
 G
 AGTAGTGAGCTGGGGATTTTACAGTGATCGGCCCAGCATTGGCGTCTACACGTGGGTCCAGACCTATGT
 G
 CCCTGGATCCTGCAGCAAATGCACCTCTAA (SEQ ID NO.: 19)
 40 MLLLLLFLAVSSLGSCSTGSPAPVPENDLVGIVGGHNTQGKWSWQVSLRIYSYHWASWVPICGGLIHP
 Q
 WVLTAHCIFRKDTDPSTYRIHTRDVLYGGRGLLNVSQIVVHPNYSVFFLGADIALLLKLATSVRTTNT
 L

AAVALPSLSLEFTSDNCWNTGWMVGLLDMLPPPYRPQQVKVLTLSNADCERQTYDAFFGAGDRKFIQ
D
DMICAGRTGRRTWKGDSSGGPLVCKKKGTWLQAGVVSWSGFYSDRPSIGVYTWVQTYVPWILQQMHL (SEQ
ID NO.: 20)

5

A NOV10 nucleic acid has a high degree of homology (92% identity) with an uncharacterized region of human chromosome 16 including clone LA16-303A1 (CHR 16; Genbank Accession No.: HS303A1), as is shown in Table 35. A NOV10 polypeptide has homology (58% identity, 66% similarity) to a human mast cell tryptase II/beta (MCTII; PatP Accession No.: W64240), as is shown in Table 36. A NOV10 polypeptide also has homology (48% identity, 63% similarity) to a mouse mast cell protease 6 precursor protein (MCP6; SwissProt Accession No.: P21845), as is shown in Table 37.

TABLE 35.

15	NOV10: 109	cccgtccccgagaatgacctggtgggcattgtggggggccacaa---caccaggggaag	165
	CHR16: 22021	cccgtcccagagaatgacctggtgggcattgtggggggccacaatgccccccggggaag	21962
	NOV10: 166	tggtcggtggcaggtcagcctgaggatctatagctaccactgggcctcctgggtgcccatc	225
20	CHR16: 21961	tggtcggtggcaggtcagcctgaggatctatagctaccactgggcctcctgggtgcccatc	21902
	NOV10: 226	tgccgggggtccctcatccacccccagtggtgctgaccgccgctcactgcattttc	282 (SEQ ID NO.: 81)
25	CHR16: 21901	tggtgggggtccctcatccacccccagtggtgctgactgctgccactgcattttc	21845 (SEQ ID NO.: 82)

30

TABLE 36.

	NOV10: 2	LLLLFLAVSSLGSCSTGSPAPVPENDLVGIVGGHNT-QGKWSWQVSLRIYSYHWASWVPI	60
	MCTII: 1	LNLLLLALPVLASRAYAAPAGQALQRVGIVGGQEA PRSKWPQVSLRV---HGPLYWMHF	57
35	NOV10: 61	CGGSLIHPQWVLTAAHCIFRKDTPSTYRIHTRDVLYGGRGLLNVSQIVVHPNYSVFFL	120
	MCTII: 58	CGGSLIHPQWVLTAAHCVGPDVKDLAALRVQLREQHLYYQDQLLPVSR IIVHPQFYTAQI	117
40	NOV10: 121	GADIALKLKATSVRTTNTLAAVALPSLSLEFTSDNCWNTGWMVGLLDMLPPPYRPQOV	180
	MCTII: 118	GADIALLELEEPVKVSSHVHTVTLPPASETFPFGMPCWVTGWGDVDNDRLPFFFLKQV	177
45	NOV10: 181	KVLTLSNADCERQTY-DAFFGAGDRKFIQDDMICAGRTGRRTWKGDSSGGPLVCKKKGTWL	240
50	MCTII: 178	KVPIMENHICDAKYHLGAYTG-DDVRIVRDDMLCAGNTRRDSQCQDSSGGPLVCKVNGTWL	236

NOV10: 241 QAGVVSWSGFYSRPSI-GVYTWVQTYVPWI 269 (SEQ ID NO.: 83)
 ***** ++ *+* * +* **
 MCTII: 237 QAGVVSWSGEGCAQPNRPGIYTRVYYLDWI 266 (SEQ ID NO.: 84)
 Where * indicates identity and + indicates similarity.

5

Table 37.

NOV10: 22 APVPENDLVGIVGGHN-TQKWSWQVSLRIYSYHWASWVPICGGS LIHPQWVLTAAHCIF 80
 ** * * ***** ++ ** ***** +* + *****
 MCP6: 22 APRPANQVRVGIVGGHEASESKWPQVSLRFKLNW---IHFCGGS LIHPQWVLTAAHCVG 78
 NOV10: 81 RKDTPSTYRIHTRDVYLYGGRGLNVSQIVVHPNYSVFFLGADIAL LKLTASVRTNTL 140
 * ++ +* *** * **++++*****+* ****+***+* * + +
 MCP6: 79 PHIKSPQLFRVQLREQYLYYGDQLLSLRIVVHPHYTAEAGGADVALLEVPVNVSTHI 138
 NOV10: 141 AAVALPSLSLEFTSDNCWNTGWMVGLLDMLPPPYRPQQVKVLTLSNADCERQTYDAFP 200
 +++* * * +** ***** + +***** +***** +*+ *+++ +
 MCP6: 139 HPISLPPASETFPPGTSCWVTGWGDIDNDEPLPPPYPLKQVKVPIVENS LDRKYHTGLY 198
 NOV10: 201 GAGDRKFIQDDMICAGRTGRRTWKGDSGGPLVCKKKGTWLQAGVVSWSGFYSRPS-IGVY 259
 * + * *+*** * * + +***** ***** +*+ **
 MCP6: 199 TGDDFPIVHDGMLCAGNTRRDSQCQDSGGPLVCKVKGTLQAGVVSWSGEGCAQPNKPGIY 258
 NOV10: 260 TWVQTYVPWI 269 (SEQ ID NO.: 85)
 * * *+ **
 MCP6: 259 TRVYYLDWI 268 (SEQ ID NO.: 86)
 Where * indicates identity and + indicates similarity.

25

30

35

40

The term mastocytosis denotes a heterogeneous group of disorders characterized by abnormal growth and accumulation of mast cells in one or more organs. Cutaneous and systemic variants of the disease have been described. Mast cell disorders have also been categorized according to other aspects, such as family history, age, course of disease, or presence of a concomitant myeloid neoplasm. However, so far, generally accepted disease criteria are missing. Recently, a number of diagnostic (disease-related) markers have been identified in mastocytosis research. These include the mast cell enzyme tryptase, CD2, and mast cell growth factor receptor c-kit (CD117). The mast cell enzyme tryptase is increasingly used as a serum- and immunohistochemical marker to estimate the actual spread of disease (burden of neoplastic mast cells). The clinical significance of novel mastocytosis markers is currently under investigation. First results indicate that they may be useful to define reliable criteria for the delineation of the disease.

The NOV10 nucleic acids and proteins of the invention are useful in potential therapeutic applications implicated in disorders characterized by abnormal growth and accumulation of mast cells in one or more organs including, but not limited to skin, ear and brain as well as other pathologies and disorders. The NOV10 nucleic acid and protein of the

invention, or fragments thereof, may further be useful in diagnostic applications, wherein the presence or amount of the NOV10 nucleic acid or the protein are to be assessed.

NOV11

5 A NOV11 sequence according to the invention is a nucleic acid sequence encoding a polypeptide related to the mast cell protease family of proteins. A NOV11 nucleic acid and its encoded polypeptide includes the sequences shown in Table 38. A NOV11 nucleic acid is localized to human chromosome 16. The disclosed nucleic acid (SEQ ID NO:21) is 858
10 nucleotides in length and contains an open reading frame (ORF) that begins with an ATG initiation codon at nucleotides 1-3 and ends with a TAG stop codon at nucleotides 856-858. The representative ORF encodes a 285 amino acid polypeptide (SEQ ID NO:22). PSORT analysis suggests that a NOV11 polypeptide is either a luminal lysosomal protein (certainty of 0.4766) or a secreted protein (certainty 0.3700). SIGNALP analysis indicates a probable signal peptide with the most likely cleavage site occurring between positions 14 and 15.

15

TABLE 38.

ATGCTGTGGCTACTGCTCCTGACCCTCCCCTGCCTGATGGGCTCTGTGCCCAGGAA
CCCAGGCGAGGGCACGGGGCGTGAGCTGGTGGGCATCACTGGGGGCTGCGACGT
CTCGGCCAGGAGGCACCCCTGGCAGGTCAGCCTGAGGTTCTACAGCATGAAGAAG
20 GGTCTGTGGGAGCCCATCTGTGGGGGCTCCCTCATCCACCCAGAGTGGGTGCTGA
CCGCCGCCCACTGCCTTTTGGAGGAGTTGGAGGCTTGCGCGTTTAGAGTGCAGGT
GGGGCAGCTGAGGCTCTATGAGGACGACCAGCGGACGAAGGTGGTTGAGATCGT
CCGTACACCCCACTACAACGAGAGCCTGTCTGCCCAGGGCGGTGCGGACATCGCC
CTGCTGAAGCTGGAGGCCCCGGTGCCGCTGTCTGAGCTCATCCACCCGGTCTCGCT
25 CCCGTCTGCCTCCCTGGACGTGCCCTCGGGGAAGACCTGCTGGGTGACCGGCTGG
GGTGTCAATTGGACGTGGAGAACTACTGCCCTGGCCCCTCAGCTTGTGGGAGGCGA
CGGTGAAGGTCAGGAGCAACGTCCTCTGTAACCAGACCTGTCGCCGCCGCTTTCC
TTCCAACCACACTGAGCGGTTTGAGCGGCTCATCAAGGACGACATGCTGTGTGCC
GGGGACGGGAACACGGCTCCTGGCCAGGCGACAACGGGGGGCCCCCTCCTGTGC
30 AGGCGGAATTGCACCTGGGTCCAGGTGGAGGTGGTGAGCTGGGGCAAACCTCTGCG
GCCTTCGCGGCTATCCCGGCATGTACACCCGCGTGACGAGCTACGTGTCCTGGATC
CGCCAGTACGTCCCGCCGTTCCCCAGACGCTAG (SEQ ID NO.: 21)

35 MLWLLLLTLPCLMGSVPRNPGE GTGRELVGITGGCDVSARRHPWQVSLRFYSMKKG
LWEPICGGLIHPWVLTAAHCLLEELEACAFRVQVGQLRLYEDDQRTKVVEIVRHPQ
YNESLSAQGGADIALLKLEAPVPLSELHPVSLPSASLDVPSGKTCWVTGWGVIGRGE
LLPWPLSLWEATVKVRSNVLCNQTCRRRFPSNHTERFERLIKDDMLCAGDGNHGSWP
GDNGGPLLCRRNCTWVQVEVVSWGKLCGLRGYPGMYTRVTSYVSWIRQYVPPFPRR
40 (SEQ ID NO.: 22)

A NOV11 nucleic acid has a high degree of homology (92% identity) with an uncharacterized region of human chromosome 16 including clone LA16-303A1 (CHR 16; Genbank Accession No.: HS303A1), as is shown in Table 39. A NOV11 polypeptide has
 5 homology (58% identity, 66% similarity) to a canine mastocytoma protease precursor (cMPP; SwissProt Accession No.: P19236), as is shown in Table 40. A NOV11 polypeptide also has homology (46% identity, 60% similarity) to a human beta tryptase precursor (BTRP; SwissProt Accession No.: P20231), as is shown in Table 41.

TABLE 39.

10	NOV11: 180	catctgtgggggctccctcatccacccagagtgggtgctgaccgccgcccactgcctttt	239
	CHR16: 21905	catctgtgggggctccctcatccacccagagtgggtgctgactgctgcccactgcatttt	21846
15	NOV11: 182	tctgtgggggctccctcatccacccagagtgggtgctgaccgccgcccactgc	234 (SEQ ID NO. 87)
	CHR16: 4196	tctgcgggggctccctcatccacccagagtgggtgctgaccgcagcgccactgc	4144 (SEQ ID NO. 88)
20			

TABLE 40.

	NOV11: 23	GTGRELVGITGGCDVSARRHPWQVSLRFYSMKKGLWEPICGGSLIHPEWVLTAACHCL-LE	82
		** *** * **++*****+ * * +* *****+*****+ **	
25	cMPP: 1 2	GTLSPKVGIVGGCKVPARRYPWQVSLRFHGMGSGQWQHICGGSLIHQPWVLTAACHCELE	71
	NOV11: 83	ELEACAFRVQVQQLRLYEDDQRTKVVEIVRHPQYNESLSAQGGADIALLKLEAPVPLSEL	142
		*** *****+ ** * **++** +* * *****+ ***	
30	cMPP: 72	GLEAATLRVQVQQLRLYDHDQLCNVTEIIRHPNFMNSWYGWDTADIALLKLEAPLTSED	131
	NOV11: 143	IHPVSLPSASLDVPSGKTCWVTGWGVIGRGELLPWPLSLWEATVKVRSNVLNQTCTRRRF	202
35		++ ***** * * * ***** * * * * * + * * * +	
	cMPP: 132	VNLVSLPSPSLIVPPGMLCWVTGWGDIADHTPLPPPYHLQEVEVPVIGNREC--CHYQ-	188
	NOV11: 203	PSNHTERFERLIKDDMLCAGDGNHGSWPGDNGGPLLCCRNCCTWVQVEVVSWGKLCGLRGY	262
40		*+ + +** ***** * * *+*****+** ***** ***** **	
	cMPP: 189	--TILEQDDEVIKQDMLCAGSEGHDSQMDSGGPLVCRWKCTWVQVGVVSWGYGCGYN-L	245
45	NOV11: 263	PGMYTRVTSYVSWIRQYVPPFP	284 (SEQ ID NO. 89)
		+ ** **++ * *	
	cMPP: 246	PGVYARVTSYVSWIHQHIPLSP	267 (SEQ ID NO. 90)
	Where * indicates identity and + indicates similarity.		

Table 41.

5	NOV11: 20	PGE ₂ GTGRELVGITGGCDVSARRHPQVSLRFYSMKKGLWEPICGGSLIHPEWVLTAAHCL	79
	BTRP: 20	PAPGQALQ ₂ RVGIVGGQEAPRSKW ₂ PQVSLRVHGP---YWMHFCGGSLIH ₂ PQVWLTAAHCV	76
10	NOV11: 80	LEEL-EACAFRVQVGQLRLYEDDQRTKVVEIVRHPQYNESLSAQGGADIALLKLEAPVPL	138
	BTRP: 77	GPDVKDLAALRVQLREQHLYYQDQLLPVSRIIVHPQF---YTAQIGADIALLELEEPVKV	133
15	NOV11: 139	SELIHPVSLPSASLDVPSGKTCWVTGWGVIGRGELLPWPLSLWEATVKVRSNVLCNQTCTCR	198
	BTRP: 134	SSHVHTVTLPPASETFPPGMP ₂ CWVTGWGDVDN ₂ DERLPPFFPLKQVKVPIMENHICDA---	190
20	NOV11: 199	RRFPSNHTE ₂ FERFERLIKDDMLCAGDGNHGSWPGDNGGPLL ₂ CRRNCTWVQVEVSWGKLCGL	258
	BTRP: 191	KYHLGAYTGDDVRIVR ₂ DDMLCAGNTRR ₂ DS ₂ CQGD ₂ SGG ₂ PLVCKVNGTWLQAGV ₂ SWGEGCAQ	250
25	NOV11: 259	RGYPGMYTRVTSYVSWIRQYVPPFP	283 (SEQ ID NO. 91)
	BTRP: 251	PNRPGIYTRVTYYLDWIHHYVPKKP	275 (SEQ ID NO. 92)

Where * indicates identity and + indicates similarity.

The term mastocytosis denotes a heterogeneous group of disorders characterized by abnormal growth and accumulation of mast cells in one or more organs. Cutaneous and systemic variants of the disease have been described. Mast cell disorders have also been categorized according to other aspects, such as family history, age, course of disease, or presence of a concomitant myeloid neoplasm. However, so far, generally accepted disease criteria are missing. Recently, a number of diagnostic (disease-related) markers have been identified in mastocytosis research. These include the mast cell enzyme tryptase, CD2, and mast cell growth factor receptor c-kit (CD117). The mast cell enzyme tryptase is increasingly used as a serum- and immunohistochemical marker to estimate the actual spread of disease (burden of neoplastic mast cells). The clinical significance of novel mastocytosis markers is currently under investigation. First results indicate that they may be useful to define reliable criteria for the delineation of the disease.

The NOV11 nucleic acids and proteins of the invention are useful in potential therapeutic applications implicated in disorders characterized by abnormal growth and accumulation of mast cells in one or more organs including, but not limited to skin, ear and brain as well as other pathologies and disorder such as hemophilia, idiopathic thrombocytopenic purpura, autoimmune disease, allergies, immunodeficiencies, transplantation, graft versus host, anemia, ataxia-telangiectasia, lymphedema, tonsillitis, hypercoagulation, and sudden infant death syndrome.

The NOV11 nucleic acid and protein of the invention, or fragments thereof, may further be useful in diagnostic applications, wherein the presence or amount of the NOV11 nucleic acid or the protein are to be assessed.

5 NOV12

A NOV12 sequence according to the invention is a nucleic acid sequence encoding a polypeptide related to the mast cell protease family of proteins. A NOV12 nucleic acid and its encoded polypeptide includes the sequences shown in Table 42. The disclosed nucleic acid (SEQ ID NO:23) is 660 nucleotides in length and encodes a 220 amino acid polypeptide (SEQ
10 ID NO:24).

TABLE 42.

TCACTGGGGGCTGCGACGTCTCGGCCAGGAGGCACCCCTGGCAGGGAGGAGTTGG
AGGCTTGCGCGTTTAGAGTGCAGGTGGGGCAGCTGAGGCTCTATGAGGACGACCA
15 GCGGACGAAGGTGGTTGAGATCGTCCGTCACCCCCAGTACAACGAGAGCCTGTCT
GCCCAGGGCGGTGCGGACATCGCCCTGCTGAAGCTGGAGGCCCGGTGCCGCTGT
CTGAGCTCATCCACCCGGTCTCGCTCCCGTCTGCCTCCCGGGACGTGCCCTCGGGG
AAGACCTGCTGGGTGACCGGCTGGGGTGTCAATTGGACGTGGAGAAGTACTGCCCT
GGCCCCCTCAGCTTGTGGGAGGCGACGGTGAAGGTCAGGAGCAACGTCCTCTGTAA
20 CCAGACCTGTCGCCGCCGCTTTCCTTCCAACCACTGAGCGGTTTGAGCGGCTCA
TCAAGGACGACATGCTGTGTGCCGGGGACGGGAACCACGGCTCCTGGCCAGGCGA
CAACGGGGGGCCCCCTCCTGTGCAGGCGGAATTGCACCTGGGTCCAGGTGGAGGTG
GTGAGCTGGGGCAAACCTCTGCGGCCTTCGCGGCTATCCCGGCATGTACACCCGCG
TGACGAGCTACGTGTCTGGATCCGCCAGTACGTCCCGCCGTTCCCCAGACGC
25 (SEQ ID NO.: 23)

SLGAATSRPGGTPGREELEACAFRVQVGQLRLYEDDQRTKVVEIVRHPQYNESLSAQ
GGADIALLKLEAPVPLSELIHPVSLPSASRDVPSGKTCWVTGWGVIGRGELLPWPLSL
WEATVKVRSNVLCNQTCRRRFPSNHTEFERLIKDDMLCAGDGNHGSWPGDNNGGPL
30 LCRRNCTWVQVEVSWGKLCGLRGYPGMYTRVTSYVSWIRQYVPPFPRR (SEQ ID
NO.: 24)

A NOV12 nucleic acid has homology (82% identity) with a canine mast cell tryptase precursor (cMCT; Genbank Accession No.: M24665), as is shown in Table 43. A NOV12
35 polypeptide has homology (54% identity, 63% similarity) to a canine mastocytoma protease precursor (cMPP; SwissProt Accession No.: P19236), as is shown in Table 44. A NOV12 polypeptide also has homology (45% identity, 59% similarity) to a human beta tryptase precursor (BTRP; SwissProt Accession No.: P20231), as is shown in Table 45.

TABLE 43.

NOV12: 178 gcgacatcgccctgctgaagctggaggccccgggtgccgctgtctgagctcatccaccg 237
 |||||
 5 cMCT: 291 gcgacatcgccctgctgaagctggaggccccctgacgctctccgaggacgtcaacctg 350
 NOV12: 238 gtctcgctccccgtctgcctccccgggacgtgccctcggggaagacctgctgggtgaccggc 297
 || || |||||
 cMCT: 351 gtgtccctccccgtctccctccctgattgtccccccgggatgctatgctgggtgaccggc 410
 10 NOV12: 298 tgggg 302 (SEQ ID NO. 93)
 |||||
 cMCT: 411 tgggg 415 (SEQ ID NO. 94)

TABLE 44.

15 NOV12: 16 EELEACAFRVQVQGLRLYEDDQRTKVVEIVRHPQYNESLSAQGGADIALLKLEAPVPLSE 75
 * * * * *
 cMPP: 71 EGLEAATLRVQVQGLRLYDHDQLCNVTEIRHPNFMNSWYGWDTADIALLKLEAPLTSE 130
 NOV12: 76 LIHPVSLPSASRDVPSGKTCWVTGWGVIGRGELLPWPLSLWEATVKVRSNVLCNQTCTRRR 135
 ++ * * * * *
 20 cMPP: 131 DVNLVSLPSPSLIVPPGMLCWVTGWGDIADHTPLPPPYHLQVEVEVPIVGNRECNC--CHYQ 188
 NOV12: 136 FPSNHTERFERLIKDDMLCAGDGNHGSWPGDNGGPLLCRRNCTWVQVEVSWGKLCGLRG 195
 * + + * * * * *
 25 cMPP: 189 ---TILEQDDEVIKQDMLCAGSEGHDSQMDSGGPLVCRWKCTWIVGVVSWGYGCGY-N 244
 NOV12: 196 YPGMYTRVTSYVSWIRQYVPPFP 218 (SEQ ID NO. 95)
 * + * * * * *
 cMPP: 245 LPGVYARVTSYVSWIHQHIPLSP 267 (SEQ ID NO. 96)
 30 Where * indicates identity and + indicates similarity.

TABLE 45.

NOV12: 14 GREELEACAFRVQVQGLRLYEDDQRTKVVEIVRHPQYNESLSAQGGADIALLKLEAPVPL 73
 * + + * * * * *
 35 BTRP: 77 GPDVKDLAALRVQLREQHLYYQDQLLPVSRIIVHPQF---YTAQIGADIALLELEEPVKV 133
 NOV12: 74 SELIHPVSLPSASRDVPSGKTCWVTGWGVIGRGELLPWPLSLWEATVKVRSNVLCNQTCTR 133
 * + * * * * *
 40 BTRP: 134 SSHVHTVTLPASETFFPGMPCWVTGWGDVNDERLPPPFPLKQVKVPIMENHICDA--- 190
 NOV12: 134 RRFPSNHTERFERLIKDDMLCAGDGNHGSWPGDNGGPLLCRRNCTWVQVEVSWGKLCGL 193
 + * * * * *
 BTRP: 191 KYHLGAYTGDDVRIVRDDMLCAGNTRDSCQGDGGPLVCKVNGTWLQAGVVSWECECAQ 250
 45 NOV12: 194 RGYPGMYTRVTSYVSWIRQYVPPFP 218 (SEQ ID NO. 97)
 * + * * * * *
 BTRP: 251 PNRPGIYTRVTYYLDWIHHYVPPKP 275 (SEQ ID NO. 98)
 Where * indicates identity and + indicates similarity.

50 The term mastocytosis denotes a heterogenous group of disorders characterized by abnormal growth and accumulation of mast cells in one or more organs. Cutaneous and systemic variants of the disease have been described. Mast cell disorders have also been categorized according to other aspects, such as family history, age, course of disease, or

presence of a concomitant myeloid neoplasm. However, so far, generally accepted disease criteria are missing. Recently, a number of diagnostic (disease-related) markers have been identified in mastocytosis research. These include the mast cell enzyme tryptase, CD2, and mast cell growth factor receptor c-kit (CD117). The mast cell enzyme tryptase is increasingly
5 used as a serum- and immunohistochemical marker to estimate the actual spread of disease (burden of neoplastic mast cells). The clinical significance of novel mastocytosis markers is currently under investigation. First results indicate that they may be useful to define reliable criteria for the delineation of the disease.

The NOV12 nucleic acids and proteins of the invention are useful in potential therapeutic
10 applications implicated in disorders characterized by abnormal growth and accumulation of mast cells in one or more organs including, but not limited to skin, ear and brain as well as other pathologies and disorder such as hemophilia, idiopathic thrombocytopenic purpura, autoimmune disease, allergies, immunodeficiencies, transplantation, graft versus host, anemia, ataxia-telangiectasia, lymphedema, tonsilitis, hypercoagulation, and sudden infant death
15 syndrome.

The NOV12 nucleic acid and protein of the invention, or fragments thereof, may further be useful in diagnostic applications, wherein the presence or amount of the NOV12 nucleic acid or the protein are to be assessed.

20 NOV13

A NOV13 sequence according to the invention is a nucleic acid sequence encoding a polypeptide related to the mast cell protease family of proteins. A NOV13 nucleic acid and its encoded polypeptide includes the sequences shown in Table 46. A NOV13 nucleic acid is localized to human chromosome 16. The disclosed nucleic acid (SEQ ID NO:25) is 843
25 nucleotides in length and contains an open reading frame (ORF) that begins with an ATG initiation codon at nucleotides 11-13 and ends with a TAG stop codon at nucleotides 835-837. The representative ORF encodes a 275 amino acid polypeptide (SEQ ID NO:26). PSORT analysis suggests that a NOV13 polypeptide is a cytoplasmic protein (certainty of 0.45). SIGNALP analysis did not identify a signal peptide. Putative untranslated regions up- and
30 down-stream of the ORF are underlined in SEQ ID NO.: 25.

TABLE 46.

TGAGAGATAAAATGGGCTCCCAGAGATGCCAGGGAGGAGGCCCGGCACGGGGC
TGAGCTGGTGGGCATCACTGGGGGCTGCGACGTCTCGGCCAGGAGGCACCCCTGG
CAGGTCAGCCTGAGGTTCTACAGCATGAAGAAGGGTCTGTGGGAGCCCATCTGTG
5 GGGGCTCCCTCATCACCCAGAGTGGGTGCTGACCGCCGCCACTGCCTTGGCAG
GGAGGAGTTGGAGGCTTGCGCGTTTTAGAGTGCAGGTGGGGCAGCTGAGGCTCTAT
GAGGACGACCAGCGGACGAAGGTGGTTGAGATCGTCCGTCACCCCCAGTACAACG
AGAGCCTGTCTGCCCAGGGCGGTGCGGACATCGCCCTGCTGAAGCTGGAGGCCCG
GGTGCCGCTGTCTGAGCTCATCACCCGGTCTCGCTCCCGTCTGCCTCCCGGCCTG
10 GGCTCCAGACGCGTCTTGATGGCTTCCTGCCGCTGCCGAGACGGATGGGCAGGA
ACTACTGCCCTGGCCCCCTCAGCTTGTGGGAGGCGACGGTGAAGGTCAGGAGCAAC
GTCCTCTGTAACCAGACCTGTCGCCGCCGCTTTCCCTCCAACCACTGAGCGGTT
TGAGCGGCTCATCAAGGACGACATGCTGTGTGCCGGGGACGGGAACCACGGCTCC
TGGCCAGGCGACAACGGGGGGCCCCCTCCTGTGCAGGCGGAATTGCACCTGGGTCC
15 AGGTGGAGGTGGTGAGCTGGGGCAAACCTCTGCGGCCTTCGCGGCTATCCCGGCAT
GTACACCCGCGTGACGAGCTACGTGTCCTGGATCCGCCAGTACGTCCCGCCGTTCC
CCAGACGCTAGCTGGG (SEQ ID NO.: 25)

20 MGSQRCQGGGPGTGRELVGITGGCDVSARRHPWQVSLRFYSMKKGLWEPICGGSLIH
PEWVLTAACHCLGREELEACA FRVQVGQLRLYEDDQRTKVVEIVRHPQYNESLSAQG
GADIALLKLEAPVPLSELHPVSLPSASRPG LQTRPGWLPAA AETDQGELLPWPLSLWE
ATVKVRSNVLCNQT CRRRFPSNHTE RFERLIKDDMLCAGDGNHGSWPGDNGG PLLC
RRNCTWVQVEVVS WGKLCGLRGYPGMYTRVTSYVSWIRQYVPPFPRR (SEQ ID NO.:
26)

A NOV13 nucleic acid has homology (84% identity) with a canine mast cell tryptase precursor (cMCT; Genbank Accession No.: M24665), as is shown in Table 47. A NOV13 polypeptide has homology (54% identity, 63% similarity) to a canine mastocytoma protease precursor (cMPP; SwissProt Accession No.:P19236), as is shown in Table 48. A NOV13 polypeptide also has homology (43% identity, 57% similarity) to a human beta tryptase precursor (BTRP; SwissProt Accession No.: P20231), as is shown in Table 49.

TABLE 47.

[illegible]

TABLE 48.

5	NOV13:	12	GTGRELVGITGGCDVSARRHPWQVSLRFYSMKKGLWEPICGGSLIHPEWVLTAAHCLGRE	71
			** *** ** * **++*****+ * * *+ *****+*****+ *	
10	cMPP:	12	GTLSPKVGIVGGCKVPARRYPWQVSLRFHGMGSGQWQHICGGSLIHPEWVLTAAHCVLE	71
15	NOV13:	72	ELEACAFRVQVGQLRLYEDDQRTKVVEIVRHPQYNESLSAQGGADIALLLKLEAPVPLSEL	131
			*** *****+ ** * **++*+ * * *****+ ***	
20	cMPP:	72	GLEAATLRVQVGQLRLYDHDQLCNVTEIIRHPNFMNSWYGWDTADIALLLKLEAPLTSED	131
25	NOV13:	132	IHPVSLPSASRPGLQTRPG---WLPAAAE TDGQELLPWFLSLWEATVKVRSNVLNQTCT	188
			++ ***** * * ** *+ + * * * * * + * ** *	
30	cMPP:	132	VNLVSLPS---PSLIVPPGMLCWVTGWGDIADHTPLPPYHLQEVEVPVGNREC---CH	186
35	NOV13:	189	RRFPSNHTERFERLIKDDMLCAGDGNHGSWPGDNGGPLLCRRNCTWVQVEVSWGKLCGL	248
			+ *+ + +** ***** * * *+*****+* ***+*+ ***** **	
40	cMPP:	187	YQ---TILEQDDEVIKQDMLCAGSEGHDSQMDSGGPLVCRWKCTWVQVGVVSWGYGCGY	243
45	NOV13:	249	RGYPGMYTRVTSYVSWIRQYVPPFP	273 (SEQ ID NO. 101)
			+ ** **++* *	
50	cMPP:	244	N-LPGVYARVTSYVSWIHQHIPLSP	267 (SEQ ID NO. 102)
			Where * indicates identity and + indicates similarity.	

TABLE 49.

35	NOV13:	1	MGSQRCQGGPGTGRELVGITGGCDVSARRHPWQVSLRFYSMKKGLWEPICGGSLIHPEW	60
			+ ** *+ *** ** + + ***** + * *****+	
40	BTRP:	12	LASRAYAAPAPGQALQRVGIVGGQEAPRSKWPWQVSLRVHGP---YWMHFCGGSLIHPEW	68
45	NOV13:	61	VLTAACHCLGREELEACAFRVQVGQLRLYEDDQRTKVVEIVRHPQYNESLSAQGGADIAL	120
			*****+* + + * **++ + ** * * **++*+* +** *****	
50	BTRP:	69	VLTAACHCVGPDVKDLAALRVQLREQHLYYQDQLLPVSRIIVHPQF---YTAQIGADIAL	125
55	NOV13:	121	KLEAPVPLSELIHPVSLPSASRPGLQTRPGWLPAAAE TDGQELLPWFLSLWEATVKVRSN	180
			+++ ** +* **++** * * **++ + * * * * * + * + *	
60	BTRP:	126	ELEPVKVSSHVHTVTLPPASETFPPGMPWCWVTGWGDVNDERLPPFPKQVKVPIMEN	185
65	NOV13:	181	VLNQTCTRRFPSNHTERFERLIKDDMLCAGDGNHGSWPGDNGGPLLCRRNCTWVQVEV	240
			+++ + ** *****+ * **++*****+* **++ **	
70	BTRP:	186	HICDA---KYHLGAYTGDDVRIVRDMLCAGNTRRDSQGDGGGPLVCKVNGTWLQAGVV	242
75	NOV13:	241	SWGKLCGLRGYPGMYTRVTSYVSWIRQYVPPFP	273 (SEQ ID NO. 103)
			+ * **++**+* ** * * *	
80	BTRP:	243	SWGEGCAQPNRPGIYTRVTTYLDWIHHYVPPKP	275 (SEQ ID NO. 104)
			Where * indicates identity and + indicates similarity.	

The term mastocytosis denotes a heterogenous group of disorders characterized by abnormal growth and accumulation of mast cells in one or more organs. Cutaneous and systemic variants of the disease have been described. Mast cell disorders have also been categorized according to other aspects, such as family history, age, course of disease, or presence of a concomitant myeloid neoplasm. However, so far, generally accepted disease

criteria are missing. Recently, a number of diagnostic (disease-related) markers have been identified in mastocytosis research. These include the mast cell enzyme tryptase, CD2, and mast cell growth factor receptor c-kit (CD117). The mast cell enzyme tryptase is increasingly used as a serum- and immunohistochemical marker to estimate the actual spread of disease (burden of neoplastic mast cells). The clinical significance of novel mastocytosis markers is currently under investigation. First results indicate that they may be useful to define reliable criteria for the delineation of the disease.

The NOV13 nucleic acids and proteins of the invention are useful in potential therapeutic applications implicated in disorders characterized by abnormal growth and accumulation of mast cells in one or more organs including, but not limited to skin, ear and brain as well as other pathologies and disorder such as hemophilia, idiopathic thrombocytopenic purpura, autoimmune disease, allergies, immunodeficiencies, transplantation, graft versus host, anemia, ataxia-telangiectasia, lymphedema, tonsillitis, hypercoagulation, and sudden infant death syndrome.

The NOV13 nucleic acid and protein of the invention, or fragments thereof, may further be useful in diagnostic applications, wherein the presence or amount of the NOV13 nucleic acid or the protein are to be assessed.

The polypeptides encoded by NOV8 and NOV11-13 represent a new family of mast cell proteases. ClustalW analysis indicates a very strong homology among these polypeptides, as is shown in Table 50.

TABLE 50.

	NOV12	-----SLGAATSRPGGTP--GRE-----
	NOV11	--MLWLLLLTLPCLMGSVPRNPGE--GRELVGITGGCDVSARRHPWQVSLRFYSMKKG
25	NOV8	MPLLPSRSLVPLSSGKTLVRPPHEPGTGRELVGITGGCDVSARRHPWQVSLRFYSMKKG
	NOV13	-----MGSQRCQ--GGPGTGRELVGITGGCDVSARRHPWQVSLRFYSMKKG
		* ***
	NOV12	-----ELEACAFRVQVQQLRLYEDDQRTKVVEIVRHPQY
30	NOV11	LWEPICGGS LIHPEWVLTAACHL--LEELEACAFRVQVQQLRLYEDDQRTKVVEIVRHPQY
	NOV8	LWEPICGGS LIHPEWVLTAACHLGPEELEACAFRVQVQQLRLYEDDQRTKVVEIVRHPQY
	NOV13	LWEPICGGS LIHPEWVLTAACHLGREELEACAFRVQVQQLRLYEDDQRTKVVEIVRHPQY

35	NOV12	NESLSAQGGADIALLKLEAPVPLSELHPVSLPSASRDVPSGKTCWVTGWGVIGRGELLP
	NOV11	NESLSAQGGADIALLKLEAPVPLSELHPVSLPSASLDVPSGKTCWVTGWGVIGRGELLP
	NOV8	NESLSAQGGADIALLKLEAPVPLSELHPVSLPSASLDVPSGKTCWVTGWGVIGRGELLP
	NOV13	NESLSAQGGADIALLKLEAPVPLSELHPVSLPSASRPGLTQTRPGWLPAAETDQGELLP
40		***** . :. *:.. . . ****

NOV12 WPLSLWEATVKVRSNVLCNQTCRRRFPSNHTEFERFERLIKDDMLCAGDGNHGSWPGDNGGP
 NOV11 WPLSLWEATVKVRSNVLCNQTCRRRFPSNHTEFERFERLIKDDMLCAGDGNHGSWPGDNGGP
 NOV8 WPLSLWEATVKVRSNVLCNQTCRRRFPSNHTEFERFERLIKDDMLCAGDERHLSPPQGDNGGP
 NOV13 WPLSLWEATVKVRSNVLCNQTCRRRFPSNHTEFERFERLIKDDMLCAGDGNHGSWPGDNGGP

5

***** . * * *****

NOV12 LLCRRNCTWVQVEVSWGKLCGLRGYPGMYTRVTSYVSWIRQYVPPFPFR (SEQ ID NO.:24)
 NOV11 LLCRRNCTWVQVEVSWGKLCGLRGYPGMYTRVTSYVSWIRQYVPPFPFR (SEQ ID NO.:22)
 NOV8 LLCRRNCTWVQVEVSWGKLCGLRGYPGMYTRVTSYVSWIRQYVPPFPFR (SEQ ID NO.:16)
 10 NOV13 LLCRRNCTWVQVEVSWGKLCGLRGYPGMYTRVTSYVSWIRQYVPPFPFR (SEQ ID NO.:26)

Where * indicates identity, : indicates strong similarity and . indicates weak similarity.

The nucleic acids and proteins of the invention are useful in potential therapeutic
 15 applications implicated in proliferative disorders, *e.g.* cancer and mastocytosis, immune
 disorders, hepatic disorders, *e.g.* cirrhosis, viral infections, *e.g.* AIDS and hepatitis, and
 disorders of the neuro-olfactory system *e.g.* trauma, surgery and/or neoplastic disorders. For
 example, a cDNA encoding the olfactory receptor protein may be useful in gene therapy for
 treating such disorders, and the olfactory receptor protein may be useful when administered to
 20 a subject in need thereof. By way of nonlimiting example, the compositions of the present
 invention will have efficacy for treatment of patients suffering from disorders of the neuro-
 olfactory system. The novel nucleic acids encoding olfactory receptor protein, and the
 olfactory receptor protein of the invention, or fragments thereof, may further be useful in the
 treatment of adenocarcinoma; lymphoma; prostate cancer; uterus cancer, immune response,
 25 AIDS, asthma, Crohn's disease, multiple sclerosis, treatment of Albright hereditary
 osteodystrophy, development of powerful assay system for functional analysis of various
 human disorders which will help in understanding of pathology of the disease, and
 development of new drug targets for various disorders. They may also be used in diagnostic
 applications, wherein the presence or amount of the nucleic acid or the protein are to be
 30 assessed. These materials are further useful in the generation of antibodies that bind
 immunospecifically to the novel substances of the invention for use in therapeutic or
 diagnostic methods.

NOVX Nucleic Acids

35 The nucleic acids of the invention include those that encode a NOVX polypeptide or
 protein. As used herein, the terms polypeptide and protein are interchangeable.

In some embodiments, a NOVX nucleic acid encodes a mature NOVX polypeptide. As
 used herein, a "mature" form of a polypeptide or protein described herein relates to the product

of a naturally occurring polypeptide or precursor form or proprotein. The naturally occurring polypeptide, precursor or proprotein includes, by way of nonlimiting example, the full-length gene product, encoded by the corresponding gene. Alternatively, it may be defined as the polypeptide, precursor or proprotein encoded by an open reading frame described herein. The product "mature" form arises, again by way of nonlimiting example, as a result of one or more naturally occurring processing steps that may take place within the cell in which the gene product arises. Examples of such processing steps leading to a "mature" form of a polypeptide or protein include the cleavage of the N-terminal methionine residue encoded by the initiation codon of an open reading frame, or the proteolytic cleavage of a signal peptide or leader sequence. Thus a mature form arising from a precursor polypeptide or protein that has residues 1 to N, where residue 1 is the N-terminal methionine, would have residues 2 through N remaining after removal of the N-terminal methionine. Alternatively, a mature form arising from a precursor polypeptide or protein having residues 1 to N, in which an N-terminal signal sequence from residue 1 to residue M is cleaved, would have the residues from residue M+1 to residue N remaining. Further as used herein, a "mature" form of a polypeptide or protein may arise from a step of post-translational modification other than a proteolytic cleavage event. Such additional processes include, by way of non-limiting example, glycosylation, myristoylation or phosphorylation. In general, a mature polypeptide or protein may result from the operation of only one of these processes, or a combination of any of them.

Among the NOVX nucleic acids is the nucleic acid whose sequence is provided in SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25, or a fragment thereof. Additionally, the invention includes mutant or variant nucleic acids of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25, or a fragment thereof, any of whose bases may be changed from the corresponding bases shown in SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25, while still encoding a protein that maintains at least one of its NOVX-like activities and physiological functions (*i.e.*, modulating angiogenesis, neuronal development). The invention further includes the complement of the nucleic acid sequence of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25, including fragments, derivatives, analogs and homologs thereof. The invention additionally includes nucleic acids or nucleic acid fragments, or complements thereto, whose structures include chemical modifications.

One aspect of the invention pertains to isolated nucleic acid molecules that encode NOVX proteins or biologically active portions thereof. Also included are nucleic acid

fragments sufficient for use as hybridization probes to identify NOVX-encoding nucleic acids (e.g., NOVX mRNA) and fragments for use as polymerase chain reaction (PCR) primers for the amplification or mutation of NOVX nucleic acid molecules. As used herein, the term "nucleic acid molecule" is intended to include DNA molecules (e.g., cDNA or genomic DNA),
5 RNA molecules (e.g., mRNA), analogs of the DNA or RNA generated using nucleotide analogs, and derivatives, fragments and homologs thereof. The nucleic acid molecule can be single-stranded or double-stranded, but preferably is double-stranded DNA.

"Probes" refer to nucleic acid sequences of variable length, preferably between at least about 10 nucleotides (nt), 100 nt, or as many as about, e.g., 6,000 nt, depending on use.

10 Probes are used in the detection of identical, similar, or complementary nucleic acid sequences. Longer length probes are usually obtained from a natural or recombinant source, are highly specific and much slower to hybridize than oligomers. Probes may be single- or double-stranded and designed to have specificity in PCR, membrane-based hybridization technologies, or ELISA-like technologies.

15 An "isolated" nucleic acid molecule is one that is separated from other nucleic acid molecules that are present in the natural source of the nucleic acid. Examples of isolated nucleic acid molecules include, but are not limited to, recombinant DNA molecules contained in a vector, recombinant DNA molecules maintained in a heterologous host cell, partially or substantially purified nucleic acid molecules, and synthetic DNA or RNA molecules.

20 Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (*i.e.*, sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated NOVX nucleic acid molecule can contain less than about 50 kb, 25 kb, 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequences which naturally flank the nucleic
25 acid molecule in genomic DNA of the cell from which the nucleic acid is derived. Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material or culture medium when produced by recombinant techniques, or of chemical precursors or other chemicals when chemically synthesized.

A nucleic acid molecule of the present invention, e.g., a nucleic acid molecule having
30 the nucleotide sequence of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25, or a complement of any of this nucleotide sequence, can be isolated using standard molecular biology techniques and the sequence information provided herein. Using all or a portion of the

nucleic acid sequence of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25, as a hybridization probe, NOVX nucleic acid sequences can be isolated using standard hybridization and cloning techniques (*e.g.*, as described in Sambrook *et al.*, eds., MOLECULAR CLONING: A LABORATORY MANUAL 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989; and Ausubel, *et al.*, eds., CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, New York, NY, 1993.)

A nucleic acid of the invention can be amplified using cDNA, mRNA or alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis. Furthermore, oligonucleotides corresponding to NOVX nucleotide sequences can be prepared by standard synthetic techniques, *e.g.*, using an automated DNA synthesizer.

As used herein, the term "oligonucleotide" refers to a series of linked nucleotide residues, which oligonucleotide has a sufficient number of nucleotide bases to be used in a PCR reaction. A short oligonucleotide sequence may be based on, or designed from, a genomic or cDNA sequence and is used to amplify, confirm, or reveal the presence of an identical, similar or complementary DNA or RNA in a particular cell or tissue. Oligonucleotides comprise portions of a nucleic acid sequence having about 10 nt, 50 nt, or 100 nt in length, preferably about 15 nt to 30 nt in length. In one embodiment, an oligonucleotide comprising a nucleic acid molecule less than 100 nt in length would further comprise at least 6 contiguous nucleotides of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25, or a complement thereof. Oligonucleotides may be chemically synthesized and may be used as probes.

In another embodiment, an isolated nucleic acid molecule of the invention comprises a nucleic acid molecule that is a complement of the nucleotide sequence shown in SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25, or a portion of this nucleotide sequence. A nucleic acid molecule that is complementary to the nucleotide sequence shown in SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25 is one that is sufficiently complementary to the nucleotide sequence shown in SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25 that it can hydrogen bond with little or no mismatches to the nucleotide sequence shown in SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25, thereby forming a stable duplex.

As used herein, the term "complementary" refers to Watson-Crick or Hoogsteen base pairing between nucleotide units of a nucleic acid molecule, and the term "binding" means the physical or chemical interaction between two polypeptides or compounds or associated polypeptides or compounds or combinations thereof. Binding includes ionic, non-ionic, Von der Waals, hydrophobic interactions, etc. A physical interaction can be either direct or indirect. Indirect interactions may be through or due to the effects of another polypeptide or compound. Direct binding refers to interactions that do not take place through, or due to, the effect of another polypeptide or compound, but instead are without other substantial chemical intermediates.

Moreover, the nucleic acid molecule of the invention can comprise only a portion of the nucleic acid sequence of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25, *e.g.*, a fragment that can be used as a probe or primer, or a fragment encoding a biologically active portion of NOVX. Fragments provided herein are defined as sequences of at least 6 (contiguous) nucleic acids or at least 4 (contiguous) amino acids, a length sufficient to allow for specific hybridization in the case of nucleic acids or for specific recognition of an epitope in the case of amino acids, respectively, and are at most some portion less than a full length sequence. Fragments may be derived from any contiguous portion of a nucleic acid or amino acid sequence of choice. Derivatives are nucleic acid sequences or amino acid sequences formed from the native compounds either directly or by modification or partial substitution. Analogs are nucleic acid sequences or amino acid sequences that have a structure similar to, but not identical to, the native compound but differs from it in respect to certain components or side chains. Analogs may be synthetic or from a different evolutionary origin and may have a similar or opposite metabolic activity compared to wild type.

Derivatives and analogs may be full length or other than full length, if the derivative or analog contains a modified nucleic acid or amino acid, as described below. Derivatives or analogs of the nucleic acids or proteins of the invention include, but are not limited to, molecules comprising regions that are substantially homologous to the nucleic acids or proteins of the invention, in various embodiments, by at least about 70%, 80%, 85%, 90%, 95%, 98%, or even 99% identity (with a preferred identity of 80-99%) over a nucleic acid or amino acid sequence of identical size or when compared to an aligned sequence in which the alignment is done by a computer homology program known in the art, or whose encoding nucleic acid is capable of hybridizing to the complement of a sequence encoding the

aforementioned proteins under stringent, moderately stringent, or low stringent conditions.

See *e.g.* Ausubel, *et al.*, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, New York, NY, 1993, and below. An exemplary program is the Gap program (Wisconsin

Sequence Analysis Package, Version 8 for UNIX, Genetics Computer Group, University

- 5 Research Park, Madison, WI) using the default settings, which uses the algorithm of Smith and Waterman (Adv. Appl. Math., 1981, 2: 482-489, which is incorporated herein by reference in its entirety).

- A "homologous nucleic acid sequence" or "homologous amino acid sequence," or variations thereof, refer to sequences characterized by a homology at the nucleotide level or
- 10 amino acid level as discussed above. Homologous nucleotide sequences encode those sequences coding for isoforms of a NOVX polypeptide. Isoforms can be expressed in different tissues of the same organism as a result of, for example, alternative splicing of RNA. Alternatively, isoforms can be encoded by different genes. In the present invention, homologous nucleotide sequences include nucleotide sequences encoding for a NOVX
- 15 polypeptide of species other than humans, including, but not limited to, mammals, and thus can include, *e.g.*, mouse, rat, rabbit, dog, cat, cow, horse, and other organisms. Homologous nucleotide sequences also include, but are not limited to, naturally occurring allelic variations and mutations of the nucleotide sequences set forth herein. A homologous nucleotide sequence does not, however, include the nucleotide sequence encoding human NOVX protein.
- 20 Homologous nucleic acid sequences include those nucleic acid sequences that encode conservative amino acid substitutions (see below) in SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26, as well as a polypeptide having NOVX activity. Biological activities of the NOVX proteins are described below. A homologous amino acid sequence does not encode the amino acid sequence of a human NOVX polypeptide.

- 25 The nucleotide sequence determined from the cloning of the human NOVX gene allows for the generation of probes and primers designed for use in identifying and/or cloning NOVX homologues in other cell types, *e.g.*, from other tissues, as well as NOVX homologues from other mammals. The probe/primer typically comprises a substantially purified oligonucleotide. The oligonucleotide typically comprises a region of nucleotide sequence that
- 30 hybridizes under stringent conditions to at least about 12, 25, 50, 100, 150, 200, 250, 300, 350 or 400 or more consecutive sense strand nucleotide sequence of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25; or an anti-sense strand nucleotide sequence of SEQ ID NO: 1, 3,

5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25; or of a naturally occurring mutant of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25.

Probes based on the human NOVX nucleotide sequence can be used to detect transcripts or genomic sequences encoding the same or homologous proteins. In various
5 embodiments, the probe further comprises a label group attached thereto, *e.g.*, the label group can be a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used as a part of a diagnostic test kit for identifying cells or tissue which misexpress a NOVX protein, such as by measuring a level of a NOVX-encoding nucleic acid in a sample of cells from a subject *e.g.*, detecting NOVX mRNA levels or determining whether
10 a genomic NOVX gene has been mutated or deleted.

A "polypeptide having a biologically active portion of NOVX" refers to polypeptides exhibiting activity similar, but not necessarily identical to, an activity of a polypeptide of the present invention, including mature forms, as measured in a particular biological assay, with or without dose dependency. A nucleic acid fragment encoding a "biologically active portion of
15 NOVX" can be prepared by isolating a portion of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25 that encodes a polypeptide having a NOVX biological activity (biological activities of the NOVX proteins are described below), expressing the encoded portion of NOVX protein (*e.g.*, by recombinant expression *in vitro*) and assessing the activity of the encoded portion of NOVX. For example, a nucleic acid fragment encoding a biologically
20 active portion of NOVX can optionally include an ATP-binding domain. In another embodiment, a nucleic acid fragment encoding a biologically active portion of NOVX includes one or more regions.

NOVX Variants

25 The invention further encompasses nucleic acid molecules that differ from the nucleotide sequences shown in SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25 due to the degeneracy of the genetic code. These nucleic acids thus encode the same NOVX protein as that encoded by the nucleotide sequence shown in SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25 *e.g.*, the polypeptide of SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20,
30 22, 24 or 26. In another embodiment, an isolated nucleic acid molecule of the invention has a nucleotide sequence encoding a protein having an amino acid sequence shown in SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26.

In addition to the human NOVX nucleotide sequence shown in SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25, it will be appreciated by those skilled in the art that DNA sequence polymorphisms that lead to changes in the amino acid sequences of NOVX may exist within a population (*e.g.*, the human population). Such genetic polymorphism in the NOVX gene may exist among individuals within a population due to natural allelic variation. As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules comprising an open reading frame encoding a NOVX protein, preferably a mammalian NOVX protein. Such natural allelic variations can typically result in 1-5% variance in the nucleotide sequence of the NOVX gene. Any and all such nucleotide variations and resulting amino acid polymorphisms in NOVX that are the result of natural allelic variation and that do not alter the functional activity of NOVX are intended to be within the scope of the invention.

Moreover, nucleic acid molecules encoding NOVX proteins from other species, and thus that have a nucleotide sequence that differs from the human sequence of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25 are intended to be within the scope of the invention. Nucleic acid molecules corresponding to natural allelic variants and homologues of the NOVX cDNAs of the invention can be isolated based on their homology to the human NOVX nucleic acids disclosed herein using the human cDNAs, or a portion thereof, as a hybridization probe according to standard hybridization techniques under stringent hybridization conditions. For example, a soluble human NOVX cDNA can be isolated based on its homology to human membrane-bound NOVX. Likewise, a membrane-bound human NOVX cDNA can be isolated based on its homology to soluble human NOVX.

Accordingly, in another embodiment, an isolated nucleic acid molecule of the invention is at least 6 nucleotides in length and hybridizes under stringent conditions to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25. In another embodiment, the nucleic acid is at least 10, 25, 50, 100, 250, 500 or 750 nucleotides in length. In another embodiment, an isolated nucleic acid molecule of the invention hybridizes to the coding region. As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences at least 60% homologous to each other typically remain hybridized to each other.

Homologs (*i.e.*, nucleic acids encoding NOVX proteins derived from species other than human) or other related sequences (*e.g.*, paralogs) can be obtained by low, moderate or high

stringency hybridization with all or a portion of the particular human sequence as a probe using methods well known in the art for nucleic acid hybridization and cloning.

As used herein, the phrase "stringent hybridization conditions" refers to conditions under which a probe, primer or oligonucleotide will hybridize to its target sequence, but to no other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures than shorter sequences. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength, pH and nucleic acid concentration) at which 50% of the probes complementary to the target sequence hybridize to the target sequence at equilibrium. Since the target sequences are generally present at excess, at T_m , 50% of the probes are occupied at equilibrium. Typically, stringent conditions will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes, primers or oligonucleotides (*e.g.*, 10 nt to 50 nt) and at least about 60°C for longer probes, primers and oligonucleotides. Stringent conditions may also be achieved with the addition of destabilizing agents, such as formamide.

Stringent conditions are known to those skilled in the art and can be found in CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. Preferably, the conditions are such that sequences at least about 65%, 70%, 75%, 85%, 90%, 95%, 98%, or 99% homologous to each other typically remain hybridized to each other. A non-limiting example of stringent hybridization conditions is hybridization in a high salt buffer comprising 6X SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA, and 500 mg/ml denatured salmon sperm DNA at 65°C. This hybridization is followed by one or more washes in 0.2X SSC, 0.01% BSA at 50°C. An isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to the sequence of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25 corresponds to a naturally occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature (*e.g.*, encodes a natural protein).

In a second embodiment, a nucleic acid sequence that is hybridizable to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17,

19, 21, 23 or 25, or fragments, analogs or derivatives thereof, under conditions of moderate stringency is provided. A non-limiting example of moderate stringency hybridization conditions are hybridization in 6X SSC, 5X Denhardt's solution, 0.5% SDS and 100 mg/ml denatured salmon sperm DNA at 55°C, followed by one or more washes in 1X SSC, 0.1%

5 SDS at 37°C. Other conditions of moderate stringency that may be used are well known in the art. See, *e.g.*, Ausubel *et al.* (eds.), 1993, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, NY, and Kriegler, 1990, GENE TRANSFER AND EXPRESSION, A LABORATORY MANUAL, Stockton Press, NY.

In a third embodiment, a nucleic acid that is hybridizable to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25, or fragments, analogs or derivatives thereof, under conditions of low stringency, is provided. A non-limiting example of low stringency hybridization conditions are hybridization in 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 mg/ml denatured salmon sperm DNA, 10% (wt/vol) dextran sulfate at 40°C, followed by one or more washes in 2X SSC, 25 mM Tris-HCl (pH 7.4), 5 mM EDTA, and 0.1% SDS at 50°C. Other conditions of low stringency that may be used are well known in the art (*e.g.*, as employed for cross-species hybridizations). See, *e.g.*, Ausubel *et al.* (eds.), 1993, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, NY, and Kriegler, 1990, GENE TRANSFER AND EXPRESSION, A LABORATORY MANUAL, Stockton Press, NY; Shilo and Weinberg, 1981, *Proc Natl Acad Sci USA* 78: 6789-6792.

Conservative mutations

In addition to naturally-occurring allelic variants of the NOVX sequence that may exist in the population, the skilled artisan will further appreciate that changes can be introduced by mutation into the nucleotide sequence of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25, thereby leading to changes in the amino acid sequence of the encoded NOVX protein, without altering the functional ability of the NOVX protein. For example, nucleotide substitutions leading to amino acid substitutions at "non-essential" amino acid residues can be made in the sequence of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25. A "non-essential" amino acid residue is a residue that can be altered from the wild-type sequence of NOVX without altering the biological activity, whereas an "essential" amino acid residue is required for biological activity. For example, amino acid residues that are conserved among

the NOVX proteins of the present invention, are predicted to be particularly unamenable to alteration.

Another aspect of the invention pertains to nucleic acid molecules encoding NOVX proteins that contain changes in amino acid residues that are not essential for activity. Such NOVX proteins differ in amino acid sequence from SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26, yet retain biological activity. In one embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence encoding a protein, wherein the protein comprises an amino acid sequence at least about 75% homologous to the amino acid sequence of SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26. Preferably, the protein encoded by the nucleic acid is at least about 80% homologous to SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26, more preferably at least about 90%, 95%, 98%, and most preferably at least about 99% homologous to SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26.

An isolated nucleic acid molecule encoding a NOVX protein homologous to the protein of can be created by introducing one or more nucleotide substitutions, additions or deletions into the nucleotide sequence of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25, such that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein.

Mutations can be introduced into the nucleotide sequence of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25 by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Preferably, conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine). Thus, a predicted nonessential amino acid residue in NOVX is replaced with another amino acid residue from the same side chain family. Alternatively, in another embodiment, mutations can be introduced randomly along all or part of a NOVX coding sequence, such as by saturation mutagenesis, and the

resultant mutants can be screened for NOVX biological activity to identify mutants that retain activity. Following mutagenesis of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25 the encoded protein can be expressed by any recombinant technology known in the art and the activity of the protein can be determined.

- 5 In one embodiment, a mutant NOVX protein can be assayed for (1) the ability to form protein:protein interactions with other NOVX proteins, other cell-surface proteins, or biologically active portions thereof, (2) complex formation between a mutant NOVX protein and a NOVX receptor; (3) the ability of a mutant NOVX protein to bind to an intracellular target protein or biologically active portion thereof; (*e.g.*, avidin proteins); (4) the ability to
10 bind NOVX protein; or (5) the ability to specifically bind an anti-NOVX protein antibody.

Antisense NOVX Nucleic Acids

- Another aspect of the invention pertains to isolated antisense nucleic acid molecules that are hybridizable to or complementary to the nucleic acid molecule comprising the
15 nucleotide sequence of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25, or fragments, analogs or derivatives thereof. An "antisense" nucleic acid comprises a nucleotide sequence that is complementary to a "sense" nucleic acid encoding a protein, *e.g.*, complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence. In specific aspects, antisense nucleic acid molecules are provided that comprise a sequence
20 complementary to at least about 10, 25, 50, 100, 250 or 500 nucleotides or an entire NOVX coding strand, or to only a portion thereof. Nucleic acid molecules encoding fragments, homologs, derivatives and analogs of a NOVX protein of SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26 or antisense nucleic acids complementary to a NOVX nucleic acid sequence of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25 are additionally
25 provided.

- In one embodiment, an antisense nucleic acid molecule is antisense to a "coding region" of the coding strand of a nucleotide sequence encoding NOVX. The term "coding region" refers to the region of the nucleotide sequence comprising codons which are translated into amino acid residues (*e.g.*, the protein coding region of human NOVX corresponds to SEQ
30 ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26). In another embodiment, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence encoding NOVX. The term "noncoding region" refers to 5' and 3' sequences which

flank the coding region that are not translated into amino acids (*i.e.*, also referred to as 5' and 3' untranslated regions).

Given the coding strand sequences encoding NOVX disclosed herein (*e.g.*, SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25), antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick or Hoogsteen base pairing. The antisense nucleic acid molecule can be complementary to the entire coding region of NOVX mRNA, but more preferably is an oligonucleotide that is antisense to only a portion of the coding or noncoding region of NOVX mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of NOVX mRNA. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45 or 50 nucleotides in length. An antisense nucleic acid of the invention can be constructed using chemical synthesis or enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (*e.g.*, an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, *e.g.*, phosphorothioate derivatives and acridine substituted nucleotides can be used.

Examples of modified nucleotides that can be used to generate the antisense nucleic acid include: 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (*i.e.*, RNA transcribed from the

inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

The antisense nucleic acid molecules of the invention are typically administered to a subject or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a NOVX protein to thereby inhibit expression of the protein, *e.g.*, by inhibiting transcription and/or translation. The hybridization can be by conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule that binds to DNA duplexes, through specific interactions in the major groove of the double helix. An example of a route of administration of antisense nucleic acid molecules of the invention includes direct injection at a tissue site. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then administered systemically. For example, for systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface, *e.g.*, by linking the antisense nucleic acid molecules to peptides or antibodies that bind to cell surface receptors or antigens. The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular concentrations of antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

In yet another embodiment, the antisense nucleic acid molecule of the invention is an α -anomeric nucleic acid molecule. An α -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gaultier *et al.* (1987) *Nucleic Acids Res* 15: 6625-6641). The antisense nucleic acid molecule can also comprise a 2'-O-methylribonucleotide (Inoue *et al.* (1987) *Nucleic Acids Res* 15: 6131-6148) or a chimeric RNA-DNA analogue (Inoue *et al.* (1987) *FEBS Lett* 215: 327-330).

Such modifications include, by way of nonlimiting example, modified bases, and nucleic acids whose sugar phosphate backbones are modified or derivatized. These modifications are carried out at least in part to enhance the chemical stability of the modified nucleic acid, such that they may be used, for example, as antisense binding nucleic acids in therapeutic applications in a subject.

NOVX Ribozymes and PNA moieties

In still another embodiment, an antisense nucleic acid of the invention is a ribozyme. Ribozymes are catalytic RNA molecules with ribonuclease activity that are capable of cleaving a single-stranded nucleic acid, such as a mRNA, to which they have a complementary region.

5 Thus, ribozymes (*e.g.*, hammerhead ribozymes (described in Haselhoff and Gerlach (1988) *Nature* 334:585-591)) can be used to catalytically cleave NOVX mRNA transcripts to thereby inhibit translation of NOVX mRNA. A ribozyme having specificity for a NOVX-encoding nucleic acid can be designed based upon the nucleotide sequence of a NOVX DNA disclosed herein (*i.e.*, SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25). For example, a

10 derivative of a Tetrahymena L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved in a NOVX-encoding mRNA. See, *e.g.*, Cech *et al.* U.S. Pat. No. 4,987,071; and Cech *et al.* U.S. Pat. No. 5,116,742. Alternatively, NOVX mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See, *e.g.*, Bartel *et al.*,

15 (1993) *Science* 261:1411-1418.

Alternatively, NOVX gene expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region of the NOVX (*e.g.*, the NOVX promoter and/or enhancers) to form triple helical structures that prevent transcription of the NOVX gene in target cells. See generally, Helene. (1991) *Anticancer Drug Des.* 6: 569-84; Helene. *et al.*

20 (1992) *Ann. N.Y. Acad. Sci.* 660:27-36; and Maher (1992) *Bioassays* 14: 807-15.

In various embodiments, the nucleic acids of NOVX can be modified at the base moiety, sugar moiety or phosphate backbone to improve, *e.g.*, the stability, hybridization, or solubility of the molecule. For example, the deoxyribose phosphate backbone of the nucleic acids can be modified to generate peptide nucleic acids (see Hyrup *et al.* (1996) *Bioorg Med*

25 *Chem* 4: 5-23). As used herein, the terms "peptide nucleic acids" or "PNAs" refer to nucleic acid mimics, *e.g.*, DNA mimics, in which the deoxyribose phosphate backbone is replaced by a pseudopeptide backbone and only the four natural nucleobases are retained. The neutral backbone of PNAs has been shown to allow for specific hybridization to DNA and RNA under conditions of low ionic strength. The synthesis of PNA oligomers can be performed using

30 standard solid phase peptide synthesis protocols as described in Hyrup *et al.* (1996) above; Perry-O'Keefe *et al.* (1996) *PNAS* 93: 14670-675.

PNAs of NOVX can be used in therapeutic and diagnostic applications. For example, PNAs can be used as antisense or antigene agents for sequence-specific modulation of gene expression by, *e.g.*, inducing transcription or translation arrest or inhibiting replication. PNAs of NOVX can also be used, *e.g.*, in the analysis of single base pair mutations in a gene by, *e.g.*,
5 PNA directed PCR clamping; as artificial restriction enzymes when used in combination with other enzymes, *e.g.*, S1 nucleases (Hyrup B. (1996) above); or as probes or primers for DNA sequence and hybridization (Hyrup *et al.* (1996), above; Perry-O'Keefe (1996), above).

In another embodiment, PNAs of NOVX can be modified, *e.g.*, to enhance their stability or cellular uptake, by attaching lipophilic or other helper groups to PNA, by the
10 formation of PNA-DNA chimeras, or by the use of liposomes or other techniques of drug delivery known in the art. For example, PNA-DNA chimeras of NOVX can be generated that may combine the advantageous properties of PNA and DNA. Such chimeras allow DNA recognition enzymes, *e.g.*, RNase H and DNA polymerases, to interact with the DNA portion while the PNA portion would provide high binding affinity and specificity. PNA-DNA
15 chimeras can be linked using linkers of appropriate lengths selected in terms of base stacking, number of bonds between the nucleobases, and orientation (Hyrup (1996) above). The synthesis of PNA-DNA chimeras can be performed as described in Hyrup (1996) above and Finn *et al.* (1996) *Nucl Acids Res* 24: 3357-63. For example, a DNA chain can be synthesized on a solid support using standard phosphoramidite coupling chemistry, and modified
20 nucleoside analogs, *e.g.*, 5'-(4-methoxytrityl) amino-5'-deoxy-thymidine phosphoramidite, can be used between the PNA and the 5' end of DNA (Mag *et al.* (1989) *Nucl Acid Res* 17: 5973-88). PNA monomers are then coupled in a stepwise manner to produce a chimeric molecule with a 5' PNA segment and a 3' DNA segment (Finn *et al.* (1996) above). Alternatively, chimeric molecules can be synthesized with a 5' DNA segment and a 3' PNA
25 segment. See, Petersen *et al.* (1975) *Bioorg Med Chem Lett* 5: 1119-11124.

In other embodiments, the oligonucleotide may include other appended groups such as peptides (*e.g.*, for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (see, *e.g.*, Letsinger *et al.*, 1989, *Proc. Natl. Acad. Sci. U.S.A.* 86:6553-6556; Lemaitre *et al.*, 1987, *Proc. Natl. Acad. Sci.* 84:648-652; PCT Publication No.
30 W088/09810) or the blood-brain barrier (see, *e.g.*, PCT Publication No. W089/10134). In addition, oligonucleotides can be modified with hybridization triggered cleavage agents (See, *e.g.*, Krol *et al.*, 1988, *BioTechniques* 6:958-976) or intercalating agents. (See, *e.g.*, Zon, 1988,

Pharm. Res. 5: 539-549). To this end, the oligonucleotide may be conjugated to another molecule, *e.g.*, a peptide, a hybridization triggered cross-linking agent, a transport agent, a hybridization-triggered cleavage agent, etc.

5 NOVX Polypeptides

A NOVX polypeptide of the invention includes the NOVX-like protein whose sequence is provided in SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26. The invention also includes a mutant or variant protein any of whose residues may be changed from the corresponding residue shown in SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24
10 or 26 while still encoding a protein that maintains its NOVX-like activities and physiological functions, or a functional fragment thereof. In some embodiments, up to 20% or more of the residues may be so changed in the mutant or variant protein. In some embodiments, the NOVX polypeptide according to the invention is a mature polypeptide.

In general, a NOVX -like variant that preserves NOVX-like function includes any
15 variant in which residues at a particular position in the sequence have been substituted by other amino acids, and further include the possibility of inserting an additional residue or residues between two residues of the parent protein as well as the possibility of deleting one or more residues from the parent sequence. Any amino acid substitution, insertion, or deletion is encompassed by the invention. In favorable circumstances, the substitution is a conservative
20 substitution as defined above.

One aspect of the invention pertains to isolated NOVX proteins, and biologically active portions thereof, or derivatives, fragments, analogs or homologs thereof. Also provided are polypeptide fragments suitable for use as immunogens to raise anti-NOVX antibodies. In one embodiment, native NOVX proteins can be isolated from cells or tissue sources by an
25 appropriate purification scheme using standard protein purification techniques. In another embodiment, NOVX proteins are produced by recombinant DNA techniques. Alternative to recombinant expression, a NOVX protein or polypeptide can be synthesized chemically using standard peptide synthesis techniques.

An "isolated" or "purified" protein or biologically active portion thereof is substantially
30 free of cellular material or other contaminating proteins from the cell or tissue source from which the NOVX protein is derived, or substantially free from chemical precursors or other chemicals when chemically synthesized. The language "substantially free of cellular material"

includes preparations of NOVX protein in which the protein is separated from cellular components of the cells from which it is isolated or recombinantly produced. In one embodiment, the language "substantially free of cellular material" includes preparations of NOVX protein having less than about 30% (by dry weight) of non-NOVX protein (also referred to herein as a "contaminating protein"), more preferably less than about 20% of non-NOVX protein, still more preferably less than about 10% of non-NOVX protein, and most preferably less than about 5% non-NOVX protein. When the NOVX protein or biologically active portion thereof is recombinantly produced, it is also preferably substantially free of culture medium, *i.e.*, culture medium represents less than about 20%, more preferably less than about 10%, and most preferably less than about 5% of the volume of the protein preparation.

The language "substantially free of chemical precursors or other chemicals" includes preparations of NOVX protein in which the protein is separated from chemical precursors or other chemicals that are involved in the synthesis of the protein. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of NOVX protein having less than about 30% (by dry weight) of chemical precursors or non-NOVX chemicals, more preferably less than about 20% chemical precursors or non-NOVX chemicals, still more preferably less than about 10% chemical precursors or non-NOVX chemicals, and most preferably less than about 5% chemical precursors or non-NOVX chemicals.

Biologically active portions of a NOVX protein include peptides comprising amino acid sequences sufficiently homologous to or derived from the amino acid sequence of the NOVX protein, *e.g.*, the amino acid sequence shown in SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26 that include fewer amino acids than the full length NOVX proteins, and exhibit at least one activity of a NOVX protein. Typically, biologically active portions comprise a domain or motif with at least one activity of the NOVX protein. A biologically active portion of a NOVX protein can be a polypeptide which is, for example, 10, 25, 50, 100 or more amino acids in length.

A biologically active portion of a NOVX protein of the present invention may contain at least one of the above-identified domains conserved between the NOVX proteins, *e.g.* TSR modules. Moreover, other biologically active portions, in which other regions of the protein are deleted, can be prepared by recombinant techniques and evaluated for one or more of the functional activities of a native NOVX protein.

In an embodiment, the NOVX protein has an amino acid sequence shown in SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26. In other embodiments, the NOVX protein is substantially homologous to SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26 and retains the functional activity of the protein of SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26 yet differs in amino acid sequence due to natural allelic variation or mutagenesis, as described in detail below. Accordingly, in another embodiment, the NOVX protein is a protein that comprises an amino acid sequence at least about 45% homologous to the amino acid sequence of SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26 and retains the functional activity of the NOVX proteins of SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26.

Determining homology between two or more sequence

To determine the percent homology of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in either of the sequences being compared for optimal alignment between the sequences). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are homologous at that position (*i.e.*, as used herein amino acid or nucleic acid "homology" is equivalent to amino acid or nucleic acid "identity").

The nucleic acid sequence homology may be determined as the degree of identity between two sequences. The homology may be determined using computer programs known in the art, such as GAP software provided in the GCG program package. See, *Needleman and Wunsch* 1970 *J Mol Biol* 48: 443-453. Using GCG GAP software with the following settings for nucleic acid sequence comparison: GAP creation penalty of 5.0 and GAP extension penalty of 0.3, the coding region of the analogous nucleic acid sequences referred to above exhibits a degree of identity preferably of at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, or 99%, with the CDS (encoding) part of the DNA sequence shown in SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25.

The term "sequence identity" refers to the degree to which two polynucleotide or polypeptide sequences are identical on a residue-by-residue basis over a particular region of comparison. The term "percentage of sequence identity" is calculated by comparing two

optimally aligned sequences over that region of comparison, determining the number of positions at which the identical nucleic acid base (*e.g.*, A, T, C, G, U, or I, in the case of nucleic acids) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the region of comparison (*i.e.*, the window size), and multiplying the result by 100 to yield the percentage of sequence identity. The term "substantial identity" as used herein denotes a characteristic of a polynucleotide sequence, wherein the polynucleotide comprises a sequence that has at least 80 percent sequence identity, preferably at least 85 percent identity and often 90 to 95 percent sequence identity, more usually at least 99 percent sequence identity as compared to a reference sequence over a comparison region. The term "percentage of positive residues" is calculated by comparing two optimally aligned sequences over that region of comparison, determining the number of positions at which the identical and conservative amino acid substitutions, as defined above, occur in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the region of comparison (*i.e.*, the window size), and multiplying the result by 100 to yield the percentage of positive residues.

Chimeric and fusion proteins

The invention also provides NOVX chimeric or fusion proteins. As used herein, a NOVX "chimeric protein" or "fusion protein" comprises a NOVX polypeptide operatively linked to a non-NOVX polypeptide. An "NOVX polypeptide" refers to a polypeptide having an amino acid sequence corresponding to NOVX, whereas a "non-NOVX polypeptide" refers to a polypeptide having an amino acid sequence corresponding to a protein that is not substantially homologous to the NOVX protein, *e.g.*, a protein that is different from the NOVX protein and that is derived from the same or a different organism. Within a NOVX fusion protein the NOVX polypeptide can correspond to all or a portion of a NOVX protein. In one embodiment, a NOVX fusion protein comprises at least one biologically active portion of a NOVX protein. In another embodiment, a NOVX fusion protein comprises at least two biologically active portions of a NOVX protein. Within the fusion protein, the term "operatively linked" is intended to indicate that the NOVX polypeptide and the non-NOVX polypeptide are fused in-frame to each other. The non-NOVX polypeptide can be fused to the N-terminus or C-terminus of the NOVX polypeptide.

For example, in one embodiment a NOVX fusion protein comprises a NOVX polypeptide operably linked to the extracellular domain of a second protein. Such fusion proteins can be further utilized in screening assays for compounds that modulate NOVX activity (such assays are described in detail below).

- 5 In another embodiment, the fusion protein is a GST-NOVX fusion protein in which the NOVX sequences are fused to the C-terminus of the GST (*i.e.*, glutathione S-transferase) sequences. Such fusion proteins can facilitate the purification of recombinant NOVX.

- In another embodiment, the fusion protein is a NOVX-immunoglobulin fusion protein in which the NOVX sequences comprising one or more domains are fused to sequences
10 derived from a member of the immunoglobulin protein family. The NOVX-immunoglobulin fusion proteins of the invention can be incorporated into pharmaceutical compositions and administered to a subject to inhibit an interaction between a NOVX ligand and a NOVX protein on the surface of a cell, to thereby suppress NOVX-mediated signal transduction *in vivo*. In one nonlimiting example, a contemplated NOVX ligand of the invention is the
15 NOVX receptor. The NOVX-immunoglobulin fusion proteins can be used to affect the bioavailability of a NOVX cognate ligand. Inhibition of the NOVX ligand/NOVX interaction may be useful therapeutically for both the treatment of proliferative and differentiative disorders, *e.g.*, cancer as well as modulating (*e.g.*, promoting or inhibiting) cell survival, as well as acute and chronic inflammatory disorders and hyperplastic wound healing, *e.g.*
20 hypertrophic scars and keloids. Moreover, the NOVX-immunoglobulin fusion proteins of the invention can be used as immunogens to produce anti-NOVX antibodies in a subject, to purify NOVX ligands, and in screening assays to identify molecules that inhibit the interaction of NOVX with a NOVX ligand.

- A NOVX chimeric or fusion protein of the invention can be produced by standard
25 recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques, *e.g.*, by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In
30 another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers that give rise to complementary overhangs between two

consecutive gene fragments that can subsequently be annealed and reamplified to generate a chimeric gene sequence (see, for example, Ausubel et al. (eds.) CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (*e.g.*, a GST polypeptide). A
5 NOVX-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the NOVX protein.

NOVX agonists and antagonists

The present invention also pertains to variants of the NOVX proteins that function as
10 either NOVX agonists (mimetics) or as NOVX antagonists. Variants of the NOVX protein can be generated by mutagenesis, *e.g.*, discrete point mutation or truncation of the NOVX protein. An agonist of the NOVX protein can retain substantially the same, or a subset of, the biological activities of the naturally occurring form of the NOVX protein. An antagonist of the NOVX protein can inhibit one or more of the activities of the naturally occurring form of
15 the NOVX protein by, for example, competitively binding to a downstream or upstream member of a cellular signaling cascade which includes the NOVX protein. Thus, specific biological effects can be elicited by treatment with a variant of limited function. In one embodiment, treatment of a subject with a variant having a subset of the biological activities of the naturally occurring form of the protein has fewer side effects in a subject relative to
20 treatment with the naturally occurring form of the NOVX proteins.

Variants of the NOVX protein that function as either NOVX agonists (mimetics) or as NOVX antagonists can be identified by screening combinatorial libraries of mutants, *e.g.*, truncation mutants, of the NOVX protein for NOVX protein agonist or antagonist activity. In one embodiment, a variegated library of NOVX variants is generated by combinatorial
25 mutagenesis at the nucleic acid level and is encoded by a variegated gene library. A variegated library of NOVX variants can be produced by, for example, enzymatically ligating a mixture of synthetic oligonucleotides into gene sequences such that a degenerate set of potential NOVX sequences is expressible as individual polypeptides, or alternatively, as a set of larger fusion proteins (*e.g.*, for phage display) containing the set of NOVX sequences
30 therein. There are a variety of methods which can be used to produce libraries of potential NOVX variants from a degenerate oligonucleotide sequence. Chemical synthesis of a degenerate gene sequence can be performed in an automatic DNA synthesizer, and the

synthetic gene then ligated into an appropriate expression vector. Use of a degenerate set of genes allows for the provision, in one mixture, of all of the sequences encoding the desired set of potential NOVX sequences. Methods for synthesizing degenerate oligonucleotides are known in the art (see, e.g., Narang (1983) *Tetrahedron* 39:3; Itakura *et al.* (1984) *Annu Rev Biochem* 53:323; Itakura *et al.* (1984) *Science* 198:1056; Ike *et al.* (1983) *Nucl Acid Res* 11:477.

Polypeptide libraries

In addition, libraries of fragments of the NOVX protein coding sequence can be used to generate a variegated population of NOVX fragments for screening and subsequent selection of variants of a NOVX protein. In one embodiment, a library of coding sequence fragments can be generated by treating a double stranded PCR fragment of a NOVX coding sequence with a nuclease under conditions wherein nicking occurs only about once per molecule, denaturing the double stranded DNA, renaturing the DNA to form double stranded DNA that can include sense/antisense pairs from different nicked products, removing single stranded portions from reformed duplexes by treatment with S1 nuclease, and ligating the resulting fragment library into an expression vector. By this method, an expression library can be derived which encodes N-terminal and internal fragments of various sizes of the NOVX protein.

Several techniques are known in the art for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene products having a selected property. Such techniques are adaptable for rapid screening of the gene libraries generated by the combinatorial mutagenesis of NOVX proteins. The most widely used techniques, which are amenable to high throughput analysis, for screening large gene libraries typically include cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and expressing the combinatorial genes under conditions in which detection of a desired activity facilitates isolation of the vector encoding the gene whose product was detected. Recursive ensemble mutagenesis (REM), a new technique that enhances the frequency of functional mutants in the libraries, can be used in combination with the screening assays to identify NOVX variants (Arkin and Yourvan (1992) *PNAS* 89:7811-7815; Delgrave *et al.* (1993) *Protein Engineering* 6:327-331).

NOVX Antibodies

Also included in the invention are antibodies to NOVX proteins, or fragments of NOVX proteins. The term "antibody" as used herein refers to immunoglobulin molecules and immunologically active portions of immunoglobulin (Ig) molecules, i.e., molecules that contain an antigen binding site that specifically binds (immunoreacts with) an antigen. Such antibodies include, but are not limited to, polyclonal, monoclonal, chimeric, single chain, F_{ab} , F_{ab} , and $F_{(ab)2}$ fragments, and an F_{ab} expression library. In general, an antibody molecule obtained from humans relates to any of the classes IgG, IgM, IgA, IgE and IgD, which differ from one another by the nature of the heavy chain present in the molecule. Certain classes have subclasses as well, such as IgG₁, IgG₂, and others. Furthermore, in humans, the light chain may be a kappa chain or a lambda chain. Reference herein to antibodies includes a reference to all such classes, subclasses and types of human antibody species.

An isolated NOVX-related protein of the invention may be intended to serve as an antigen, or a portion or fragment thereof, and additionally can be used as an immunogen to generate antibodies that immunospecifically bind the antigen, using standard techniques for polyclonal and monoclonal antibody preparation. The full-length protein can be used or, alternatively, the invention provides antigenic peptide fragments of the antigen for use as immunogens. An antigenic peptide fragment comprises at least 6 amino acid residues of the amino acid sequence of the full length protein, such as an amino acid sequence shown in from SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26, and encompasses an epitope thereof such that an antibody raised against the peptide forms a specific immune complex with the full length protein or with any fragment that contains the epitope. Preferably, the antigenic peptide comprises at least 10 amino acid residues, or at least 15 amino acid residues, or at least 20 amino acid residues, or at least 30 amino acid residues. Preferred epitopes encompassed by the antigenic peptide are regions of the protein that are located on its surface; commonly these are hydrophilic regions.

In certain embodiments of the invention, at least one epitope encompassed by the antigenic peptide is a region of NOVX-related protein that is located on the surface of the protein, *e.g.*, a hydrophilic region. A hydrophobicity analysis of the human NOVX-related protein sequence will indicate which regions of a NOVX-related protein are particularly hydrophilic and, therefore, are likely to encode surface residues useful for targeting antibody

production. As a means for targeting antibody production, hydropathy plots showing regions of hydrophilicity and hydrophobicity may be generated by any method well known in the art, including, for example, the Kyte Doolittle or the Hopp Woods methods, either with or without Fourier transformation. See, *e.g.*, Hopp and Woods, 1981, *Proc. Nat. Acad. Sci. USA* 78:

5 3824-3828; Kyte and Doolittle 1982, *J. Mol. Biol.* 157: 105-142, each of which is incorporated herein by reference in its entirety. Antibodies that are specific for one or more domains within an antigenic protein, or derivatives, fragments, analogs or homologs thereof, are also provided herein.

A protein of the invention, or a derivative, fragment, analog, homolog or ortholog
10 thereof, may be utilized as an immunogen in the generation of antibodies that immunospecifically bind these protein components.

Various procedures known within the art may be used for the production of polyclonal or monoclonal antibodies directed against a protein of the invention, or against derivatives, fragments, analogs homologs or orthologs thereof (see, for example, *Antibodies: A Laboratory*
15 *Manual*, Harlow E, and Lane D, 1988, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, incorporated herein by reference). Some of these antibodies are discussed below.

Polyclonal Antibodies

For the production of polyclonal antibodies, various suitable host animals (*e.g.*, rabbit,
20 goat, mouse or other mammal) may be immunized by one or more injections with the native protein, a synthetic variant thereof, or a derivative of the foregoing. An appropriate immunogenic preparation can contain, for example, the naturally occurring immunogenic protein, a chemically synthesized polypeptide representing the immunogenic protein, or a recombinantly expressed immunogenic protein. Furthermore, the protein may be conjugated
25 to a second protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. The preparation can further include an adjuvant. Various adjuvants used to increase the immunological response include, but are not limited to, Freund's (complete and incomplete), mineral gels (*e.g.*, aluminum
30 hydroxide), surface active substances (*e.g.*, lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, dinitrophenol, etc.), adjuvants usable in humans such as Bacille Calmette-Guerin and *Corynebacterium parvum*, or similar immunostimulatory agents.

Additional examples of adjuvants which can be employed include MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate).

The polyclonal antibody molecules directed against the immunogenic protein can be isolated from the mammal (e.g., from the blood) and further purified by well known techniques, such as affinity chromatography using protein A or protein G, which provide primarily the IgG fraction of immune serum. Subsequently, or alternatively, the specific antigen which is the target of the immunoglobulin sought, or an epitope thereof, may be immobilized on a column to purify the immune specific antibody by immunoaffinity chromatography. Purification of immunoglobulins is discussed, for example, by D. Wilkinson (The Scientist, published by The Scientist, Inc., Philadelphia PA, Vol. 14, No. 8 (April 17, 2000), pp. 25-28).

Monoclonal Antibodies

The term "monoclonal antibody" (MAb) or "monoclonal antibody composition", as used herein, refers to a population of antibody molecules that contain only one molecular species of antibody molecule consisting of a unique light chain gene product and a unique heavy chain gene product. In particular, the complementarity determining regions (CDRs) of the monoclonal antibody are identical in all the molecules of the population. MAbs thus contain an antigen binding site capable of immunoreacting with a particular epitope of the antigen characterized by a unique binding affinity for it.

Monoclonal antibodies can be prepared using hybridoma methods, such as those described by Kohler and Milstein, Nature, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes can be immunized in vitro.

The immunizing agent will typically include the protein antigen, a fragment thereof or a fusion protein thereof. Generally, either peripheral blood lymphocytes are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, Monoclonal Antibodies: Principles and Practice, Academic Press, (1986) pp. 59-103). Immortalized cell lines are usually transformed mammalian cells, particularly myeloma

cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells can be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine
5 phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

Preferred immortalized cell lines are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a
10 medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, California and the American Type Culture Collection, Manassas, Virginia. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies (Kozbor, J. Immunol., 133:3001 (1984); Brodeur
15 et al., Monoclonal Antibody Production Techniques and Applications, Marcel Dekker, Inc., New York, (1987) pp. 51-63).

The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against the antigen. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by
20 immunoprecipitation or by an in vitro binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, Anal. Biochem., 107:220 (1980). Preferably, antibodies having a high degree of specificity and a high binding affinity for the target antigen
25 are isolated.

After the desired hybridoma cells are identified, the clones can be subcloned by limiting dilution procedures and grown by standard methods. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively, the hybridoma cells can be grown *in vivo* as ascites in a mammal.

30 The monoclonal antibodies secreted by the subclones can be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures

such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

The monoclonal antibodies can also be made by recombinant DNA methods, such as those described in U.S. Patent No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA can be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also can be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences (U.S. Patent No. 4,816,567; Morrison, Nature 368, 812-13 (1994)) or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

20 Humanized Antibodies

The antibodies directed against the protein antigens of the invention can further comprise humanized antibodies or human antibodies. These antibodies are suitable for administration to humans without engendering an immune response by the human against the administered immunoglobulin. Humanized forms of antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) that are principally comprised of the sequence of a human immunoglobulin, and contain minimal sequence derived from a non-human immunoglobulin. Humanization can be performed following the method of Winter and co-workers (Jones et al., Nature, 321:522-525 (1986); Riechmann et al., Nature, 332:323-327 (1988); Verhoeven et al., Science, 239:1534-1536 (1988)), by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. (See also U.S. Patent No. 5,225,539.) In some instances, Fv framework residues of the human immunoglobulin are replaced by

corresponding non-human residues. Humanized antibodies can also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the framework regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin (Jones et al., 1986; Riechmann et al., 1988; and Presta, Curr. Op. Struct. Biol., 2:593-596 (1992)).

Human Antibodies

Fully human antibodies relate to antibody molecules in which essentially the entire sequences of both the light chain and the heavy chain, including the CDRs, arise from human genes. Such antibodies are termed "human antibodies", or "fully human antibodies" herein. Human monoclonal antibodies can be prepared by the trioma technique; the human B-cell hybridoma technique (see Kozbor, et al., 1983 *Immunol Today* 4: 72) and the EBV hybridoma technique to produce human monoclonal antibodies (see Cole, et al., 1985 In: *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc., pp. 77-96). Human monoclonal antibodies may be utilized in the practice of the present invention and may be produced by using human hybridomas (see Cote, et al., 1983. *Proc Natl Acad Sci USA* 80: 2026-2030) or by transforming human B-cells with Epstein Barr Virus in vitro (see Cole, et al., 1985 In: *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc., pp. 77-96).

In addition, human antibodies can also be produced using additional techniques, including phage display libraries (Hoogenboom and Winter, J. Mol. Biol., 227:381 (1991); Marks et al., J. Mol. Biol., 222:581 (1991)). Similarly, human antibodies can be made by introducing human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in Marks et al. (Bio/Technology 10, 779-783 (1992)); Lonberg et al. (Nature 368 856-859 (1994)); Morrison (Nature 368, 812-13 (1994)); Fishwild

et al, (Nature Biotechnology 14, 845-51 (1996)); Neuberger (Nature Biotechnology 14, 826 (1996)); and Lonberg and Huszar (Intern. Rev. Immunol. 13 65-93 (1995)).

Human antibodies may additionally be produced using transgenic nonhuman animals which are modified so as to produce fully human antibodies rather than the animal's endogenous antibodies in response to challenge by an antigen. (See PCT publication WO94/02602). The endogenous genes encoding the heavy and light immunoglobulin chains in the nonhuman host have been incapacitated, and active loci encoding human heavy and light chain immunoglobulins are inserted into the host's genome. The human genes are incorporated, for example, using yeast artificial chromosomes containing the requisite human DNA segments. An animal which provides all the desired modifications is then obtained as progeny by crossbreeding intermediate transgenic animals containing fewer than the full complement of the modifications. The preferred embodiment of such a nonhuman animal is a mouse, and is termed the XenomouseTM as disclosed in PCT publications WO 96/33735 and WO 96/34096. This animal produces B cells which secrete fully human immunoglobulins. The antibodies can be obtained directly from the animal after immunization with an immunogen of interest, as, for example, a preparation of a polyclonal antibody, or alternatively from immortalized B cells derived from the animal, such as hybridomas producing monoclonal antibodies. Additionally, the genes encoding the immunoglobulins with human variable regions can be recovered and expressed to obtain the antibodies directly, or can be further modified to obtain analogs of antibodies such as, for example, single chain Fv molecules.

An example of a method of producing a nonhuman host, exemplified as a mouse, lacking expression of an endogenous immunoglobulin heavy chain is disclosed in U.S. Patent No. 5,939,598. It can be obtained by a method including deleting the J segment genes from at least one endogenous heavy chain locus in an embryonic stem cell to prevent rearrangement of the locus and to prevent formation of a transcript of a rearranged immunoglobulin heavy chain locus, the deletion being effected by a targeting vector containing a gene encoding a selectable marker; and producing from the embryonic stem cell a transgenic mouse whose somatic and germ cells contain the gene encoding the selectable marker.

A method for producing an antibody of interest, such as a human antibody, is disclosed in U.S. Patent No. 5,916,771. It includes introducing an expression vector that contains a nucleotide sequence encoding a heavy chain into one mammalian host cell in culture,

introducing an expression vector containing a nucleotide sequence encoding a light chain into another mammalian host cell, and fusing the two cells to form a hybrid cell. The hybrid cell expresses an antibody containing the heavy chain and the light chain.

In a further improvement on this procedure, a method for identifying a clinically
5 relevant epitope on an immunogen, and a correlative method for selecting an antibody that binds immunospecifically to the relevant epitope with high affinity, are disclosed in PCT publication WO 99/53049.

F_{ab} Fragments and Single Chain Antibodies

10 According to the invention, techniques can be adapted for the production of single-chain antibodies specific to an antigenic protein of the invention (see e.g., U.S. Patent No. 4,946,778). In addition, methods can be adapted for the construction of F_{ab} expression libraries (see e.g., Huse, et al., 1989 Science 246: 1275-1281) to allow rapid and effective
15 identification of monoclonal F_{ab} fragments with the desired specificity for a protein or derivatives, fragments, analogs or homologs thereof. Antibody fragments that contain the idiotypes to a protein antigen may be produced by techniques known in the art including, but not limited to: (i) an F_{(ab)₂} fragment produced by pepsin digestion of an antibody molecule; (ii) an F_{ab} fragment generated by reducing the disulfide bridges of an F_{(ab)₂} fragment; (iii) an F_{ab} fragment generated by the treatment of the antibody molecule with papain and a reducing
20 agent and (iv) F_v fragments.

Bispecific Antibodies

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the
25 binding specificities is for an antigenic protein of the invention. The second binding target is any other antigen, and advantageously is a cell-surface protein or receptor or receptor subunit.

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different
30 specificities (Milstein and Cuello, *Nature*, 305:537-539 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct

bispecific structure. The purification of the correct molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed in WO 93/08829, published 13 May 1993, and in Traunecker *et al.*, 1991 *EMBO J.*, 10:3655-3659.

Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. For further details of generating bispecific antibodies see, for example, Suresh *et al.*, Methods in Enzymology, 121:210 (1986).

According to another approach described in WO 96/27011, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the CH3 region of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g. tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g. F(ab')₂ bispecific antibodies). Techniques for generating bispecific antibodies from antibody fragments have been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan *et al.*, Science 229:81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative

to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

Additionally, Fab' fragments can be directly recovered from *E. coli* and chemically coupled to form bispecific antibodies. Shalaby et al., J. Exp. Med. 175:217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')₂ molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling in vitro to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

Various techniques for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny et al., J. Immunol. 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger et al., Proc. Natl. Acad. Sci. USA 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the V_H and V_L domains of one fragment are forced to pair with the complementary V_L and V_H domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See, Gruber et al., J. Immunol. 152:5368 (1994).

Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt et al., J. Immunol. 147:60 (1991).

Exemplary bispecific antibodies can bind to two different epitopes, at least one of which originates in the protein antigen of the invention. Alternatively, an anti-antigenic arm of an immunoglobulin molecule can be combined with an arm which binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (e.g. CD2, CD3, CD28, or B7), or Fc receptors for IgG (Fc R), such as Fc RI (CD64), Fc RII (CD32) and Fc RIII (CD16) so as to focus cellular defense mechanisms to the cell expressing the particular antigen. Bispecific

antibodies can also be used to direct cytotoxic agents to cells which express a particular antigen. These antibodies possess an antigen-binding arm and an arm which binds a cytotoxic agent or a radionuclide chelator, such as EOTUBE, DPTA, DOTA, or TETA. Another bispecific antibody of interest binds the protein antigen described herein and further binds
5 tissue factor (TF).

Heteroconjugate Antibodies

Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such
10 antibodies have, for example, been proposed to target immune system cells to unwanted cells (U.S. Patent No. 4,676,980), and for treatment of HIV infection (WO 91/00360; WO 92/200373; EP 03089). It is contemplated that the antibodies can be prepared in vitro using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins can be constructed using a disulfide exchange reaction or by
15 forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate and those disclosed, for example, in U.S. Patent No. 4,676,980.

Effector Function Engineering

It can be desirable to modify the antibody of the invention with respect to effector
20 function, so as to enhance, e.g., the effectiveness of the antibody in treating cancer. For example, cysteine residue(s) can be introduced into the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated can have improved internalization capability and/or increased complement-mediated cell killing and
25 antibody-dependent cellular cytotoxicity (ADCC). See Caron et al., J. Exp Med., 176: 1191-1195 (1992) and Shopes, J. Immunol., 148: 2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity can also be prepared using heterobifunctional cross-linkers as described in Wolff et al. Cancer Research, 53: 2560-2565 (1993). Alternatively, an antibody can be engineered that has dual Fc regions and can thereby have enhanced complement lysis
30 and ADCC capabilities. See Stevenson et al., Anti-Cancer Drug Design, 3: 219-230 (1989).

Immunoconjugates

The invention also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (e.g., an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (i.e., a radioconjugate).

Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, Phytolaca americana proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include ^{212}Bi , ^{131}I , ^{131}In , ^{90}Y , and ^{186}Re .

Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta et al., Science, 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO94/11026.

In another embodiment, the antibody can be conjugated to a "receptor" (such as streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (e.g., avidin) that is in turn conjugated to a cytotoxic agent.

NOVX Recombinant Expression Vectors and Host Cells

Another aspect of the invention pertains to vectors, preferably expression vectors, containing a nucleic acid encoding a NOVX protein, or derivatives, fragments, analogs or homologs thereof. As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (*e.g.*, bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (*e.g.*, non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively-linked. Such vectors are referred to herein as "expression vectors". In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" can be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (*e.g.*, replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

The recombinant expression vectors of the invention comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, that is operatively-linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, "operably-linked" is intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner that allows for expression of the nucleotide sequence (*e.g.*, in an *in vitro* transcription/translation system or in a host cell when the vector is introduced into the host cell).

The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (*e.g.*, polyadenylation signals). Such regulatory sequences are described, for example, in Goeddel, GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY 185, Academic Press, San Diego, Calif. (1990). Regulatory sequences include

those that direct constitutive expression of a nucleotide sequence in many types of host cell and those that direct expression of the nucleotide sequence only in certain host cells (e.g., tissue-specific regulatory sequences). It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. The expression vectors of the invention can be introduced into host cells to thereby produce proteins or peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein (e.g., NOVX proteins, mutant forms of NOVX proteins, fusion proteins, etc.).

The recombinant expression vectors of the invention can be designed for expression of NOVX proteins in prokaryotic or eukaryotic cells. For example, NOVX proteins can be expressed in bacterial cells such as *Escherichia coli*, insect cells (using baculovirus expression vectors) yeast cells or mammalian cells. Suitable host cells are discussed further in Goeddel, GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY 185, Academic Press, San Diego, Calif. (1990). Alternatively, the recombinant expression vector can be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

Expression of proteins in prokaryotes is most often carried out in *Escherichia coli* with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein encoded therein, usually to the amino terminus of the recombinant protein. Such fusion vectors typically serve three purposes: (i) to increase expression of recombinant protein; (ii) to increase the solubility of the recombinant protein; and (iii) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase. Typical fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith and Johnson, 1988. *Gene* 67: 31-40), pMAL (New England Biolabs, Beverly, Mass.) and pRIT5 (Pharmacia, Piscataway, N.J.) that fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein.

Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amrann *et al.*, (1988) *Gene* 69:301-315) and pET 11d (Studier *et al.*, GENE EXPRESSION

TECHNOLOGY: METHODS IN ENZYMOLOGY 185, Academic Press, San Diego, Calif. (1990) 60-89).

One strategy to maximize recombinant protein expression in *E. coli* is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant protein. *See, e.g.*, Gottesman, GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY 185, Academic Press, San Diego, Calif. (1990) 119-128. Another strategy is to alter the nucleic acid sequence of the nucleic acid to be inserted into an expression vector so that the individual codons for each amino acid are those preferentially utilized in *E. coli* (*see, e.g.*, Wada, *et al.*, 1992. *Nucl. Acids Res.* 20: 2111-2118). Such alteration of nucleic acid sequences of the invention can be carried out by standard DNA synthesis techniques.

In another embodiment, the NOVX expression vector is a yeast expression vector. Examples of vectors for expression in yeast *Saccharomyces cerevisiae* include pYepSec1 (Baldari, *et al.*, 1987. *EMBO J.* 6: 229-234), pMFa (Kurjan and Herskowitz, 1982. *Cell* 30: 933-943), pJRY88 (Schultz *et al.*, 1987. *Gene* 54: 113-123), pYES2 (Invitrogen Corporation, San Diego, Calif.), and picZ (InVitrogen Corp, San Diego, Calif.).

Alternatively, NOVX can be expressed in insect cells using baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (*e.g.*, SF9 cells) include the pAc series (Smith, *et al.*, 1983. *Mol. Cell. Biol.* 3: 2156-2165) and the pVL series (Lucklow and Summers, 1989. *Virology* 170: 31-39).

In yet another embodiment, a nucleic acid of the invention is expressed in mammalian cells using a mammalian expression vector. Examples of mammalian expression vectors include pCDM8 (Seed, 1987. *Nature* 329: 840) and pMT2PC (Kaufman, *et al.*, 1987. *EMBO J.* 6: 187-195). When used in mammalian cells, the expression vector's control functions are often provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma, adenovirus 2, cytomegalovirus, and simian virus 40. For other suitable

promoters include the albumin promoter (liver-specific; Pinkert, *et al.*, 1987. *Genes Dev.* 1: 268-277), lymphoid-specific promoters (Calame and Eaton, 1988. *Adv. Immunol.* 43: 235-275), in particular promoters of T cell receptors (Winoto and Baltimore, 1989. *EMBO J.* 8: 729-733) and immunoglobulins (Banerji, *et al.*, 1983. *Cell* 33: 729-740; Queen and
5 Baltimore, 1983. *Cell* 33: 741-748), neuron-specific promoters (*e.g.*, the neurofilament promoter; Byrne and Ruddle, 1989. *Proc. Natl. Acad. Sci. USA* 86: 5473-5477), pancreas-specific promoters (Edlund, *et al.*, 1985. *Science* 230: 912-916), and mammary gland-specific promoters (*e.g.*, milk whey promoter; U.S. Pat. No. 4,873,316 and European Application Publication No. 264,166). Developmentally-regulated promoters are also
10 encompassed, *e.g.*, the murine hox promoters (Kessel and Gruss, 1990. *Science* 249: 374-379) and the α -fetoprotein promoter (Campes and Tilghman, 1989. *Genes Dev.* 3: 537-546).

The invention further provides a recombinant expression vector comprising a DNA molecule of the invention cloned into the expression vector in an antisense orientation. That is, the DNA molecule is operatively-linked to a regulatory sequence in a manner that allows
15 for expression (by transcription of the DNA molecule) of an RNA molecule that is antisense to NOVX mRNA. Regulatory sequences operatively linked to a nucleic acid cloned in the antisense orientation can be chosen that direct the continuous expression of the antisense RNA molecule in a variety of cell types, for instance viral promoters and/or enhancers, or regulatory sequences can be chosen that direct constitutive, tissue specific or cell type specific expression
20 of antisense RNA. The antisense expression vector can be in the form of a recombinant plasmid, phagemid or attenuated virus in which antisense nucleic acids are produced under the control of a high efficiency regulatory region, the activity of which can be determined by the cell type into which the vector is introduced. For a discussion of the regulation of gene expression using antisense genes *see, e.g.*, Weintraub, *et al.*, "Antisense RNA as a molecular
25 tool for genetic analysis," *Reviews-Trends in Genetics*, Vol. 1(1) 1986.

Another aspect of the invention pertains to host cells into which a recombinant expression vector of the invention has been introduced. The terms "host cell" and "recombinant host cell" are used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but also to the progeny or potential progeny of such a
30 cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

A host cell can be any prokaryotic or eukaryotic cell. For example, NOVX protein can be expressed in bacterial cells such as *E. coli*, insect cells, yeast or mammalian cells (such as human, Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to those skilled in the art.

- 5 Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (*e.g.*, DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation.
- 10 Suitable methods for transforming or transfecting host cells can be found in Sambrook, *et al.* (MOLECULAR CLONING: A LABORATORY MANUAL, 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989), and other laboratory manuals.

- 15 For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (*e.g.*, resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Various selectable markers include those that confer resistance to drugs, such as G418, hygromycin and methotrexate. Nucleic acid encoding a
- 20 selectable marker can be introduced into a host cell on the same vector as that encoding NOVX or can be introduced on a separate vector. Cells stably transfected with the introduced nucleic acid can be identified by drug selection (*e.g.*, cells that have incorporated the selectable marker gene will survive, while the other cells die).

- 25 A host cell of the invention, such as a prokaryotic or eukaryotic host cell in culture, can be used to produce (*i.e.*, express) NOVX protein. Accordingly, the invention further provides methods for producing NOVX protein using the host cells of the invention. In one embodiment, the method comprises culturing the host cell of invention (into which a recombinant expression vector encoding NOVX protein has been introduced) in a suitable medium such that NOVX protein is produced. In another embodiment, the method further
- 30 comprises isolating NOVX protein from the medium or the host cell.

Transgenic NOVX Animals

The host cells of the invention can also be used to produce non-human transgenic animals. For example, in one embodiment, a host cell of the invention is a fertilized oocyte or an embryonic stem cell into which NOVX protein-coding sequences have been introduced. Such host cells can then be used to create non-human transgenic animals in which exogenous NOVX sequences have been introduced into their genome or homologous recombinant animals in which endogenous NOVX sequences have been altered. Such animals are useful for studying the function and/or activity of NOVX protein and for identifying and/or evaluating modulators of NOVX protein activity. As used herein, a "transgenic animal" is a non-human animal, preferably a mammal, more preferably a rodent such as a rat or mouse, in which one or more of the cells of the animal includes a transgene. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, amphibians, etc. A transgene is exogenous DNA that is integrated into the genome of a cell from which a transgenic animal develops and that remains in the genome of the mature animal, thereby directing the expression of an encoded gene product in one or more cell types or tissues of the transgenic animal. As used herein, a "homologous recombinant animal" is a non-human animal, preferably a mammal, more preferably a mouse, in which an endogenous NOVX gene has been altered by homologous recombination between the endogenous gene and an exogenous DNA molecule introduced into a cell of the animal, *e.g.*, an embryonic cell of the animal, prior to development of the animal.

A transgenic animal of the invention can be created by introducing NOVX-encoding nucleic acid into the male pronuclei of a fertilized oocyte (*e.g.*, by microinjection, retroviral infection) and allowing the oocyte to develop in a pseudopregnant female foster animal. Sequences including SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25 can be introduced as a transgene into the genome of a non-human animal. Alternatively, a non-human homologue of the human NOVX gene, such as a mouse NOVX gene, can be isolated based on hybridization to the human NOVX cDNA (described further *supra*) and used as a transgene. Intronic sequences and polyadenylation signals can also be included in the transgene to increase the efficiency of expression of the transgene. A tissue-specific regulatory sequence(s) can be operably-linked to the NOVX transgene to direct expression of NOVX protein to particular cells. Methods for generating transgenic animals via embryo

manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866; 4,870,009; and 4,873,191; and Hogan, 1986. In: MANIPULATING THE MOUSE EMBRYO, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. Similar methods are used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of the NOVX transgene in its genome and/or expression of NOVX mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying a transgene-encoding NOVX protein can further be bred to other transgenic animals carrying other transgenes.

10 To create a homologous recombinant animal, a vector is prepared which contains at least a portion of a NOVX gene into which a deletion, addition or substitution has been introduced to thereby alter, *e.g.*, functionally disrupt, the NOVX gene. The NOVX gene can be a human gene (*e.g.*, the DNA of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25), but more preferably, is a non-human homologue of a human NOVX gene. For example, a mouse homologue of human NOVX gene of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25 can be used to construct a homologous recombination vector suitable for altering an endogenous NOVX gene in the mouse genome. In one embodiment, the vector is designed such that, upon homologous recombination, the endogenous NOVX gene is functionally disrupted (*i.e.*, no longer encodes a functional protein; also referred to as a "knock out" vector).

20 Alternatively, the vector can be designed such that, upon homologous recombination, the endogenous NOVX gene is mutated or otherwise altered but still encodes functional protein (*e.g.*, the upstream regulatory region can be altered to thereby alter the expression of the endogenous NOVX protein). In the homologous recombination vector, the altered portion of the NOVX gene is flanked at its 5'- and 3'-termini by additional nucleic acid of the NOVX gene to allow for homologous recombination to occur between the exogenous NOVX gene carried by the vector and an endogenous NOVX gene in an embryonic stem cell. The additional flanking NOVX nucleic acid is of sufficient length for successful homologous recombination with the endogenous gene. Typically, several kilobases of flanking DNA (both at the 5'- and 3'-termini) are included in the vector. *See, e.g.*, Thomas, *et al.*, 1987. *Cell* 51: 503 for a description of homologous recombination vectors. The vector is then introduced into an embryonic stem cell line (*e.g.*, by electroporation) and cells in which the introduced NOVX

gene has homologously-recombined with the endogenous NOVX gene are selected. *See, e.g., Li, et al., 1992. Cell* 69: 915.

The selected cells are then injected into a blastocyst of an animal (*e.g., a mouse*) to form aggregation chimeras. *See, e.g., Bradley, 1987. In: TERATOCARCINOMAS AND*
5 *EMBRYONIC STEM CELLS: A PRACTICAL APPROACH*, Robertson, ed. IRL, Oxford, pp. 113-152. A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term. Progeny harboring the homologously-recombined DNA in their germ cells can be used to breed animals in which all cells of the animal contain the homologously-recombined DNA by germline transmission of the transgene. Methods for
10 constructing homologous recombination vectors and homologous recombinant animals are described further in Bradley, 1991. *Curr. Opin. Biotechnol.* 2: 823-829; PCT International Publication Nos.: WO 90/11354; WO 91/01140; WO 92/0968; and WO 93/04169.

In another embodiment, transgenic non-humans animals can be produced that contain selected systems that allow for regulated expression of the transgene. One example of such a
15 system is the cre/loxP recombinase system of bacteriophage P1. For a description of the cre/loxP recombinase system, *See, e.g., Lakso, et al., 1992. Proc. Natl. Acad. Sci. USA* 89: 6232-6236. Another example of a recombinase system is the FLP recombinase system of *Saccharomyces cerevisiae*. *See, O'Gorman, et al., 1991. Science* 251:1351-1355. If a cre/loxP recombinase system is used to regulate expression of the transgene, animals containing
20 transgenes encoding both the Cre recombinase and a selected protein are required. Such animals can be provided through the construction of "double" transgenic animals, *e.g., by* mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase.

Clones of the non-human transgenic animals described herein can also be produced
25 according to the methods described in Wilmut, *et al., 1997. Nature* 385: 810-813. In brief, a cell (*e.g., a somatic cell*) from the transgenic animal can be isolated and induced to exit the growth cycle and enter G₀ phase. The quiescent cell can then be fused, *e.g., through the use of* electrical pulses, to an enucleated oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops to
30 morula or blastocyte and then transferred to pseudopregnant female foster animal. The offspring borne of this female foster animal will be a clone of the animal from which the cell (*e.g., the somatic cell*) is isolated.

Pharmaceutical Compositions

The NOVX nucleic acid molecules, NOVX proteins, and anti-NOVX antibodies (also referred to herein as "active compounds") of the invention, and derivatives, fragments, analogs and homologs thereof, can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier. As used herein, "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. Suitable carriers are described in the most recent edition of Remington's Pharmaceutical Sciences, a standard reference text in the field, which is incorporated herein by reference. Preferred examples of such carriers or diluents include, but are not limited to, water, saline, finger's solutions, dextrose solution, and 5% human serum albumin. Liposomes and non-aqueous vehicles such as fixed oils may also be used. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

The antibodies disclosed herein can also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein et al., Proc. Natl. Acad. Sci. USA, 82: 3688 (1985); Hwang et al., Proc. Natl. Acad. Sci. USA, 77: 4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Patent No. 5,013,556.

Particularly useful liposomes can be generated by the reverse-phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol, and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin et al., J. Biol. Chem., 257: 286-288 (1982) via a disulfide-interchange reaction. A chemotherapeutic agent (such as Doxorubicin) is optionally contained within the liposome. See Gabizon et al., J. National Cancer Inst., 81(19): 1484 (1989).

A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, *e.g.*, intravenous, intradermal, subcutaneous, oral (*e.g.*, inhalation), transdermal (*i.e.*, topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, 5 intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid (EDTA); buffers such as acetates, citrates or phosphates, 10 and agents for the adjustment of tonicity such as sodium chloride or dextrose. The pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous 15 solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringeability exists. It must be stable under 20 the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by 25 the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the 30 composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound (*e.g.*, a NOVX protein or anti-NOVX antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, methods of preparation are vacuum drying and freeze-drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant, *e.g.*, a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

The compounds can also be prepared in the form of suppositories (*e.g.*, with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

In one embodiment, the active compounds are prepared with carriers that will protect
5 the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be
10 obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

15 It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The
20 specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

The nucleic acid molecules of the invention can be inserted into vectors and used as
25 gene therapy vectors. Gene therapy vectors can be delivered to a subject by, for example, intravenous injection, local administration (*see, e.g.*, U.S. Patent No. 5,328,470) or by stereotactic injection (*see, e.g.*, Chen, *et al.*, 1994. *Proc. Natl. Acad. Sci. USA* 91: 3054-3057). The pharmaceutical preparation of the gene therapy vector can include the gene therapy vector in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery
30 vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from recombinant cells, *e.g.*, retroviral vectors, the pharmaceutical preparation can include one or more cells that produce the gene delivery system.

Antibodies specifically binding a protein of the invention, as well as other molecules identified by the screening assays disclosed herein, can be administered for the treatment of various disorders in the form of pharmaceutical compositions. Principles and considerations involved in preparing such compositions, as well as guidance in the choice of components are provided, for example, in Remington : The Science And Practice Of Pharmacy 19th ed. (Alfonso R. Gennaro, et al., editors) Mack Pub. Co., Easton, Pa. : 1995; Drug Absorption Enhancement : Concepts, Possibilities, Limitations, And Trends, Harwood Academic Publishers, Langhorne, Pa., 1994; and Peptide And Protein Drug Delivery (Advances In Parenteral Sciences, Vol. 4), 1991, M. Dekker, New York. If the antigenic protein is intracellular and whole antibodies are used as inhibitors, internalizing antibodies are preferred. However, liposomes can also be used to deliver the antibody, or an antibody fragment, into cells. Where antibody fragments are used, the smallest inhibitory fragment that specifically binds to the binding domain of the target protein is preferred. For example, based upon the variable-region sequences of an antibody, peptide molecules can be designed that retain the ability to bind the target protein sequence. Such peptides can be synthesized chemically and/or produced by recombinant DNA technology. See, *e.g.*, Marasco *et al.*, 1993 *Proc. Natl. Acad. Sci. USA*, 90: 7889-7893. The formulation herein can also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Alternatively, or in addition, the composition can comprise an agent that enhances its function, such as, for example, a cytotoxic agent, cytokine, chemotherapeutic agent, or growth-inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended. The active ingredients can also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles, and nanocapsules) or in macroemulsions.

The formulations to be used for *iv vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

Sustained-release preparations can be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the

antibody, which matrices are in the form of shaped articles, e.g., films, or microcapsules.

Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, 5 degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods.

- 10 The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

Screening and Detection Methods

- The isolated nucleic acid molecules of the invention can be used to express NOVX 15 protein (e.g., via a recombinant expression vector in a host cell in gene therapy applications), to detect NOVX mRNA (e.g., in a biological sample) or a genetic lesion in a NOVX gene, and to modulate NOVX activity, as described further, below. In addition, the NOVX proteins can be used to screen drugs or compounds that modulate the NOVX protein activity or expression as well as to treat disorders characterized by insufficient or excessive production of NOVX 20 protein or production of NOVX protein forms that have decreased or aberrant activity compared to NOVX wild-type protein. In addition, the anti-NOVX antibodies of the invention can be used to detect and isolate NOVX proteins and modulate NOVX activity. For example, NOVX activity includes growth and differentiation, antibody production, and tumor growth.

- The invention further pertains to novel agents identified by the screening assays 25 described herein and uses thereof for treatments as described, *supra*.

Screening Assays

- The invention provides a method (also referred to herein as a "screening assay") for identifying modulators, *i.e.*, candidate or test compounds or agents (*e.g.*, peptides, 30 peptidomimetics, small molecules or other drugs) that bind to NOVX proteins or have a stimulatory or inhibitory effect on, *e.g.*, NOVX protein expression or NOVX protein activity. The invention also includes compounds identified in the screening assays described herein.

In one embodiment, the invention provides assays for screening candidate or test compounds which bind to or modulate the activity of the membrane-bound form of a NOVX protein or polypeptide or biologically-active portion thereof. The test compounds of the invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the "one-bead one-compound" library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds. See, e.g., Lam, 1997. *Anticancer Drug Design* 12: 145.

A "small molecule" as used herein, is meant to refer to a composition that has a molecular weight of less than about 5 kD and most preferably less than about 4 kD. Small molecules can be, e.g., nucleic acids, peptides, polypeptides, peptidomimetics, carbohydrates, lipids or other organic or inorganic molecules. Libraries of chemical and/or biological mixtures, such as fungal, bacterial, or algal extracts, are known in the art and can be screened with any of the assays of the invention.

Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt, *et al.*, 1993. *Proc. Natl. Acad. Sci. U.S.A.* 90: 6909; Erb, *et al.*, 1994. *Proc. Natl. Acad. Sci. U.S.A.* 91: 11422; Zuckermann, *et al.*, 1994. *J. Med. Chem.* 37: 2678; Cho, *et al.*, 1993. *Science* 261: 1303; Carrell, *et al.*, 1994. *Angew. Chem. Int. Ed. Engl.* 33: 2059; Carrell, *et al.*, 1994. *Angew. Chem. Int. Ed. Engl.* 33: 2061; and Gallop, *et al.*, 1994. *J. Med. Chem.* 37: 1233.

Libraries of compounds may be presented in solution (e.g., Houghten, 1992. *Biotechniques* 13: 412-421), or on beads (Lam, 1991. *Nature* 354: 82-84), on chips (Fodor, 1993. *Nature* 364: 555-556), bacteria (Ladner, U.S. Patent No. 5,223,409), spores (Ladner, U.S. Patent 5,233,409), plasmids (Cull, *et al.*, 1992. *Proc. Natl. Acad. Sci. USA* 89: 1865-1869) or on phage (Scott and Smith, 1990. *Science* 249: 386-390; Devlin, 1990. *Science* 249: 404-406; Cwirla, *et al.*, 1990. *Proc. Natl. Acad. Sci. U.S.A.* 87: 6378-6382; Felici, 1991. *J. Mol. Biol.* 222: 301-310; Ladner, U.S. Patent No. 5,233,409.).

In one embodiment, an assay is a cell-based assay in which a cell which expresses a membrane-bound form of NOVX protein, or a biologically-active portion thereof, on the cell surface is contacted with a test compound and the ability of the test compound to bind to a

NOVX protein determined. The cell, for example, can be of mammalian origin or a yeast cell.

Determining the ability of the test compound to bind to the NOVX protein can be accomplished, for example, by coupling the test compound with a radioisotope or enzymatic label such that binding of the test compound to the NOVX protein or biologically-active

- 5 portion thereof can be determined by detecting the labeled compound in a complex. For example, test compounds can be labeled with ^{125}I , ^{35}S , ^{14}C , or ^3H , either directly or indirectly, and the radioisotope detected by direct counting of radioemission or by scintillation counting. Alternatively, test compounds can be enzymatically-labeled with, for example, horseradish peroxidase, alkaline phosphatase, or luciferase, and the enzymatic label detected by
- 10 determination of conversion of an appropriate substrate to product. In one embodiment, the assay comprises contacting a cell which expresses a membrane-bound form of NOVX protein, or a biologically-active portion thereof, on the cell surface with a known compound which binds NOVX to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with a NOVX protein, wherein
- 15 determining the ability of the test compound to interact with a NOVX protein comprises determining the ability of the test compound to preferentially bind to NOVX protein or a biologically-active portion thereof as compared to the known compound.

- In another embodiment, an assay is a cell-based assay comprising contacting a cell expressing a membrane-bound form of NOVX protein, or a biologically-active portion thereof,
- 20 on the cell surface with a test compound and determining the ability of the test compound to modulate (*e.g.*, stimulate or inhibit) the activity of the NOVX protein or biologically-active portion thereof. Determining the ability of the test compound to modulate the activity of NOVX or a biologically-active portion thereof can be accomplished, for example, by determining the ability of the NOVX protein to bind to or interact with a NOVX target
- 25 molecule. As used herein, a "target molecule" is a molecule with which a NOVX protein binds or interacts in nature, for example, a molecule on the surface of a cell which expresses a NOVX interacting protein, a molecule on the surface of a second cell, a molecule in the extracellular milieu, a molecule associated with the internal surface of a cell membrane or a cytoplasmic molecule. A NOVX target molecule can be a non-NOVX molecule or a NOVX
- 30 protein or polypeptide of the invention. In one embodiment, a NOVX target molecule is a component of a signal transduction pathway that facilitates transduction of an extracellular signal (*e.g.* a signal generated by binding of a compound to a membrane-bound NOVX

molecule) through the cell membrane and into the cell. The target, for example, can be a second intercellular protein that has catalytic activity or a protein that facilitates the association of downstream signaling molecules with NOVX.

Determining the ability of the NOVX protein to bind to or interact with a NOVX target molecule can be accomplished by one of the methods described above for determining direct binding. In one embodiment, determining the ability of the NOVX protein to bind to or interact with a NOVX target molecule can be accomplished by determining the activity of the target molecule. For example, the activity of the target molecule can be determined by detecting induction of a cellular second messenger of the target (*i.e.* intracellular Ca^{2+} , diacylglycerol, IP_3 , etc.), detecting catalytic/enzymatic activity of the target an appropriate substrate, detecting the induction of a reporter gene (comprising a NOVX-responsive regulatory element operatively linked to a nucleic acid encoding a detectable marker, *e.g.*, luciferase), or detecting a cellular response, for example, cell survival, cellular differentiation, or cell proliferation.

In yet another embodiment, an assay of the invention is a cell-free assay comprising contacting a NOVX protein or biologically-active portion thereof with a test compound and determining the ability of the test compound to bind to the NOVX protein or biologically-active portion thereof. Binding of the test compound to the NOVX protein can be determined either directly or indirectly as described above. In one such embodiment, the assay comprises contacting the NOVX protein or biologically-active portion thereof with a known compound which binds NOVX to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with a NOVX protein, wherein determining the ability of the test compound to interact with a NOVX protein comprises determining the ability of the test compound to preferentially bind to NOVX or biologically-active portion thereof as compared to the known compound.

In still another embodiment, an assay is a cell-free assay comprising contacting NOVX protein or biologically-active portion thereof with a test compound and determining the ability of the test compound to modulate (*e.g.* stimulate or inhibit) the activity of the NOVX protein or biologically-active portion thereof. Determining the ability of the test compound to modulate the activity of NOVX can be accomplished, for example, by determining the ability of the NOVX protein to bind to a NOVX target molecule by one of the methods described above for determining direct binding. In an alternative embodiment, determining the ability of

the test compound to modulate the activity of NOVX protein can be accomplished by determining the ability of the NOVX protein further modulate a NOVX target molecule. For example, the catalytic/enzymatic activity of the target molecule on an appropriate substrate can be determined as described above.

5 In yet another embodiment, the cell-free assay comprises contacting the NOVX protein or biologically-active portion thereof with a known compound which binds NOVX protein to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with a NOVX protein, wherein determining the ability of the test compound to interact with a NOVX protein comprises determining the
10 ability of the NOVX protein to preferentially bind to or modulate the activity of a NOVX target molecule.

The cell-free assays of the invention are amenable to use of both the soluble form or the membrane-bound form of NOVX protein. In the case of cell-free assays comprising the membrane-bound form of NOVX protein, it may be desirable to utilize a solubilizing agent
15 such that the membrane-bound form of NOVX protein is maintained in solution. Examples of such solubilizing agents include non-ionic detergents such as n-octylglucoside, n-dodecylglucoside, n-dodecylmaltoside, octanoyl-N-methylglucamide, decanoyl-N-methylglucamide, Triton® X-100, Triton® X-114, Thesit®, Isotridecypoly(ethylene glycol ether)_n, N-dodecyl--N,N-dimethyl-3-ammonio-1-propane sulfonate,
20 3-(3-cholamidopropyl) dimethylamminiol-1-propane sulfonate (CHAPS), or 3-(3-cholamidopropyl)dimethylamminiol-2-hydroxy-1-propane sulfonate (CHAPSO).

In more than one embodiment of the above assay methods of the invention, it may be desirable to immobilize either NOVX protein or its target molecule to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate
25 automation of the assay. Binding of a test compound to NOVX protein, or interaction of NOVX protein with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and micro-centrifuge tubes. In one embodiment, a fusion protein can be provided that adds a domain that allows one or both of the proteins to be
30 bound to a matrix. For example, GST-NOVX fusion proteins or GST-target fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtiter plates, that are then combined with the test compound or the

test compound and either the non-adsorbed target protein or NOVX protein, and the mixture is incubated under conditions conducive to complex formation (*e.g.*, at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components, the matrix immobilized in the case of beads, complex
5 determined either directly or indirectly, for example, as described, *supra*. Alternatively, the complexes can be dissociated from the matrix, and the level of NOVX protein binding or activity determined using standard techniques.

Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either the NOVX protein or its target
10 molecule can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated NOVX protein or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well-known within the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, Ill.), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies reactive with NOVX protein or target molecules, but
15 which do not interfere with binding of the NOVX protein to its target molecule, can be derivatized to the wells of the plate, and unbound target or NOVX protein trapped in the wells by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the NOVX protein or target molecule, as well as enzyme-linked
20 assays that rely on detecting an enzymatic activity associated with the NOVX protein or target molecule.

In another embodiment, modulators of NOVX protein expression are identified in a method wherein a cell is contacted with a candidate compound and the expression of NOVX mRNA or protein in the cell is determined. The level of expression of NOVX mRNA or
25 protein in the presence of the candidate compound is compared to the level of expression of NOVX mRNA or protein in the absence of the candidate compound. The candidate compound can then be identified as a modulator of NOVX mRNA or protein expression based upon this comparison. For example, when expression of NOVX mRNA or protein is greater (*i.e.*, statistically significantly greater) in the presence of the candidate compound than in its
30 absence, the candidate compound is identified as a stimulator of NOVX mRNA or protein expression. Alternatively, when expression of NOVX mRNA or protein is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate

compound is identified as an inhibitor of NOVX mRNA or protein expression. The level of NOVX mRNA or protein expression in the cells can be determined by methods described herein for detecting NOVX mRNA or protein.

In yet another aspect of the invention, the NOVX proteins can be used as "bait proteins" in a two-hybrid assay or three hybrid assay (*see, e.g.*, U.S. Patent No. 5,283,317; Zervos, *et al.*, 1993. *Cell* 72: 223-232; Madura, *et al.*, 1993. *J. Biol. Chem.* 268: 12046-12054; Bartel, *et al.*, 1993. *Biotechniques* 14: 920-924; Iwabuchi, *et al.*, 1993. *Oncogene* 8: 1693-1696; and Brent WO 94/10300), to identify other proteins that bind to or interact with NOVX ("NOVX-binding proteins" or "NOVX-bp") and modulate NOVX activity. Such NOVX-binding proteins are also likely to be involved in the propagation of signals by the NOVX proteins as, for example, upstream or downstream elements of the NOVX pathway.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. In one construct, the gene that codes for NOVX is fused to a gene encoding the DNA binding domain of a known transcription factor (*e.g.*, GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact, *in vivo*, forming a NOVX-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (*e.g.*, LacZ) that is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene that encodes the protein which interacts with NOVX.

The invention further pertains to novel agents identified by the aforementioned screening assays and uses thereof for treatments as described herein.

Detection Assays

Portions or fragments of the cDNA sequences identified herein (and the corresponding complete gene sequences) can be used in numerous ways as polynucleotide reagents. By way of example, and not of limitation, these sequences can be used to: (i) identify an individual

from a minute biological sample (tissue typing); and (ii) aid in forensic identification of a biological sample. Some of these applications are described in the subsections, below.

Tissue Typing

5 The NOVX sequences of the invention can be used to identify individuals from minute biological samples. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identification. The sequences of the invention are useful as additional DNA markers for RFLP ("restriction fragment length polymorphisms," described in U.S. Patent No. 5,272,057).

10 Furthermore, the sequences of the invention can be used to provide an alternative technique that determines the actual base-by-base DNA sequence of selected portions of an individual's genome. Thus, the NOVX sequences described herein can be used to prepare two PCR primers from the 5'- and 3'-termini of the sequences. These primers can then be used to amplify an individual's DNA and subsequently sequence it.

15 Panels of corresponding DNA sequences from individuals, prepared in this manner, can provide unique individual identifications, as each individual will have a unique set of such DNA sequences due to allelic differences. The sequences of the invention can be used to obtain such identification sequences from individuals and from tissue. The NOVX sequences of the invention uniquely represent portions of the human genome. Allelic variation occurs to
20 some degree in the coding regions of these sequences, and to a greater degree in the noncoding regions. It is estimated that allelic variation between individual humans occurs with a frequency of about once per each 500 bases. Much of the allelic variation is due to single nucleotide polymorphisms (SNPs), which include restriction fragment length polymorphisms (RFLPs).

25 Each of the sequences described herein can, to some degree, be used as a standard against which DNA from an individual can be compared for identification purposes. Because greater numbers of polymorphisms occur in the noncoding regions, fewer sequences are necessary to differentiate individuals. The noncoding sequences can comfortably provide positive individual identification with a panel of perhaps 10 to 1,000 primers that each yield a
30 noncoding amplified sequence of 100 bases. If predicted coding sequences, such as those in SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25 are used, a more appropriate number of primers for positive individual identification would be 500-2,000.

Predictive Medicine

The invention also pertains to the field of predictive medicine in which diagnostic assays, prognostic assays, pharmacogenomics, and monitoring clinical trials are used for prognostic (predictive) purposes to thereby treat an individual prophylactically. Accordingly, one aspect of the invention relates to diagnostic assays for determining NOVX protein and/or nucleic acid expression as well as NOVX activity, in the context of a biological sample (*e.g.*, blood, serum, cells, tissue) to thereby determine whether an individual is afflicted with a disease or disorder, or is at risk of developing a disorder, associated with aberrant NOVX expression or activity. Disorders associated with aberrant NOVX expression of activity include, for example, disorders of olfactory loss, *e.g.* trauma, HIV illness, neoplastic growth, and neurological disorders, *e.g.* Parkinson's disease and Alzheimer's disease.

The invention also provides for prognostic (or predictive) assays for determining whether an individual is at risk of developing a disorder associated with NOVX protein, nucleic acid expression or activity. For example, mutations in a NOVX gene can be assayed in a biological sample. Such assays can be used for prognostic or predictive purpose to thereby prophylactically treat an individual prior to the onset of a disorder characterized by or associated with NOVX protein, nucleic acid expression, or biological activity.

Another aspect of the invention provides methods for determining NOVX protein, nucleic acid expression or activity in an individual to thereby select appropriate therapeutic or prophylactic agents for that individual (referred to herein as "pharmacogenomics"). Pharmacogenomics allows for the selection of agents (*e.g.*, drugs) for therapeutic or prophylactic treatment of an individual based on the genotype of the individual (*e.g.*, the genotype of the individual examined to determine the ability of the individual to respond to a particular agent.)

Yet another aspect of the invention pertains to monitoring the influence of agents (*e.g.*, drugs, compounds) on the expression or activity of NOVX in clinical trials.

These and other agents are described in further detail in the following sections.

Diagnostic Assays

An exemplary method for detecting the presence or absence of NOVX in a biological sample involves obtaining a biological sample from a test subject and contacting the biological

sample with a compound or an agent capable of detecting NOVX protein or nucleic acid (*e.g.*, mRNA, genomic DNA) that encodes NOVX protein such that the presence of NOVX is detected in the biological sample. An agent for detecting NOVX mRNA or genomic DNA is a labeled nucleic acid probe capable of hybridizing to NOVX mRNA or genomic DNA. The nucleic acid probe can be, for example, a full-length NOVX nucleic acid, such as the nucleic acid of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25, or a portion thereof, such as an oligonucleotide of at least 15, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically hybridize under stringent conditions to NOVX mRNA or genomic DNA. Other suitable probes for use in the diagnostic assays of the invention are described herein.

One agent for detecting NOVX protein is an antibody capable of binding to NOVX protein, preferably an antibody with a detectable label. Antibodies directed against a protein of the invention may be used in methods known within the art relating to the localization and/or quantitation of the protein (*e.g.*, for use in measuring levels of the protein within appropriate physiological samples, for use in diagnostic methods, for use in imaging the protein, and the like). In a given embodiment, antibodies against the proteins, or derivatives, fragments, analogs or homologs thereof, that contain the antigen binding domain, are utilized as pharmacologically-active compounds.

An antibody specific for a protein of the invention can be used to isolate the protein by standard techniques, such as immunoaffinity chromatography or immunoprecipitation. Such an antibody can facilitate the purification of the natural protein antigen from cells and of recombinantly produced antigen expressed in host cells. Moreover, such an antibody can be used to detect the antigenic protein (*e.g.*, in a cellular lysate or cell supernatant) in order to evaluate the abundance and pattern of expression of the antigenic protein. Antibodies directed against the protein can be used diagnostically to monitor protein levels in tissue as part of a clinical testing procedure, *e.g.*, to, for example, determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling (*i.e.*, physically linking) the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine,

dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

5 Antibodies can be polyclonal, or more preferably, monoclonal. An intact antibody, or a fragment thereof (*e.g.*, Fab or F(ab')_2) can be used. The term "labeled", with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (*i.e.*, physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly
10 labeled. Examples of indirect labeling include detection of a primary antibody using a fluorescently-labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently-labeled streptavidin. The term "biological sample" is intended to include tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject. That is, the detection method of the invention
15 can be used to detect NOVX mRNA, protein, or genomic DNA in a biological sample *in vitro* as well as *in vivo*. For example, *in vitro* techniques for detection of NOVX mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro* techniques for detection of NOVX protein include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations, and immunofluorescence. *In vitro* techniques for detection of NOVX
20 genomic DNA include Southern hybridizations. Furthermore, *in vivo* techniques for detection of NOVX protein include introducing into a subject a labeled anti-NOVX antibody. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques.

 In one embodiment, the biological sample contains protein molecules from the test
25 subject. Alternatively, the biological sample can contain mRNA molecules from the test subject or genomic DNA molecules from the test subject. A preferred biological sample is a peripheral blood leukocyte sample isolated by conventional means from a subject.

 In one embodiment, the methods further involve obtaining a control biological sample from a control subject, contacting the control sample with a compound or agent capable of
30 detecting NOVX protein, mRNA, or genomic DNA, such that the presence of NOVX protein, mRNA or genomic DNA is detected in the biological sample, and comparing the presence of

NOVX protein, mRNA or genomic DNA in the control sample with the presence of NOVX protein, mRNA or genomic DNA in the test sample.

The invention also encompasses kits for detecting the presence of NOVX in a biological sample. For example, the kit can comprise: a labeled compound or agent capable of
5 detecting NOVX protein or mRNA in a biological sample; means for determining the amount of NOVX in the sample; and means for comparing the amount of NOVX in the sample with a standard. The compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect NOVX protein or nucleic acid.

10 Prognostic Assays

The diagnostic methods described herein can furthermore be utilized to identify subjects having or at risk of developing a disease or disorder associated with aberrant NOVX expression or activity. For example, the assays described herein, such as the preceding diagnostic assays or the following assays, can be utilized to identify a subject having or at risk
15 of developing a disorder associated with NOVX protein, nucleic acid expression or activity. Such disorders include for example, disorders of olfactory loss, *e.g.* trauma, HIV illness, neoplastic growth, and neurological disorders, *e.g.* Parkinson's disease and Alzheimer's disease.

Alternatively, the prognostic assays can be utilized to identify a subject having or at
20 risk for developing a disease or disorder. Thus, the invention provides a method for identifying a disease or disorder associated with aberrant NOVX expression or activity in which a test sample is obtained from a subject and NOVX protein or nucleic acid (*e.g.*, mRNA, genomic DNA) is detected, wherein the presence of NOVX protein or nucleic acid is diagnostic for a subject having or at risk of developing a disease or disorder associated with
25 aberrant NOVX expression or activity. As used herein, a "test sample" refers to a biological sample obtained from a subject of interest. For example, a test sample can be a biological fluid (*e.g.*, serum), cell sample, or tissue.

Furthermore, the prognostic assays described herein can be used to determine whether a subject can be administered an agent (*e.g.*, an agonist, antagonist, peptidomimetic, protein,
30 peptide, nucleic acid, small molecule, or other drug candidate) to treat a disease or disorder associated with aberrant NOVX expression or activity. For example, such methods can be used to determine whether a subject can be effectively treated with an agent for a disorder.

Thus, the invention provides methods for determining whether a subject can be effectively treated with an agent for a disorder associated with aberrant NOVX expression or activity in which a test sample is obtained and NOVX protein or nucleic acid is detected (*e.g.*, wherein the presence of NOVX protein or nucleic acid is diagnostic for a subject that can be administered the agent to treat a disorder associated with aberrant NOVX expression or activity).

The methods of the invention can also be used to detect genetic lesions in a NOVX gene, thereby determining if a subject with the lesioned gene is at risk for a disorder characterized by aberrant cell proliferation and/or differentiation. In various embodiments, the methods include detecting, in a sample of cells from the subject, the presence or absence of a genetic lesion characterized by at least one of an alteration affecting the integrity of a gene encoding a NOVX-protein, or the misexpression of the NOVX gene. For example, such genetic lesions can be detected by ascertaining the existence of at least one of: (i) a deletion of one or more nucleotides from a NOVX gene; (ii) an addition of one or more nucleotides to a NOVX gene; (iii) a substitution of one or more nucleotides of a NOVX gene, (iv) a chromosomal rearrangement of a NOVX gene; (v) an alteration in the level of a messenger RNA transcript of a NOVX gene, (vi) aberrant modification of a NOVX gene, such as of the methylation pattern of the genomic DNA, (vii) the presence of a non-wild-type splicing pattern of a messenger RNA transcript of a NOVX gene, (viii) a non-wild-type level of a NOVX protein, (ix) allelic loss of a NOVX gene, and (x) inappropriate post-translational modification of a NOVX protein. As described herein, there are a large number of assay techniques known in the art which can be used for detecting lesions in a NOVX gene. A preferred biological sample is a peripheral blood leukocyte sample isolated by conventional means from a subject. However, any biological sample containing nucleated cells may be used, including, for example, buccal mucosal cells.

In certain embodiments, detection of the lesion involves the use of a probe/primer in a polymerase chain reaction (PCR) (*see, e.g.*, U.S. Patent Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (*see, e.g.*, Landegran, *et al.*, 1988. *Science* 241: 1077-1080; and Nakazawa, *et al.*, 1994. *Proc. Natl. Acad. Sci. USA* 91: 360-364), the latter of which can be particularly useful for detecting point mutations in the NOVX-gene (*see, Abravaya, et al.*, 1995. *Nucl. Acids Res.* 23: 675-682). This method can include the steps of collecting a sample of cells from a patient, isolating nucleic

acid (e.g., genomic, mRNA or both) from the cells of the sample, contacting the nucleic acid sample with one or more primers that specifically hybridize to a NOVX gene under conditions such that hybridization and amplification of the NOVX gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. It is anticipated that PCR and/or LCR
5 may be desirable to use as a preliminary amplification step in conjunction with any of the techniques used for detecting mutations described herein.

Alternative amplification methods include: self sustained sequence replication (*see*, Guatelli, *et al.*, 1990. *Proc. Natl. Acad. Sci. USA* 87: 1874-1878), transcriptional amplification
10 system (*see*, Kwoh, *et al.*, 1989. *Proc. Natl. Acad. Sci. USA* 86: 1173-1177); Q β Replicase (*see*, Lizardi, *et al.*, 1988. *BioTechnology* 6: 1197), or any other nucleic acid amplification method, followed by the detection of the amplified molecules using techniques well known to those of skill in the art. These detection schemes are especially useful for the detection of nucleic acid molecules if such molecules are present in very low numbers.

15 In an alternative embodiment, mutations in a NOVX gene from a sample cell can be identified by alterations in restriction enzyme cleavage patterns. For example, sample and control DNA is isolated, amplified (optionally), digested with one or more restriction endonucleases, and fragment length sizes are determined by gel electrophoresis and compared. Differences in fragment length sizes between sample and control DNA indicates mutations in
20 the sample DNA. Moreover, the use of sequence specific ribozymes (*see, e.g.*, U.S. Patent No. 5,493,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site.

In other embodiments, genetic mutations in NOVX can be identified by hybridizing a sample and control nucleic acids, e.g., DNA or RNA, to high-density arrays containing
25 hundreds or thousands of oligonucleotides probes. *See, e.g.*, Cronin, *et al.*, 1996. *Human Mutation* 7: 244-255; Kozal, *et al.*, 1996. *Nat. Med.* 2: 753-759. For example, genetic mutations in NOVX can be identified in two dimensional arrays containing light-generated DNA probes as described in Cronin, *et al., supra*. Briefly, a first hybridization array of probes can be used to scan through long stretches of DNA in a sample and control to identify base
30 changes between the sequences by making linear arrays of sequential overlapping probes. This step allows the identification of point mutations. This is followed by a second hybridization array that allows the characterization of specific mutations by using smaller,

specialized probe arrays complementary to all variants or mutations detected. Each mutation array is composed of parallel probe sets, one complementary to the wild-type gene and the other complementary to the mutant gene.

In yet another embodiment, any of a variety of sequencing reactions known in the art can be used to directly sequence the NOVX gene and detect mutations by comparing the
5 sequence of the sample NOVX with the corresponding wild-type (control) sequence. Examples of sequencing reactions include those based on techniques developed by Maxim and Gilbert, 1977. *Proc. Natl. Acad. Sci. USA* 74: 560 or Sanger, 1977. *Proc. Natl. Acad. Sci. USA* 74: 5463. It is also contemplated that any of a variety of automated sequencing procedures can
10 be utilized when performing the diagnostic assays (*see, e.g.,* Naeve, *et al.*, 1995. *Biotechniques* 19: 448), including sequencing by mass spectrometry (*see, e.g.,* PCT International Publication No. WO 94/16101; Cohen, *et al.*, 1996. *Adv. Chromatography* 36: 127-162; and Griffin, *et al.*, 1993. *Appl. Biochem. Biotechnol.* 38: 147-159).

Other methods for detecting mutations in the NOVX gene include methods in which
15 protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA heteroduplexes. *See, e.g.,* Myers, *et al.*, 1985. *Science* 230: 1242. In general, the art technique of "mismatch cleavage" starts by providing heteroduplexes of formed by hybridizing (labeled) RNA or DNA containing the wild-type NOVX sequence with potentially mutant RNA or DNA obtained from a tissue sample. The double-stranded duplexes are treated
20 with an agent that cleaves single-stranded regions of the duplex such as which will exist due to basepair mismatches between the control and sample strands. For instance, RNA/DNA duplexes can be treated with RNase and DNA/DNA hybrids treated with S₁ nuclease to enzymatically digesting the mismatched regions. In other embodiments, either DNA/DNA or RNA/DNA duplexes can be treated with hydroxylamine or osmium tetroxide and with
25 piperidine in order to digest mismatched regions. After digestion of the mismatched regions, the resulting material is then separated by size on denaturing polyacrylamide gels to determine the site of mutation. *See, e.g.,* Cotton, *et al.*, 1988. *Proc. Natl. Acad. Sci. USA* 85: 4397; Saleeba, *et al.*, 1992. *Methods Enzymol.* 217: 286-295. In an embodiment, the control DNA or RNA can be labeled for detection.

30 In still another embodiment, the mismatch cleavage reaction employs one or more proteins that recognize mismatched base pairs in double-stranded DNA (so called "DNA mismatch repair" enzymes) in defined systems for detecting and mapping point mutations in

NOVX cDNAs obtained from samples of cells. For example, the mutY enzyme of *E. coli* cleaves A at G/A mismatches and the thymidine DNA glycosylase from HeLa cells cleaves T at G/T mismatches. *See, e.g., Hsu, et al., 1994. Carcinogenesis 15: 1657-1662.* According to an exemplary embodiment, a probe based on a NOVX sequence, *e.g., a wild-type NOVX*
5 *sequence, is hybridized to a cDNA or other DNA product from a test cell(s). The duplex is treated with a DNA mismatch repair enzyme, and the cleavage products, if any, can be detected from electrophoresis protocols or the like. See, e.g., U.S. Patent No. 5,459,039.*

In other embodiments, alterations in electrophoretic mobility will be used to identify mutations in NOVX genes. For example, single strand conformation polymorphism (SSCP)
10 *may be used to detect differences in electrophoretic mobility between mutant and wild type nucleic acids. See, e.g., Orita, et al., 1989. Proc. Natl. Acad. Sci. USA: 86: 2766; Cotton, 1993. Mutat. Res. 285: 125-144; Hayashi, 1992. Genet. Anal. Tech. Appl. 9: 73-79.* Single-stranded DNA fragments of sample and control NOVX nucleic acids will be denatured and allowed to renature. The secondary structure of single-stranded nucleic acids varies
15 *according to sequence, the resulting alteration in electrophoretic mobility enables the detection of even a single base change. The DNA fragments may be labeled or detected with labeled probes. The sensitivity of the assay may be enhanced by using RNA (rather than DNA), in which the secondary structure is more sensitive to a change in sequence. In one embodiment, the subject method utilizes heteroduplex analysis to separate double stranded heteroduplex*
20 *molecules on the basis of changes in electrophoretic mobility. See, e.g., Keen, et al., 1991. Trends Genet. 7: 5.*

In yet another embodiment, the movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient gel electrophoresis (DGGE). *See, e.g., Myers, et al., 1985. Nature 313: 495.* When DGGE is
25 *used as the method of analysis, DNA will be modified to insure that it does not completely denature, for example by adding a GC clamp of approximately 40 bp of high-melting GC-rich DNA by PCR. In a further embodiment, a temperature gradient is used in place of a denaturing gradient to identify differences in the mobility of control and sample DNA. See, e.g., Rosenbaum and Reissner, 1987. Biophys. Chem. 265: 12753.*

30 *Examples of other techniques for detecting point mutations include, but are not limited to, selective oligonucleotide hybridization, selective amplification, or selective primer extension. For example, oligonucleotide primers may be prepared in which the known*

mutation is placed centrally and then hybridized to target DNA under conditions that permit hybridization only if a perfect match is found. *See, e.g., Saiki, et al., 1986. Nature 324: 163; Saiki, et al., 1989. Proc. Natl. Acad. Sci. USA 86: 6230.* Such allele specific oligonucleotides are hybridized to PCR amplified target DNA or a number of different mutations when the
5 oligonucleotides are attached to the hybridizing membrane and hybridized with labeled target DNA.

Alternatively, allele specific amplification technology that depends on selective PCR amplification may be used in conjunction with the instant invention. Oligonucleotides used as primers for specific amplification may carry the mutation of interest in the center of the
10 molecule (so that amplification depends on differential hybridization; *see, e.g., Gibbs, et al., 1989. Nucl. Acids Res. 17: 2437-2448*) or at the extreme 3'-terminus of one primer where, under appropriate conditions, mismatch can prevent, or reduce polymerase extension (*see, e.g., Prossner, 1993. Tibtech. 11: 238*). In addition it may be desirable to introduce a novel restriction site in the region of the mutation to create cleavage-based detection. *See, e.g.,*
15 Gasparini, *et al., 1992. Mol. Cell Probes 6: 1.* It is anticipated that in certain embodiments amplification may also be performed using *Taq* ligase for amplification. *See, e.g., Barany, 1991. Proc. Natl. Acad. Sci. USA 88: 189.* In such cases, ligation will occur only if there is a perfect match at the 3'-terminus of the 5' sequence, making it possible to detect the presence of a known mutation at a specific site by looking for the presence or absence of amplification.

20 The methods described herein may be performed, for example, by utilizing pre-packaged diagnostic kits comprising at least one probe nucleic acid or antibody reagent described herein, which may be conveniently used, *e.g.,* in clinical settings to diagnose patients exhibiting symptoms or family history of a disease or illness involving a NOVX gene.

Furthermore, any cell type or tissue, preferably peripheral blood leukocytes, in which
25 NOVX is expressed may be utilized in the prognostic assays described herein. However, any biological sample containing nucleated cells may be used, including, for example, buccal mucosal cells.

Pharmacogenomics

Agents, or modulators that have a stimulatory or inhibitory effect on NOVX activity
30 (*e.g., NOVX gene expression*), as identified by a screening assay described herein can be administered to individuals to treat (prophylactically or therapeutically) disorders (*e.g.* proliferative disorders, *e.g.* cancer and mastocytosis, immune disorders, hepatic disorders, *e.g.*

cirrhosis, viral infections, *e.g.* AIDS and hepatitis, and disorders of the neuro-olfactory system *e.g.* trauma, surgery and/or neoplastic disorders). In conjunction with such treatment, the pharmacogenomics (*i.e.*, the study of the relationship between an individual's genotype and that individual's response to a foreign compound or drug) of the individual may be considered.

- 5 Differences in metabolism of therapeutics can lead to severe toxicity or therapeutic failure by altering the relation between dose and blood concentration of the pharmacologically active drug. Thus, the pharmacogenomics of the individual permits the selection of effective agents (*e.g.*, drugs) for prophylactic or therapeutic treatments based on a consideration of the individual's genotype. Such pharmacogenomics can further be used to determine appropriate
- 10 dosages and therapeutic regimens. Accordingly, the activity of NOVX protein, expression of NOVX nucleic acid, or mutation content of NOVX genes in an individual can be determined to thereby select appropriate agent(s) for therapeutic or prophylactic treatment of the individual.

- Pharmacogenomics deals with clinically significant hereditary variations in the
- 15 response to drugs due to altered drug disposition and abnormal action in affected persons. See *e.g.*, Eichelbaum, 1996. *Clin. Exp. Pharmacol. Physiol.*, 23: 983-985; Linder, 1997. *Clin. Chem.*, 43: 254-266. In general, two types of pharmacogenetic conditions can be differentiated. Genetic conditions transmitted as a single factor altering the way drugs act on the body (altered drug action) or genetic conditions transmitted as single factors altering the
- 20 way the body acts on drugs (altered drug metabolism). These pharmacogenetic conditions can occur either as rare defects or as polymorphisms. For example, glucose-6-phosphate dehydrogenase (G6PD) deficiency is a common inherited enzymopathy in which the main clinical complication is hemolysis after ingestion of oxidant drugs (anti-malarials, sulfonamides, analgesics, nitrofurans) and consumption of fava beans.

- 25 As an illustrative embodiment, the activity of drug metabolizing enzymes is a major determinant of both the intensity and duration of drug action. The discovery of genetic polymorphisms of drug metabolizing enzymes (*e.g.*, N-acetyltransferase 2 (NAT 2) and cytochrome P450 enzymes CYP2D6 and CYP2C19) has provided an explanation as to why some patients do not obtain the expected drug effects or show exaggerated drug response and
- 30 serious toxicity after taking the standard and safe dose of a drug. These polymorphisms are expressed in two phenotypes in the population, the extensive metabolizer (EM) and poor metabolizer (PM). The prevalence of PM is different among different populations. For

example, the gene coding for CYP2D6 is highly polymorphic and several mutations have been identified in PM, which all lead to the absence of functional CYP2D6. Poor metabolizers of CYP2D6 and CYP2C19 quite frequently experience exaggerated drug response and side effects when they receive standard doses. If a metabolite is the active therapeutic moiety, PM show no therapeutic response, as demonstrated for the analgesic effect of codeine mediated by its CYP2D6-formed metabolite morphine. At the other extreme are the so called ultra-rapid metabolizers who do not respond to standard doses. Recently, the molecular basis of ultra-rapid metabolism has been identified to be due to CYP2D6 gene amplification.

Thus, the activity of NOVX protein, expression of NOVX nucleic acid, or mutation content of NOVX genes in an individual can be determined to thereby select appropriate agent(s) for therapeutic or prophylactic treatment of the individual. In addition, pharmacogenetic studies can be used to apply genotyping of polymorphic alleles encoding drug-metabolizing enzymes to the identification of an individual's drug responsiveness phenotype. This knowledge, when applied to dosing or drug selection, can avoid adverse reactions or therapeutic failure and thus enhance therapeutic or prophylactic efficiency when treating a subject with a NOVX modulator, such as a modulator identified by one of the exemplary screening assays described herein.

Monitoring of Effects During Clinical Trials

Monitoring the influence of agents (*e.g.*, drugs, compounds) on the expression or activity of NOVX (*e.g.*, the ability to modulate aberrant cell proliferation) can be applied not only in basic drug screening, but also in clinical trials. For example, the effectiveness of an agent determined by a screening assay as described herein to increase NOVX gene expression, protein levels, or upregulate NOVX activity, can be monitored in clinical trials of subjects exhibiting decreased NOVX gene expression, protein levels, or downregulated NOVX activity. Alternatively, the effectiveness of an agent determined by a screening assay to decrease NOVX gene expression, protein levels, or downregulate NOVX activity, can be monitored in clinical trials of subjects exhibiting increased NOVX gene expression, protein levels, or upregulated NOVX activity. In such clinical trials, the expression or activity of NOVX and, preferably, other genes that have been implicated in, for example, a cellular proliferation or immune disorder can be used as a "read out" or markers of the immune responsiveness of a particular cell.

By way of example, and not of limitation, genes, including NOVX, that are modulated in cells by treatment with an agent (*e.g.*, compound, drug or small molecule) that modulates NOVX activity (*e.g.*, identified in a screening assay as described herein) can be identified.

Thus, to study the effect of agents on cellular proliferation disorders, for example, in a clinical trial, cells can be isolated and RNA prepared and analyzed for the levels of expression of NOVX and other genes implicated in the disorder. The levels of gene expression (*i.e.*, a gene expression pattern) can be quantified by Northern blot analysis or RT-PCR, as described herein, or alternatively by measuring the amount of protein produced, by one of the methods as described herein, or by measuring the levels of activity of NOVX or other genes. In this manner, the gene expression pattern can serve as a marker, indicative of the physiological response of the cells to the agent. Accordingly, this response state may be determined before, and at various points during, treatment of the individual with the agent.

In one embodiment, the invention provides a method for monitoring the effectiveness of treatment of a subject with an agent (*e.g.*, an agonist, antagonist, protein, peptide, peptidomimetic, nucleic acid, small molecule, or other drug candidate identified by the screening assays described herein) comprising the steps of (i) obtaining a pre-administration sample from a subject prior to administration of the agent; (ii) detecting the level of expression of a NOVX protein, mRNA, or genomic DNA in the preadministration sample; (iii) obtaining one or more post-administration samples from the subject; (iv) detecting the level of expression or activity of the NOVX protein, mRNA, or genomic DNA in the post-administration samples; (v) comparing the level of expression or activity of the NOVX protein, mRNA, or genomic DNA in the pre-administration sample with the NOVX protein, mRNA, or genomic DNA in the post administration sample or samples; and (vi) altering the administration of the agent to the subject accordingly. For example, increased administration of the agent may be desirable to increase the expression or activity of NOVX to higher levels than detected, *i.e.*, to increase the effectiveness of the agent. Alternatively, decreased administration of the agent may be desirable to decrease expression or activity of NOVX to lower levels than detected, *i.e.*, to decrease the effectiveness of the agent.

Methods of Treatment

The invention provides for both prophylactic and therapeutic methods of treating a subject at risk of (or susceptible to) a disorder or having a disorder associated with aberrant NOVX expression or activity. Disorders associated with aberrant NOVX expression include,

for example, proliferative disorders, *e.g.* cancer and mastocytosis, immune disorders, hepatic disorders, *e.g.* cirrhosis, viral infections, *e.g.* AIDS and hepatitis, and disorders of the neuro-olfactory system *e.g.* surgery and/or neoplastic disorders of olfactory loss, *e.g.* trauma, HIV illness, neoplastic growth, and neurological disorders, *e.g.* Parkinson's disease and

5 Alzheimer's disease.

These methods of treatment will be discussed more fully, below.

Disease and Disorders

Diseases and disorders that are characterized by increased (relative to a subject not
10 suffering from the disease or disorder) levels or biological activity may be treated with
Therapeutics that antagonize (*i.e.*, reduce or inhibit) activity. Therapeutics that antagonize
activity may be administered in a therapeutic or prophylactic manner. Therapeutics that may
be utilized include, but are not limited to: (i) an aforementioned peptide, or analogs,
derivatives, fragments or homologs thereof; (ii) antibodies to an aforementioned peptide; (iii)
15 nucleic acids encoding an aforementioned peptide; (iv) administration of antisense nucleic acid
and nucleic acids that are "dysfunctional" (*i.e.*, due to a heterologous insertion within the
coding sequences of coding sequences to an aforementioned peptide) that are utilized to
"knockout" endogenous function of an aforementioned peptide by homologous recombination
(*see, e.g.*, Capecchi, 1989. *Science* 244: 1288-1292); or (v) modulators (*i.e.*, inhibitors,
20 agonists and antagonists, including additional peptide mimetic of the invention or antibodies
specific to a peptide of the invention) that alter the interaction between an aforementioned
peptide and its binding partner.

Diseases and disorders that are characterized by decreased (relative to a subject not
suffering from the disease or disorder) levels or biological activity may be treated with
25 Therapeutics that increase (*i.e.*, are agonists to) activity. Therapeutics that upregulate activity
may be administered in a therapeutic or prophylactic manner. Therapeutics that may be
utilized include, but are not limited to, an aforementioned peptide, or analogs, derivatives,
fragments or homologs thereof; or an agonist that increases bioavailability.

Increased or decreased levels can be readily detected by quantifying peptide and/or
30 RNA, by obtaining a patient tissue sample (*e.g.*, from biopsy tissue) and assaying it *in vitro* for
RNA or peptide levels, structure and/or activity of the expressed peptides (or mRNAs of an
aforementioned peptide). Methods that are well-known within the art include, but are not

limited to, immunoassays (*e.g.*, by Western blot analysis, immunoprecipitation followed by sodium dodecyl sulfate (SDS) polyacrylamide gel electrophoresis, immunocytochemistry, etc.) and/or hybridization assays to detect expression of mRNAs (*e.g.*, Northern assays, dot blots, *in situ* hybridization, and the like).

5

Prophylactic Methods

In one aspect, the invention provides a method for preventing, in a subject, a disease or condition associated with an aberrant NOVX expression or activity, by administering to the subject an agent that modulates NOVX expression or at least one NOVX activity. Subjects at risk for a disease that is caused or contributed to by aberrant NOVX expression or activity can be identified by, for example, any or a combination of diagnostic or prognostic assays as described herein. Administration of a prophylactic agent can occur prior to the manifestation of symptoms characteristic of the NOVX aberrancy, such that a disease or disorder is prevented or, alternatively, delayed in its progression. Depending upon the type of NOVX aberrancy, for example, a NOVX agonist or NOVX antagonist agent can be used for treating the subject. The appropriate agent can be determined based on screening assays described herein. The prophylactic methods of the invention are further discussed in the following subsections.

20 Therapeutic Methods

Another aspect of the invention pertains to methods of modulating NOVX expression or activity for therapeutic purposes. The modulatory method of the invention involves contacting a cell with an agent that modulates one or more of the activities of NOVX protein activity associated with the cell. An agent that modulates NOVX protein activity can be an agent as described herein, such as a nucleic acid or a protein, a naturally-occurring cognate ligand of a NOVX protein, a peptide, a NOVX peptidomimetic, or other small molecule. In one embodiment, the agent stimulates one or more NOVX protein activity. Examples of such stimulatory agents include active NOVX protein and a nucleic acid molecule encoding NOVX that has been introduced into the cell. In another embodiment, the agent inhibits one or more NOVX protein activity. Examples of such inhibitory agents include antisense NOVX nucleic acid molecules and anti-NOVX antibodies. These modulatory methods can be performed *in vitro* (*e.g.*, by culturing the cell with the agent) or, alternatively, *in vivo* (*e.g.*, by administering

the agent to a subject). As such, the invention provides methods of treating an individual afflicted with a disease or disorder characterized by aberrant expression or activity of a NOVX protein or nucleic acid molecule. In one embodiment, the method involves administering an agent (*e.g.*, an agent identified by a screening assay described herein), or combination of
5 agents that modulates (*e.g.*, up-regulates or down-regulates) NOVX expression or activity. In another embodiment, the method involves administering a NOVX protein or nucleic acid molecule as therapy to compensate for reduced or aberrant NOVX expression or activity.

Stimulation of NOVX activity is desirable in situations in which NOVX is abnormally downregulated and/or in which increased NOVX activity is likely to have a beneficial effect.

10 One example of such a situation is where a subject has a disorder characterized by aberrant cell proliferation and/or differentiation (*e.g.*, cancer or immune associated). Another example of such a situation is where the subject has an immunodeficiency disease (*e.g.*, AIDS).

Antibodies of the invention, including polyclonal, monoclonal, humanized and fully human antibodies, may be used as therapeutic agents. Such agents will generally be employed to
15 treat or prevent a disease or pathology in a subject. An antibody preparation, preferably one having high specificity and high affinity for its target antigen, is administered to the subject and will generally have an effect due to its binding with the target. Such an effect may be one of two kinds, depending on the specific nature of the interaction between the given antibody molecule and the target antigen in question. In the first instance, administration of the
20 antibody may abrogate or inhibit the binding of the target with an endogenous ligand to which it naturally binds. In this case, the antibody binds to the target and masks a binding site of the naturally occurring ligand, wherein the ligand serves as an effector molecule. Thus the receptor mediates a signal transduction pathway for which ligand is responsible.

Alternatively, the effect may be one in which the antibody elicits a physiological result
25 by virtue of binding to an effector binding site on the target molecule. In this case the target, a receptor having an endogenous ligand which may be absent or defective in the disease or pathology, binds the antibody as a surrogate effector ligand, initiating a receptor-based signal transduction event by the receptor.

A therapeutically effective amount of an antibody of the invention relates generally to
30 the amount needed to achieve a therapeutic objective. As noted above, this may be a binding interaction between the antibody and its target antigen that, in certain cases, interferes with the functioning of the target, and in other cases, promotes a physiological response. The amount

required to be administered will furthermore depend on the binding affinity of the antibody for its specific antigen, and will also depend on the rate at which an administered antibody is depleted from the free volume other subject to which it is administered. Common ranges for therapeutically effective dosing of an antibody or antibody fragment of the invention may be, by way of nonlimiting example, from about 0.1 mg/kg body weight to about 50 mg/kg body weight. Common dosing frequencies may range, for example, from twice daily to once a week.

Determination of the Biological Effect of the Therapeutic

In various embodiments of the invention, suitable *in vitro* or *in vivo* assays are performed to determine the effect of a specific Therapeutic and whether its administration is indicated for treatment of the affected tissue.

In various specific embodiments, *in vitro* assays may be performed with representative cells of the type(s) involved in the patient's disorder, to determine if a given Therapeutic exerts the desired effect upon the cell type(s). Compounds for use in therapy may be tested in suitable animal model systems including, but not limited to rats, mice, chicken, cows, monkeys, rabbits, and the like, prior to testing in human subjects. Similarly, for *in vivo* testing, any of the animal model system known in the art may be used prior to administration to human subjects.

The invention will be further described in the following examples, which do not limit the scope of the invention described in the claims.

EXAMPLES

Example 1.: Method of Identifying the Nucleic Acids of the Present Invention.

Novel nucleic acid sequences were identified by TblastN using CuraGen Corporation's sequence file run against the Genomic Daily Files made available by GenBank. The nucleic acids were further predicted by the program GenScan™, including selection of exons. These were further modified by means of similarities using BLAST searches. The sequences were then manually corrected for apparent inconsistencies, thereby obtaining the sequences encoding the full-length protein.

Example 2.: Preparation of the mammalian expression vector pCEP4/Sec

The oligonucleotide primers,
pSec-V5-His Forward
CTCGTCCTCGAGGGTAAGCCTATCCCTAAC (SEQ ID NO: 105) and
pSec-V5-His Reverse
5 CTCGTCGGGCCCCTGATCAGCGGGTTTAAAC (SEQ ID NO: 106),

were designed to amplify a fragment from the pcDNA3.1-V5His (Invitrogen, Carlsbad, CA) expression vector that includes V5 and His6. The PCR product was digested with XhoI and ApaI and ligated into the XhoI/ApaI digested pSecTag2 B vector harboring an Ig kappa leader sequence
10 (Invitrogen, Carlsbad CA). The correct structure of the resulting vector, pSecV5His, including an in-frame Ig-kappa leader and V5-His6 was verified by DNA sequence analysis. The vector pSecV5His was digested with PmeI and NheI to provide a fragment retaining the above elements in the correct frame. The PmeI-NheI fragment was ligated into the BamHI/Klenow and NheI treated vector pCEP4 (Invitrogen, Carlsbad, CA). The resulting vector was named pCEP4/Sec and includes an in-frame Ig
15 kappa leader, a site for insertion of a clone of interest, V5 and His6 under control of the PCMV and/or the PT7 promoter. pCEP4/Sec is an expression vector that allows heterologous protein expression and secretion by fusing any protein to the Ig Kappa chain signal peptide. Detection and purification of the expressed protein are aided by the presence of the V5 epitope tag and 6xHis tag at the C-terminus (Invitrogen, Carlsbad, CA).

20

Example 3. Molecular cloning of the full length clone 83350421_EXT (NOV1)

Oligonucleotide primers were designed to PCR amplify a DNA segment representing an ORF coding for the full length 83350421_EXT. The forward primer includes an in frame BamHI restriction site and a consensus Kozak sequence. The reverse primer contains an in
25 frame Xho restriction site. The sequences of the primers are the following:

83350421 Forw: GGATCCACCATGAGTGAGCTTGTAAGAGCAAGATCC (SEQ ID NO:107), and

83350421 Rev: CTCGAGTGGTTGCGCATCACCTGCTTCCAGCAC (SEQ ID NO:108).

30 PCR reactions were set up using 5 ng cDNA template consisting of equal portions of human testis, fetal brain, mammary, skeletal muscle derived cDNA, 1 microM of each of 10219646 MatF and 10219646 Reverse primers, 5 micromoles dNTP (Clontech Laboratories,

Palo Alto CA) and 1 microliter of 50xAdvantage-HF 2 polymerase (Clontech Laboratories) in 50 microliter volume. The following reaction conditions were used:

- a) 96°C 3 minutes
- b) 96°C 30 seconds denaturation
- 5 c) 70°C 30 seconds, primer annealing. This temperature was gradually decreased by 1°C/cycle
- d) 72°C 1 minute extension.
- Repeat steps b-d 10 times
- 10 e) 96°C 30 seconds denaturation
- f) 60°C 30 seconds annealing
- g) 72°C 1 minute extension
- Repeat steps e-g 25 times
- h) 72°C 5 minutes final extension

15 An amplified product having a size of approximately 300 bp was detected by agarose gel electrophoresis. The construct is called pCR2.1-83350421-S747-3A. The product was isolated and ligated into the pCR2.1 vector (Invitrogen Corp, Carlsbad, CA). The DNA sequence of the cloned insert differs from that of clone 83350421_EXT_REVCOMP reported in Table 2. The sequence of the cloned insert in pCR2.1-83350421-S747-3A (SEQ ID NO: 20 109) is shown in Table 51.

Table 51. The nucleotide sequence of the insert in pCR2.1-83350421-S747-3A.

CTCGAGTGGTTGCGCATCACCTGCTTCCAGCACTTTAGTGAGATCAAAAGTGGGC
ATAATACCCTCCCTGACATCAGGACCATCTCCAGGCTCATCCTCTATCTTAAGCAG
AGCCAGTTCCTGTTGAAAAGCTTCCATGTCAGGCCCTTGAAAAGCAGGCACTGCTT
25 GATTTTCAATCTCCCCACTAGGTGCAATACCCTGATTATCAGTTGGTGGTTCTCTT
CTTGACGTTTTTCTCAGTGGGCTCCTGGACAATCACAGATCCAACCGGCTGGGAA
GACTCTTGGTCATTTCTTCTGAGGATTGGGATCTTGCTCTTACAAGCTCACTC
ATGGTGGATCC (SEQ ID NO:109)

30 It was determined this sequence codes for a 111 amino acid residue polypeptide as shown in Table 52. The cloned sequence has an insertion, compared to the sequence of the protein of clone 83350421_EXT_REVCOMP reported in Table 2. The insertion introduces 17 extra residues into the polypeptide, shown in Table 52 (SEQ ID NO:110) (indicated by underlined bold font).

35 **Table 52.** The amino acid sequence of the polypeptide coded by the insert of pCR2.1-83350421-S747-3A.

MSELVRARSQSSERGNQESSQPVGSVIVQEPTEEKRQEEEPPTDNQGIAPSGEIQAVPAFQGPDM
AFQQLALLKIEDEPGDGPVREGIMPTFDLTKVLEAGDAQP (SEQ ID NO:110)

Example 4.: Expression of NOV 1 (83350421) in human embryonic kidney 293 cells.

The BamHI-XhoI fragment containing the 83350421 sequence was isolated from clone pCR2.1-83350421-S747-3A (Example 3) and subcloned into BamHI-XhoI digested pCEP4/Sec (Example 2) to generate expression vector pCEP4/Sec-83350421. The pCEP4/Sec-83350421 vector was transfected into human embryonic kidney 293 cells (ATCC No. CRL-1573, Manassas, VA) using the LipofectaminePlus reagent following the manufacturer's instructions (Gibco/BRL/Life Technologies, Rockville, MD). The cell pellet and supernatant, respectively, were harvested 72 hours after transfection and examined for 83350421 expression by Western blotting (reducing conditions) with an anti-V5 antibody. Fig. 1 shows that 83350421 is expressed as a polypeptide with an apparent molecular weight of about 30 kDa protein in the 293 cell pellet, based on use SeeBlue Molecular Weight Standards (Invitrogen). However, secretion of the protein by 293 cells was not observed.

15

20




Figure 1. Expression of NOV1 (83350421) protein in 293 cells.

Example 5. Molecular cloning of the full length clone NOV5 (ba403c19_A)

Oligonucleotide primers were designed to amplify a DNA segment representing the full length NOV5 (ba403c19_A). The forward primer includes an in frame BamHI restriction site and the consensus Kozak sequence, and the reverse primer contains an in frame XhoI restriction site. The sequences of the PCR primers are the following: Ba403c19_A Alt-FL-Forward: GGATCCACCATGATTCAAAAGTGTGGCTTGAGATCC (SEQ ID NO:111), and Ba403c19_A Reverse: CTCGAGTTTCCTCCTGAATAGAGCTGTAAATTG (SEQ ID NO:112).

PCR reactions were set up using a total of 5ng mixture of cDNA template containing equal amounts of cDNAs derived from human fetal brain, human testis, human mammary and human skeletal muscle tissues, 1 microM of each of the Ba403c19_A Alt-FL-Forward, and Ba403c19_A Reverse primers, 5 micromoles dNTP (Clontech Laboratories, Palo Alto CA) and 1 microliter of 50xAdvantage-HF 2 polymerase (Clontech Laboratories) in 50 microliter volume. The following reaction conditions were used:

- a) 96°C 3 minutes
- b) 96°C 30 seconds denaturation
- c) 70°C 30 seconds, primer annealing. This temperature was gradually decreased by 1°C/cycle
- d) 72°C 1 minute extension.
- Repeat steps b-d 10 times
- e) 96°C 30 seconds denaturation
- f) 60°C 30 seconds annealing
- g) 72°C 1 minute extension
- Repeat steps e-g 25 times
- h) 72°C 10 minutes final extension

PCR products having the expected size of approximately 600 bp were isolated from agarose gel and ligated to pCR2.1 vector (Invitrogen, Carlsbad, CA). The clone is called pCR2.1-cg Ba403c19-S551-6e. The cloned insert was sequenced, using vector specific, M13 Forward(-40) and M13 Reverse primers as well as the following gene specific primers:

Ba403c19_A S1: GGACTTGATCAGCAAGCAGAG (SEQ ID NO:113) and
Ba403c19_A S2: CTCTGCTTGCTGATCAAGTCC (SEQ ID NO:114).

The cloned sequence (SEQ ID NO:115) differs by one nucleotide (shown in underlined bold font in Table 53) from that presented for clone Ba403c19_A in Table 16 (SEQ ID NO:9).

Table 53. Nucleotide sequence of the clone pCR2.1-cg Ba403c19-S551-6e.

ATGATTCAAAAGTGTGTTGGCTTGAGATCCTTATGGGTATATTCATTGCTGGCACCCTATCCCTGGAC
TGTAACCTTACTGAACGTTACCTGAGAAGAGTCACCTGGCAAATCTGAGACATCTGAGTAGTATGAGC
AATTCATTTCTGTAGAATGTCTACGAGAAAACATAGCTTTTGTAGTTGCCCAAGAGTTTCTGCAATAC
ACCCAACCTATGAAGAGGGACATCAAGAAGGCCTTCTATGAAATGTCCCTACAGGCCTTCAACATCTTC

AGCCAACACACCTTCAAATATTGGAAAGAGAGACACCTCAAACAAATCCAAATAGGACTTGATCAGCAA
 GCAGAGTACCTGAACCAATGCTTGGAGGAAGAGCGGAGAATGAAAATGAAGACATGAAAGAAATGAAAGAG
 AATGAGATGAAACCCCTCAGAAGCCAGGGTCCCCCAGCTGAGCAGCCTGGAAGTGAAGAGATATTTCCAC
 AGGATAGACAATTTCTGAAAGAAAAGAAATACAGTGACTGTGCCTGGGAGATTGTCCGAGTGGAAATC
 5 AGAAGATGTTTGTATTACTTTTACAAATTTACAGCTCTATTTCAGGAGGAAA (SEQ ID
 NO:115)

This base change causes a corresponding change of one amino acid in the polypeptide coded
 by the insert of pCR2.1-cg Ba403c19-S551-6e (SEQ ID NO:116) (shown in underlined bold
 10 font in Table 54) from the sequence for the NOV5 (ba403c19_A) polypeptide shown in Table
 16 (SEQ ID NO: 10).

**Table 54. Amino acid sequence of the polypeptide coded by the insert of
 pCR2.1-cg Ba403c19-S551-6e.**

15 MIQKCLWLEILMGIFLAGTSLDCNLLNVHLRRVTWQNLRLHSSMSNSFPVECLRENI
 AFELPQEFLQYTQPMKRDIKKAIFYEMSLQAFNIFSQHTFKYWKERHLKQIQIGLDQQA
 EYLNQCLEEDENENEDMKEMKENEMKPSEARVPQLSSLELRRYFHRIDNFLKEKKYS
 DCAWEIVRVEIRRCLYYFYKFTALFRRK (SEQ ID NO:116)

20 **Example 6. Molecular cloning of a mature form of NOV5 (ba403c19_A)**

Oligonucleotide primers were designed to PCR amplify a DNA
 segment representing the cDNA coding for a mature form of the
 25 ba403c19_A sequence. The forward primer includes an in frame BamHI
 restriction site, and the reverse primer contains an in frame XhoI
 restriction site. The sequences of the PCR primers are the following:

Ba403c19_A Mat-Forward: GGATCCCTGGACTGTAACCTACTGAACGTTCCACC (SEQ ID
 30 NO:117) and

BA403c19_A REVERSE: CTCGAGTTTCCTCCTGAATAGAGCTGTAAATTTG (SEQ ID NO:118).

PCR reactions were set up using a total of 5ng mixture of cDNA template containing equal amounts of cDNAs derived from human fetal brain, human testis, human mammary and human skeletal muscle tissues, 1 microM of each of the Ba403c19_A Mat-Forward, and Ba403c19_A Reverse primers, 5 micromoles dNTP (Clontech Laboratories, Palo Alto CA) and 1 microliter of 50xAdvantage-HF 2 polymerase (Clontech Laboratories) in 50 microliter volume. The following reaction conditions were used:

- a) 96°C 3 minutes
- b) 96°C 30 seconds denaturation
- c) 70°C 30 seconds, primer annealing. This temperature was gradually decreased by 1°C/cycle
- d) 72°C 1minute extension.
Repeat steps b-d 10 times
- e) 96°C 30 seconds denaturation
- f) 60°C 30 seconds annealing
- g) 72°C 1 minute extension
Repeat steps e-g 25 times
- h) 72°C 10 minutes final extension

PCR products having the expected size of approximately 600 bp were isolated from agarose gel and ligated to pCR2.1 vector (Invitrogen, Carlsbad, CA). The clone is called pCR2.1-cg Ba403c19-S546-1b. The cloned insert was sequenced using vector specific, M13 Forward(-40) and M13 Reverse primers as well as the gene specific primers Ba403c19_A S1 (SEQ ID NO:113) and Ba403c19_A S2 (SEQ ID NO:114) used in Example 5.

The cloned sequence was verified as an ORF coding for a polypeptide representing a mature form of clone Ba403c19 (see Table 55; SEQ ID NO:119). The sequence differs by the same single nucleotide as found in Example 5 (shown in underlined bold font in Table 53) from that presented for clone Ba403c19_A in Table 16 (SEQ ID NO: 9).

Table 55. Nucleotide sequence of the clone pCR2.1-cg Ba403c19-S546-1b.

CTGGACTGTAACCTACTGAACGTTACCTGAGAAGAGTCACCTGGCAAAATCTGA
 GACATCTGAGTAGTATGAGCAATTCATTTCTGTAGAATGTCTACGAGAAAACAT
 5 AGCTTTTGAGTTGCCCCAAGAGTTTCTGCAATACACCCAACCTATGAAGAGGGAC
 ATCAAGAAGGCCTTCTATGAAATGTCCCTACAGGCCTTCAACATCTTCAGCCAAC
 ACACCTTCAAATATTGGAAAGAGAGACACCTCAAACAAATCCAAATAGGACTTGA
 TCAGCAAGCAGAGTACCTGAACCAATGCTTGGAGGAAGACGAGAATGAAAATGA
 AGACATGAAAGAAATGAAAGAGAATGAGATGAAACCCTCAGAAGCCAGGGTCCC
 10 CCAGCTGAGCAGCCTGGAACCTGAGGAGATATTTCCACAGGATAGACAATTTCTCTG
 AAAGAAAAGAAATACAGTGACTGTGCCTGGGAGATTGTCCGAGTGGAAATCAGA
 AGATGTTTGTATTACTTTTACAAATTTACAGCTCTATTACAGGAGGAAA (SEQ ID
 NO:119)

15 This base change causes the same corresponding change of one amino acid as found
 in Example 5 for the polypeptide coded by pCR2.1-cg Ba403c19-S546-1b (shown in
 underlined bold font in Table 56; SEQ ID NO:120) from the sequence for the ba403c19_A
 polypeptide shown in Table 16 (SEQ ID NO: 10).

20 **Table 56. Amino acid sequence of the polypeptide coded by the insert of
 pCR2.1-cg Ba403c19-S546-1b.**

LDCNLLNVHLRRVTWQNLRLHSSMSNSFPVECLRENIAFELPQEFLQYTQPMKRDIKK
 AFYEMSLQAFNIFSQHTFKYWKERHLKQIQIGLDQQA EYLNQCLEED**EN**ENEDMKEM
 KENEMKPSEARVPQLSSLELRRYFHRIDNFLKEKKYSDCAWEIVRVEIRRCLYYFYKF
 25 TALFRRK (SEQ ID NO:120)

**Example 7.: Expression of a mature form of clone BA403c19_A (NOV5) in human
 embryonic kidney 293 cells.**

The BamHI-XhoI fragment containing the BA403c19_A sequence was isolated from
 30 pCR2.1-cg Ba403c19-S546-1b (Example 6) and subcloned into BamHI-XhoI digested
 pCEP4/Sec (Example 2) to generate the expression vector pCEP4/Sec-BA403c19_A. The
 pCEP4/Sec-BA403c19_A vector was transfected into human embryonic kidney 293 cells
 (ATCC No. CRL-1573, Manassas, VA) using the LipofectaminePlus reagent following the
 manufacturer's instructions (Gibco/BRL/Life Technologies, Rockville, MD). The cell pellet
 35 and supernatant, respectively, were harvested 72 hours after transfection and examined for
 BA403c19_A expression by Western blotting (reducing conditions) with an anti-V5 antibody.

Fig. 2 shows that in the supernatant BA403c19_A is expressed as a polypeptide with an apparent molecular weight of about 35 kDa secreted by 293 cells, based on use of SeeBlue Molecular Weight Standards (Invitrogen).



Fig. 2. NOV5 (BA403c19_A) protein secreted by 293 cells.

OTHER EMBODIMENTS

While the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

What is claimed is:

1. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:
 - a) a mature form of the amino acid sequence selected from the group consisting of SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26;
 - b) a variant of a mature form of the amino acid sequence selected from the group consisting of SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26, wherein any amino acid in the mature form is changed to a different amino acid, provided that no more than 15% of the amino acid residues in the sequence of the mature form are so changed;
 - c) the amino acid sequence selected from the group consisting of SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26;
 - d) a variant of the amino acid sequence selected from the group consisting of SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26 wherein any amino acid specified in the chosen sequence is changed to a different amino acid, provided that no more than 15% of the amino acid residues in the sequence are so changed; and
 - e) a fragment of any of a) through d).
2. The polypeptide of claim 1 that is a naturally occurring allelic variant of the sequence selected from the group consisting of SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26.
3. The polypeptide of claim 2, wherein the variant is the translation of a single nucleotide polymorphism.
4. The polypeptide of claim 1 that is a variant polypeptide described therein, wherein any amino acid specified in the chosen sequence is changed to provide a conservative substitution.
5. An isolated nucleic acid molecule comprising a nucleic acid sequence encoding a polypeptide comprising an amino acid sequence selected from the group consisting of:

- a) a mature form of the amino acid sequence given SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26;
 - b) a variant of a mature form of the amino acid sequence selected from the group consisting of SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26 wherein any amino acid in the mature form of the chosen sequence is changed to a different amino acid, provided that no more than 15% of the amino acid residues in the sequence of the mature form are so changed;
 - c) the amino acid sequence selected from the group consisting of SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26;
 - d) a variant of the amino acid sequence selected from the group consisting of SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26, in which any amino acid specified in the chosen sequence is changed to a different amino acid, provided that no more than 15% of the amino acid residues in the sequence are so changed;
 - e) a nucleic acid fragment encoding at least a portion of a polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26 or any variant of said polypeptide wherein any amino acid of the chosen sequence is changed to a different amino acid, provided that no more than 10% of the amino acid residues in the sequence are so changed; and
 - f) the complement of any of said nucleic acid molecules.
6. The nucleic acid molecule of claim 5, wherein the nucleic acid molecule comprises the nucleotide sequence of a naturally occurring allelic nucleic acid variant.
7. The nucleic acid molecule of claim 5 that encodes a variant polypeptide, wherein the variant polypeptide has the polypeptide sequence of a naturally occurring polypeptide variant.
8. The nucleic acid molecule of claim 5, wherein the nucleic acid molecule comprises a single nucleotide polymorphism encoding said variant polypeptide.

9. The nucleic acid molecule of claim 5, wherein said nucleic acid molecule comprises a nucleotide sequence selected from the group consisting of
- a) the nucleotide sequence selected from the group consisting of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25;
 - b) a nucleotide sequence wherein one or more nucleotides in the nucleotide sequence selected from the group consisting of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25 is changed from that selected from the group consisting of the chosen sequence to a different nucleotide provided that no more than 15% of the nucleotides are so changed;
 - c) a nucleic acid fragment of the sequence selected from the group consisting of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25; and
 - d) a nucleic acid fragment wherein one or more nucleotides in the nucleotide sequence selected from the group consisting of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25 is changed from that selected from the group consisting of the chosen sequence to a different nucleotide provided that no more than 15% of the nucleotides are so changed.
10. The nucleic acid molecule of claim 5, wherein said nucleic acid molecule hybridizes under stringent conditions to the nucleotide sequence selected from the group consisting of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 or 25, or a complement of said nucleotide sequence.
11. The nucleic acid molecule of claim 5, wherein the nucleic acid molecule comprises a nucleotide sequence in which any nucleotide specified in the coding sequence of the chosen nucleotide sequence is changed from that selected from the group consisting of the chosen sequence to a different nucleotide provided that no more than 15% of the nucleotides in the chosen coding sequence are so changed, an isolated second polynucleotide that is a complement of the first polynucleotide, or a fragment of any of them.
12. A vector comprising the nucleic acid molecule of claim 11.
13. The vector of claim 12, further comprising a promoter operably linked to said nucleic acid molecule.

14. A cell comprising the vector of claim 12.
15. An antibody that binds immunospecifically to the polypeptide of claim 1.
16. The antibody of claim 15, wherein said antibody is a monoclonal antibody.
17. The antibody of claim 15, wherein the antibody is a humanized antibody.
18. A method for determining the presence or amount of the polypeptide of claim 1 in a sample, the method comprising:
 - (a) providing said sample;
 - (b) introducing said sample to an antibody that binds immunospecifically to the polypeptide; and
 - (c) determining the presence or amount of antibody bound to said polypeptide, thereby determining the presence or amount of polypeptide in said sample.
19. A method for determining the presence or amount of the nucleic acid molecule of claim 5 in a sample, the method comprising:
 - (a) providing said sample;
 - (b) introducing said sample to a probe that binds to said nucleic acid molecule; and
 - (c) determining the presence or amount of said probe bound to said nucleic acid molecule, thereby determining the presence or amount of the nucleic acid molecule in said sample.
20. A method of identifying an agent that binds to the polypeptide of claim 1, the method comprising:
 - (a) introducing said polypeptide to said agent; and
 - (b) determining whether said agent binds to said polypeptide.

21. A method for identifying a potential therapeutic agent for use in treatment of a pathology, wherein the pathology is related to aberrant expression or aberrant physiological interactions of the polypeptide of claim 1, the method comprising:
- (a) providing a cell expressing the polypeptide of claim 1 and having a property or function ascribable to the polypeptide;
 - (b) contacting the cell with a composition comprising a candidate substance; and
 - (c) determining whether the substance alters the property or function ascribable to the polypeptide;
- whereby, if an alteration observed in the presence of the substance is not observed when the cell is contacted with a composition devoid of the substance, the substance is identified as a potential therapeutic agent.
22. A method for modulating the activity of the polypeptide of claim 1, the method comprising introducing a cell sample expressing the polypeptide of said claim with a compound that binds to said polypeptide in an amount sufficient to modulate the activity of the polypeptide.
23. A method of treating or preventing a pathology associated with the polypeptide of claim 1, said method comprising administering the polypeptide of claim 1 to a subject in which such treatment or prevention is desired in an amount sufficient to treat or prevent said pathology in said subject.
24. The method of claim 23, wherein said subject is a human.
25. A method of treating or preventing a pathology associated with the polypeptide of claim 1, said method comprising administering to a subject in which such treatment or prevention is desired a NOVX nucleic acid in an amount sufficient to treat or prevent said pathology in said subject.
26. The method of claim 25, wherein said subject is a human.

27. A method of treating or preventing a pathology associated with the polypeptide of claim 1, said method comprising administering to a subject in which such treatment or prevention is desired a NOVX antibody in an amount sufficient to treat or prevent said pathology in said subject.
28. The method of claim 27, wherein the subject is a human.
29. A pharmaceutical composition comprising the polypeptide of claim 1 and a pharmaceutically acceptable carrier.
30. A pharmaceutical composition comprising the nucleic acid molecule of claim 5 and a pharmaceutically acceptable carrier.
31. A pharmaceutical composition comprising the antibody of claim 15 and a pharmaceutically acceptable carrier.
32. A kit comprising in one or more containers, the pharmaceutical composition of claim 29.
33. A kit comprising in one or more containers, the pharmaceutical composition of claim 30.
34. A kit comprising in one or more containers, the pharmaceutical composition of claim 31.
35. The use of a therapeutic in the manufacture of a medicament for treating a syndrome associated with a human disease, the disease selected from a pathology associated with the polypeptide of claim 1, wherein said therapeutic is the polypeptide of claim 1.
36. The use of a therapeutic in the manufacture of a medicament for treating a syndrome associated with a human disease, the disease selected from a pathology associated with the polypeptide of claim 1, wherein said therapeutic is a NOVX nucleic acid.

37. The use of a therapeutic in the manufacture of a medicament for treating a syndrome associated with a human disease, the disease selected from a pathology associated with the polypeptide of claim 1, wherein said therapeutic is a NOVX antibody.
38. A method for screening for a modulator of activity or of latency or predisposition to a pathology associated with the polypeptide of claim 1, said method comprising:
- a) administering a test compound to a test animal at increased risk for a pathology associated with the polypeptide of claim 1, wherein said test animal recombinantly expresses the polypeptide of claim 1;
 - b) measuring the activity of said polypeptide in said test animal after administering the compound of step (a); and
 - c) comparing the activity of said protein in said test animal with the activity of said polypeptide in a control animal not administered said polypeptide, wherein a change in the activity of said polypeptide in said test animal relative to said control animal indicates the test compound is a modulator of latency of, or predisposition to, a pathology associated with the polypeptide of claim 1.
39. The method of claim 38, wherein said test animal is a recombinant test animal that expresses a test protein transgene or expresses said transgene under the control of a promoter at an increased level relative to a wild-type test animal, and wherein said promoter is not the native gene promoter of said transgene.
40. A method for determining the presence of or predisposition to a disease associated with altered levels of the polypeptide of claim 1 in a first mammalian subject, the method comprising:
- a) measuring the level of expression of the polypeptide in a sample from the first mammalian subject; and
 - b) comparing the amount of said polypeptide in the sample of step (a) to the amount of the polypeptide present in a control sample from a second mammalian subject known not to have, or not to be predisposed to, said disease,
- wherein an alteration in the expression level of the polypeptide in the first subject as compared to the control sample indicates the presence of or predisposition to said disease.

41. A method for determining the presence of or predisposition to a disease associated with altered levels of the nucleic acid molecule of claim 5 in a first mammalian subject, the method comprising:
- a) measuring the amount of the nucleic acid in a sample from the first mammalian subject; and
 - b) comparing the amount of said nucleic acid in the sample of step (a) to the amount of the nucleic acid present in a control sample from a second mammalian subject known not to have or not be predisposed to, the disease;
- wherein an alteration in the level of the nucleic acid in the first subject as compared to the control sample indicates the presence of or predisposition to the disease.
42. A method of treating a pathological state in a mammal, the method comprising administering to the mammal a polypeptide in an amount that is sufficient to alleviate the pathological state, wherein the polypeptide is a polypeptide having an amino acid sequence at least 95% identical to a polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26 or a biologically active fragment thereof.
43. A method of treating a pathological state in a mammal, the method comprising administering to the mammal the antibody of claim 15 in an amount sufficient to alleviate the pathological state.